

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

aSF757
.2
.A77
2003

States
Department of
Agriculture

Agricultural
Research
Service

Animal Welfare
Information Center

March 2005



2003 ARS Immunology Research Workshop Proceedings

National Agricultural Library Cataloging Record:

ARS Immunology Research Workshop (2003 : Bethesda, Maryland)

2003 ARS Immunology Research Workshop proceedings.

1. Animals—Immunology—Research—Congresses. 2. Natural immunity—Congresses.


3. Immunologic diseases in animals—Treatment--Congresses.

4. Animals—Diseases—Nutritional aspects—Congresses.

I. Allen, Tim (John Timothy), 1953- II. Callahan, Rosemary.

III. United States. Agricultural Research Service. IV. Title.


SF757.2



2003 ARS Immunology Research Workshop Proceedings

A workshop sponsored by the U.S. Department of Agriculture, Agricultural Research Service at Bethesda, Maryland, on December 1-4, 2003.

Also includes a bibliography of representative ARS immunology research published since 2000.



Compiled and edited by:

Tim Allen
United States Department of Agriculture
Agricultural Research Service
National Agricultural Library
Animal Welfare Information Center
Beltsville, Maryland

Rosemary Callahan
United States Department of Agriculture
Agricultural Research Service
National Program Staff
Beltsville, Maryland

Abstract

Allen, Tim, Callahan, Rosemary, eds. 2005. 2003 ARS Immunology Research Workshop Proceedings. U.S. Department of Agriculture, Agricultural Research Service, and National Agricultural Library, Animal Welfare Information Center.

The information in this proceedings provides an insight into ongoing and recently published animal and human immunology research performed by the U.S. Department of Agriculture's Agricultural Research Service (ARS). It also contains final reports produced by conference work groups that will provide a roadmap for future ARS immunology research efforts. The work groups discussed four broad topics: 1) immunity and disease resistance; 2) immunointervention strategies; 3) immunity and the pathogenesis of disease; and 4) immunity and nutrition. The purpose of the workshop was to examine the state of current research in animal immunology and determine where to concentrate future efforts to advance national research programs in animal health, animal production, aquaculture, food safety, and human nutrition.

Keywords: immunity and disease resistance, immunointervention strategies, immunity and the pathogenesis of disease, immunity and nutrition.

Mention of trade names or commercial products in this report is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

To ensure timely distribution, this report was reproduced essentially as supplied by the authors. It received no publication editing and design.

While supplies last, single copies of this publication may be obtained at no cost from Tim Allen, U.S. Department of Agriculture, National Agricultural Library, Animal Welfare Information Center, 10301 Baltimore Avenue, 4th Floor, Beltsville, MD 20705.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotope, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Issued March 2005

Table of Contents

Executive Summary	1
Welcome Letter	4
Agenda.....	5
Keynote Addresses	11
Jay A. Berzofsky	11
National Cancer Institute, NIH, Bethesda, Maryland	11
Vaccine strategies to improve CTL avidity, memory, and efficacy	11
Arthur M. Krieg	12
Coley Pharmaceutical Group, Wellesley, Massachusetts	12
Enhancing vaccines with adjuvants that activate dendritic cells and B cells in vivo	12
Alan Sher ¹	13
C. Feng ² , C. Collazo ¹ , and G.A. Taylor ³	13
¹ Laboratory of Parasitic Diseases, NIAID, NIH, Bethesda, Maryland	13
The p47 GTPases: a family of IFN induced proteins that regulate host resistance to intracellular pathogens	13
Plenary Sessions	14
Innate Immunity.....	14
Immune Regulation	17
Immunogenetics.....	21
Immunity and Nutrition	24
Immunity and Disease Resistance	28
Immunointervention.....	31
Poster Sessions	35
Innate Immunity.....	35
Immune Regulation	38
Immunogenetics.....	43
Immunity and the Pathogenesis of Disease	44
Immunity and Nutrition	54
Immunity and Disease Resistance	60
Immunointervention Strategies.....	63
Working Group Reports.....	69
Immunity and Disease Resistance	70
Immunointervention.....	75
Immunity and the Pathogenesis of Diseases	81
Nutrition and Immune Function Interactions.....	83
Auburn, Alabama	88
Aquatic Animal Health Research.....	88
Fayetteville, Arkansas	93
Poultry Production and Products Safety Research	93
Little Rock, Arkansas.....	96
Arkansas Children's Nutrition Center	96
Davis, California	97
Western Human Nutrition Research Center	97

Athens, Georgia	100
Southeast Poultry Research Laboratory.....	100
West Lafayette, Indiana.....	105
Livestock Behavior Research Unit	105
Crop Production and Pest Control Research	105
Ames, Iowa	106
National Animal Disease Center	106
New Orleans, Louisiana	123
Southern Regional Research Center.....	123
Beltsville, Maryland	125
Henry A. Wallace Beltsville Agricultural Research Center.....	125
Boston, Massachusetts	140
The Jean Mayer USDA Human Nutrition Research Center on Aging	140
East Lansing, Michigan	142
The Avian Disease and Oncology Laboratory	142
Mississippi State, Mississippi	147
Stoneville, Mississippi	147
Columbia, Missouri.....	148
Animal Physiology Research Unit	148
Donald Danforth Plant Science Center, Plant Genetics Research Unit.....	149
Clay Center, Nebraska	149
Roman L. Hruska U.S. Meat Animal Research Center.....	149
Orient Point, New York.....	150
Plum Island Animal Disease Center	150
College Station, Texas	153
Southern Plains Agricultural Research Center	153
Houston, Texas	156
Children's Nutrition Research Center	156
Logan, Utah.....	157
Poisonous Plant Research	157
Pullman, Washington	158
Animal Disease Research Unit.....	158
Kearneysville, West Virginia	161
National Center for Cool and Cold Water Aquaculture	161
Laramie, Wyoming	162
Arthropod-Borne Animal Diseases Research Laboratory.....	162
APPENDIX	164
List of Participants	165
Author Index.....	179

2003 ARS Immunology Research Workshop

Executive Summary

The 2003 ARS Immunology Research Workshop was sponsored by the United States Department of Agriculture (USDA), Agricultural Research Service (ARS), in Bethesda, Maryland, December 1-4, 2003. The purpose of the workshop was to exchange recent scientific breakthroughs in animal immunology and determine where ARS and its partners should concentrate future efforts to significantly advance national research programs in animal health, animal production, aquaculture, food safety, and human nutrition.

The workshop participants included ARS immunologists working in livestock, wildlife, avian and marine species, as well as immunologists working in other Federal laboratories and universities across the United States. The first day of the workshop provided a series of presentations covering high profile topics in basic immunology research with emphasis on cutting-edge discoveries in innate immunology, immune regulation, and immunogenetics. The second day focused on the applications of immunology research in nutrition, disease resistance, and immunointervention. This exchange of information set the stage for the third day, when all participants were divided into the following four working groups: 1) immunity and disease resistance; 2) immunointervention strategies; 3) immunity and the pathogenesis of diseases; and 4) immunity and nutrition. The challenge set forth for the working groups was to identify research objectives within specific areas of animal immunology that could significantly advance research programs in animal health, animal production, aquaculture, food safety, and human nutrition.

The disease resistance working group was asked to identify research objectives that will lead to the discovery and characterization of immune mechanisms associated with heritable disease resistant traits and that may in parallel be used as targets to design novel biotherapeutics and highly efficacious vaccines. The immunointervention working group was tasked with providing research objectives that will significantly increase our understanding of protective immunity and lead to the rational design of highly efficacious vaccines and diagnostics. The goal for the pathogenesis working group was to provide research priorities to advance our understanding of immune responses that lead to pathogenesis, either inherently programmed within the immune system, or driven by metabolic and infectious disease pathways. The human nutrition working group was asked to identify research objectives in animal immunology that were most likely to make significant contributions to our understanding of health and nutrition.

A significant outcome of the workshop was that all four working groups identified the lack of veterinary immunological reagents as a key stumbling block to advancing the field of animal immunology. The lack of immunological reagents was also considered paramount to make significant progress in many of our national research programs. For instance, the lack of efficient tools for evaluating cell-mediated immunity in cattle and swine makes the discovery and development of highly efficacious vaccines more difficult to attain and demonstrate. Many of the workshop participants felt that cell-mediated immunity should be at the core of strategies to discover new tools against the diseases that resist traditional vaccine approaches. The overwhelming consensus was that the void in our immunological tool chest is the rate-limiting step in providing the next generation of vaccines.

In today's economy it is highly unlikely that the private sector will invest in the basic research necessary to develop the tools for testing cell-mediated immunity. Accordingly, 26 participants agreed to look at organizing a community wide effort, post-workshop, to discover, develop, and distribute quality-controlled immunological reagents for veterinary science research.

As a result of the final working sessions, ARS immunologists and their partners identified a total of 66 research objectives to incorporate into future research action plans. The research objectives with the highest priority are noted here:

- Establish national and international networks to prioritize and develop immune toolkits for animal species of economic importance. Coordinate tool development so that priority reagents become available. Work with commercial sources to market products.
- Identify critical components of innate immunity; Use innate immune pathways for interventions/traditional and therapeutic vaccines and for development of new adjuvants and improvement of vaccine efficacy.
- Identify and characterize host genes that control important immunologic responses; determine whether alleles of certain genes would enhance animal health
- Use information from microbial genomics to define substances from pathogens that stimulate/regulate immune responses at mucosal surfaces.
- Develop broad spectrum, targeted delivery systems (target dendritic cells or others) for immunomodulator and/or peptide and/or DNA delivery to effector cells.
- Optimize use of existing vaccines and immunomodulators for intervention at embryogenesis, birth, weaning, or prior to stress (shipping, comingling, slaughter).
- Elucidate factors affecting efficacy of current adjuvant systems
- Define the affects of stress/diet/age on immune responses at mucosal surfaces.
- Identify specific nutrients with beneficial immune modulating properties that optimize immune function. Define specific nutrient requirements for effective and appropriate vaccination and immunological memory, and for amelioration of periods of immune deficiency, such as during pregnancy, parturition, pediatric development, aging, and stress.
- Define probiotic microbial populations that affect immune development and functional resistance at mucosal surfaces and on the efficacy of systemic immunity, and the diets (prebiotics) that support appropriate resident or ingested microbial populations.
- Develop new methods for detecting and measuring cell-mediated immunity; identify microbial antigens and regulatory events critical for the development and maintenance of cell-mediated immunity.
- Utilize genomic information to reveal new approaches for disease control or biotherapeutic production and identify novel nutrient and immune interactions.
- Define gene expression patterns and regulatory phenomena that affect the level of host resistance at mucosal surfaces.

In addition, the workshop participants selected the following research objectives for their potential to substantially impact ARS national research programs in animal health, animal production, aquaculture, food safety, and human nutrition:

- Novel approaches for disease diagnostics; Biomarkers of immunointervention success will enhance our capacity to screen for the most efficacious method to elicit a protective immune response.

- Improved methods to limit immune-associated disease and persistence of pathogens thereby increasing production and decrease spread of disease, respectively.
- Improved methods to limit factors (e.g., environmental, husbandry, nutritional) that negatively impact neonatal immune development.
- Definition of pathways for interventions, for development of both traditional and therapeutic vaccines, and for preparation of novel mucosally targeted biotherapeutics
- Increased understanding of how innate immunity impacts vaccine responses; improved knowledge base for adjuvant and immune modulator design/selection for vaccines and biotherapeutics.
- Improved vaccine efficacy; verification of efficacy of mucosal delivery vectors and adjuvants; targeting the neonate and maternal immunity.
- Broad-based vaccine delivery systems applicable to both production animals and wildlife that account for the practical realities of immunization in the field.
- Provide identification of diets that enhance immune development and reduce inflammation.
- Better understanding of the interaction between dietary choices and environment on immune function and optimal health.
- The emergence of genetic profiles will include dietary recommendations for human life style and animal management to optimize health.
- By leveraging knowledge in databases from human disease research, we will build a comparative immunology database that can be a guide to antigen selection for use in vaccines.

Welcome Letter



United States Department of Agriculture
Agricultural Research Service
National Program Staff

December 1, 2003

Dear Workshop Attendee:

On behalf of the National Program Staff, welcome to the "2003 ARS Immunology Research Workshop." We are delighted that you could join us for this exciting event. The scientists participating in the workshop include Immunologists working in very different fields and we hope the diversity of disease models, animal species, and approaches to solving problems discussed over the next three days will be beneficial to your research. We also hope you will have many opportunities to establish strategic collaborations over the next three days that will build on our understanding of Immunology and its many applications to animal and human health.

We have four goals for the workshop:

1. To exchange new information and ideas in immunology to maximize the impact of our research programs and solve problems of high National priority.
2. To enhance interactions and form strategic collaborations between scientists engaged in immunology research.
3. To provide a forum for stakeholders, customers, and partners to give input and direction on current and future ARS immunology research programs.
4. To initiate the preparation of action plans for consideration at future ARS National Program workshops.

The workshop has been structured to facilitate interactions and the exchange of new ideas between scientists conducting research in Federal laboratories, academia, and the private sector. Working group sessions will be held on the last day to assess future research direction and prepare research plans. We are very fortunate to have some of the most prominent experts in immunology working in animal and human health participate in this workshop. We have tried to format this workshop to promote constructive exchange of the most recent data and new ideas.

Once again, we are delighted that you could join us for this exciting event and we thank you for your participation in this workshop.

Sincerely,

Cyril Gerard Gay, D.V.M., Ph.D
National Program Leader, Animal Health and Safety



Animal Production, Product Value, and Safety
5601 Sunnyside Avenue • Beltsville, MD 20705-5138

2003 ARS Immunology Research Workshop

Agenda

December 1-4, 2003

Bethesda, Maryland

December 1, 2003

Kick Off

18:00 – Welcome Address, Cyril Gerard Gay, National Program Leader, Animal Health and Safety, ARS, Beltsville, Maryland

Goals, Partnerships, and Opportunities

18:30 - Keynote Presentation, Jay Berzofsky, NCI/NIH/HHS, Bethesda, Maryland

Vaccine Strategies to Maximize CTL Avidity, Memory, and Efficacy

19:30 - Reception

December 2, 2003

Plenary Sessions

Innate Immunity

Moderator: James A. Roth, Director, Center for Food Security and Public Health, College of Veterinary Medicine, Iowa State University, Ames, Iowa

08:00 - Matthew Fenton, University of Maryland School of Medicine, Baltimore, Maryland

Role of Toll-like Receptors in the Host Response Against Mycobacterial Infection

08:30 - Susan D. Eicher, ARS, Livestock Behavior Research Unit, West Lafayette, Indiana

Toll-like Receptor and Acute Phase Cytokine Expression in Neonatal Dairy Calves

09:00 - Rami Dalloul, ARS, Animal Parasite Diseases Laboratory, Beltsville, Maryland

Oligodeoxynucleotides (ODN)-induced Activation of Macrophage-Mediated Innate Immunity Against Intracellular Pathogens, *Eimeria* And *Salmonella*, and the Development Of Novel Immunomodulatory Strategies Using Embryo Injection of Cpg-Odns in Poultry.

09:30 - Elida M. Bautista, ARS, Foot and Mouth Disease Research, Orient Point, New York

Analysis of Swine Skin-Derived Dendritic Cells in the Induction of Innate Immune Responses to Foot-and-Mouth Disease Virus (FMDV)

10:00 - Interactive Poster Session (38 posters)

- Innate Immunity
- Immune Regulation
- Immunogenetics
- Immunity and the Pathogenesis of Diseases

Immune Regulation

Moderator: William Golde, Immunologist, Plum Island Animal Research Center, ARS, Orient Point, New York

11:00 - Willi Born, National Jewish Medical and Research Center, Denver, Colorado

Regulatory Role of Gamma/Delta T Cells in Allergic Airway Hypereactivity

11:30 - David Woodland, Trudeau Institute, Saranac Lake, New York

Regulation of Memory T Cell Responses in Viral Infections of the Respiratory Tract

12:00 - Lunch and Keynote Presentation: Arthur Kreig, Coley Pharmaceuticals, Wellesley, Massachusetts

Enhancing Vaccines with Adjuvants that Activate Dendritic Cells and B Cells In Vivo

13:30 - Wendy Brown, Washington State University, Pullman, Washington

Regulation of the CD4⁺ T Lymphocyte Response by *Anaplasma marginale*

14:00 - Wenbin Tuo, ARS, Animal Parasite Diseases Laboratory, Beltsville, Maryland

Bovine T Cell Response to Antigens of *Neospora caninum*

14:30 - Marc Estes, University of Texas, Galveston, Texas

Progress Towards Developing Effective Subunit Vaccine Responses by CD8⁺ Memory and Effector T Cells in a Bovine Model of Tuberculosis

15:00 – Jesse P. Goff, ARS, Periparturient Diseases of Cattle Research Unit, Ames, Iowa

Effects of the Mammary Gland on the Composition and Function of Peripheral Blood Mononuclear Leukocyte Populations in Periparturient Dairy Cows

15:30 – Break

Immunogenetics

Moderator: Peter L. Nara, Director of Research and Development, Biological Mimetics, Inc., Frederick, Maryland

16:00 - Annie DeGroot, Epivax, Inc, Providence, Rhode Island

Accelerating Vaccine Design: New In Silico Epitope Mapping Methods

16:30 - Henry Hunt, ARS, Avian Diseases and Oncology Laboratory, East Lansing, Michigan
Marek's Disease and the MHC: How Viruses Subvert the Immune System for Their Advantage

17:00 - Lynn M. Herrmann, ARS, Animal Disease Research Unit, Pullman, Washington
Expression of Specific MHC Class II DRB1 Alleles Associates with Sheep Persistently Infected with Ovine Progressive Pneumonia Virus (OPPV)

17:30 - John Butler, University of Iowa, Iowa
From Basic Porcine Immunology to Practical Application

December 3, 2003

Plenary Sessions

Immunity and Nutrition

Moderator: Ildy Katona, Department of Pediatrics, Uniformed Services University of the Health Sciences in Bethesda, Maryland

07:30 - Joe Urban, ARS, Nutrient Requirements and Functions Laboratory, Beltsville, Maryland
The Effect of Human-Derived Probiotic Bacteria on the Immune and Intestinal Function of Pigs

08:00 - Kirk Klasing, University of California, Davis, California
The Care and Feeding of an Immune System: How Much is Needed and What are the Priorities?

08:30 - Simin Meydani, ARS, Nutritional Immunology Laboratory, Boston, Massachusetts
Aging, Immune Response, and Infectious Disease: Molecular Mechanisms and Reversal by Nutrient Intervention

09:00 - Brian Nonnecke, Periparturient Diseases of Cattle Research Unit, Ames, Iowa
Nutritional Plane Affects Antigen-Induced Proliferation of Lymphocyte Subsets in Milk Replacer-Fed Calves Vaccinated with *M. bovis* Bacillus Calmette-Guerin (BCG)

09:30 - Harry Dawson, Nutrient Requirements and Functions Laboratory, Beltsville, Maryland
Real-Time PCR Arrays can Delineate Immunological and Nutrition-Related Gene Expression in Swine

10:00 - Interactive Poster Session (32 posters)

- Immunity and nutrition
- Immunity and disease resistance
- Immunointervention strategies

Immunity and Disease Resistance

Moderator: Muquarrab Qureshi, National Program Leader, Animal Genomics, CSREES, USDA, Washington D.C.

11:00 – Michael Murtaugh, University of Minnesota,
Mechanisms of Innate and Adaptive Immunity in Swine

11:30 - Lou Gasbarre, ARS, Bovine Functional Genomics, Beltsville, Maryland
Genetic and Immunologic Basis for Resistance to GI Nematode Infections

12:00 - Lunch and Keynote Presentation: Alan Sher, NIAID/NIH/HHS, Bethesda, Maryland
The P47 Gtpases: A Family of IFN Induced Proteins that Regulate Host Resistance to Intracellular Pathogens

13:30 - Pamela Ferro, ARS, Food and Feed Safety Research, College Station, Texas
Heterophils Isolated from Chickens Resistant to Extraintestinal *Salmonella enteritidis* Infection Express Higher Levels of Pro-inflammatory Cytokine mRNA Following Infection than Heterophils from Susceptible Chickens

14:00 - Hyun Lillehoj, ARS, Animal Parasite Diseases Laboratory, Beltsville, Maryland
High-Throughput Molecular Approaches to Identify Coccidiosis Disease Resistance Genes and Protective Immunological Mechanisms in Avian Coccidiosis

Immunointervention

Moderator: Ronald Schultz, Professor and Chair, Department of Pathobiological Sciences, University of Wisconsin, Madison, Wisconsin

14:30 – Cynthia Baldwin, University of Massachusetts,
Protective Killed *Leptospira* Vaccine Induces Potent Type 1 (Th1) Immunity Comprising Responses by CD4 and WC1+ $\gamma\delta$ T Lymphocytes

15:00 – Evelyn Dean-Nystrom, ARS, Food Safety and Enteric Diseases Research Unit, Ames, Iowa
Development of an Oral Vaccine for *Escherichia coli* O157:H7

15:30 - Break

16:00 - Wongi Min, ARS, Animal Parasite Diseases Laboratory, Beltsville, Maryland
Adjuvant Effect of IL-1 β , IL-2, IL-8, IL-15, IFN- α , IFN- γ , TGF- β 4 and Lymphotoxin on DNA Vaccination Against *Eimeria acervulina*

16:30 - Haiqi He, ARS, Food and Feed Safety Research, College Station, Texas

Stimulation of Chicken Leukocytes and Reduction of *Salmonella* Organ Invasion in Neonatal Chicken by Immunostimulatory Cpg-Oligodeoxynucleotide

17:00 – M. Suresh, Assistant Professor of Immunology, University of Wisconsin, Madison, Wisconsin
Regulation of Memory CD8 T Cell Generation by Cytokines

December 4, 2003

Research Priorities Working Groups. *The workshop participants will be divided into teams and asked to identify research priorities, strategic collaborations, and innovative research plans that will substantially increase our knowledge of immunology and deliver solutions to disease problems of high national priority in animal and human health. The recommendations of the participants will form the basis for developing ARS National Program action plans. Focus will be given to the following research areas:*

- Immunity and disease resistance
- Immunointervention strategies
- Immunity and the Pathogenesis of Diseases
- Nutrition and Immune Function Interactions

08:30 Working Group Instructions

- Robert Heckert, National Program Leader, Animal Health, ARS, Beltsville, Maryland
- Lewis Smith, National Program Leader, Aquaculture & Animal Well-Being, ARS, Beltsville, Maryland

09:00 Working Group Sessions

- **Immunity and Disease Resistance**

Working Group Leaders:

- Lanie Bilodeau, Immunologists, Catfish Genetics Research Unit, ARS, Stoneville, Mississippi
- Lou Gasbarre, Research Leader, Bovine Functional Genomics Research Unit, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- Michael H. Kogut, Immunologist, Southern Plains Agricultural Research Center, ARS, College Station, Texas
- Joan Lunney, Immunologist, Animal Parasite Diseases Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland

- **Immunointervention Strategies**

Working Group Leaders:

- Marcus Kehrli, Research Leader, Virus and Prion Diseases Research Laboratory, National Animal Disease Center, ARS, Ames, Iowa
- Hyun Lillehoy, Immunologist, Animal Parasite Diseases Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- David White, Veterinary Medical Officer, Arthropod-borne Animal Diseases Research Laboratory, ARS, Laramie, Wyoming

- **Immunity and the Pathogenesis of Diseases**

Working Group Leaders:

- Don Knowles, Research Leader, Animal Disease Research Unit, ARS, Pullman, Washington
- David Swayne, Research Leader, Southeast Poultry Research Laboratory, ARS, Athens, Georgia
- Ray Waters, Veterinary Medical Officer, Bacterial Diseases of Livestock Research Unit, National Animal Disease Center, ARS, Ames, Iowa

- **Nutrition and Immune Function Interactions**

Working Group Leaders:

- Theodore Elsasser, Research Animal Scientist, Growth Biology Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- Simin Meydani, Chief, Nutritional Immunology Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging, ARS, Tufts University, Boston, Massachusetts
- Joe Urban, Research Leader, Nutrient Requirements and Functions Laboratory, Beltsville Human Nutrition Research Center, ARS, Beltsville, Maryland

12:00 **Lunch**

13:00 **Working Group Reports**

- Immunity and Disease Resistance
- Immunointervention Strategies
- Immunity and the Pathogenesis of Diseases
- Nutrition and Immune Function Interaction

15:00 - **Conclusion and Post-meeting Instructions**

- Cyril Gerard Gay, National Program Leader, Animal Health and Safety, ARS, Beltsville, Maryland

Keynote Addresses

December 1, 2003

Jay A. Berzofsky

Molecular Immunogenetics & Vaccine Research Section, Metabolism Branch,
National Cancer Institute, NIH, Bethesda, Maryland

Vaccine strategies to improve CTL avidity, memory, and efficacy

Cytokines are potentially useful not only as vaccine components to increase the magnitude of the immune response, but also to steer it in desired directions. We found that certain combinations such as GM-CSF and IL-12 synergized to increase cytotoxic T lymphocyte (CTL) activity. We showed that GM-CSF worked to increase antigen presenting cell numbers and activity in the draining lymph nodes. We reasoned that CD40L might synergize with GM-CSF by maturing those presenting cells. Indeed, GM-CSF and CD40L were a potent combination to increase CTL responses. Also, we had seen that NKT cells making IL-13 dampened the CTL response to tumors, so we asked if blockade of IL-13 or elimination of NKT cells would also increase the CTL response to a peptide vaccine. We found that either of these treatments further enhanced the CTL response and the ability to protect against viral challenge. Thus, a push-pull approach in which a cytokine and a costimulatory molecule are combined to push the response and disruption of an inhibitory pathway is used to pull the response can maximize CTL induction.

In addition to increasing the quantity of the CTL, we sought to also improve the quality of the CTL induced by a vaccine. We previously found that avidity of CTL is critical for effective

viral clearance, but selective induction of high avidity CTL has been difficult to achieve with vaccines. Likewise, persistent CTL memory is essential for long-lasting CTL-mediated vaccine efficacy. Thus, we have sought strategies to selectively induce longer-lived high avidity CTL with vaccines targeting HIV protein antigens. First, increasing costimulation (signal 2) allowed us to immunize with lower doses of antigen (signal 1) to selectively induce high avidity CTL in vivo. This was achieved using a recombinant poxvirus vector expressing a triad of costimulatory molecules (B7-1, ICAM-1, and LFA-3) (TRICOM) to increase expression of these molecules on splenocytes pulsed with an HIV-1 CTL epitope peptide. Second, since IL-15 was known to contribute to maintenance of CTL memory, we asked whether IL-15 at the time of immunization would alter the CTL response. Expression of IL-15 by a recombinant vaccinia vector along with HIV-1 gp160 induced long-lived memory CTL that persisted at least 14 months after a single boost. The memory cells expressed higher levels of IL-15R α and showed greater proliferation to IL-15 in vitro and homeostatic proliferation in vivo, as well as higher avidity for antigen. Thus, IL-15 at the time of priming selects or induces a different phenotype CTL with greater avidity and longevity. These vaccine strategies can provide more effective CTL immunity by enhancing the numbers, avidity, and memory of the CTL induced, allowing for greater and longer lasting protection.

December 2, 2003

Arthur M. Krieg

Coley Pharmaceutical Group, Wellesley, Massachusetts

Enhancing vaccines with adjuvants that activate dendritic cells and B cells in vivo

The innate immune system has evolved “pattern recognition receptors” that detect molecules present in various classes of pathogens, but not in our own cells. The best known of these pattern recognition receptors belong to the family of Toll-like receptors (TLRs), which appear to have evolved to activate appropriate protective immune responses when a pathogen is detected. One such pathogen-associated pattern is unmethylated CpG dinucleotides in bacterial or viral DNA, which are detected by TLR9, thereby stimulating innate and adaptive immunity. These immune effects can be mimicked by synthetic oligodeoxynucleotides containing CpG motifs (CpG ODN). In humans and most other species, at least two types of immune cells express TLR9 and are activated directly by CpG ODN; B cells, and plasmacytoid dendritic cells (pDC). CpG ODN activate B cells to proliferate and secrete immunoglobulin; and pDC to secrete a variety of Th1-like cytokines, chemokines, and type I interferons, and to express increased costimulatory molecules. When activated by CpG, pDC gain the ability to stimulate Th1-like T cell responses. CpG ODN

induce immune defense mechanisms that protect against challenge with a wide range of infectious pathogens, and have shown therapeutic activity in animal models of allergic disease and cancer. CpG ODN are also extremely effective vaccine adjuvants, inducing Th1 responses in mice, primates, and other species with both specific antibody and CTL. Phase I human clinical trial results indicate that a CpG ODN, 7909, is relatively well tolerated. As an adjuvant for a hepatitis B vaccine, CpG 7909 appears to induce earlier seroconversion with the production of increased levels of specific antibody and cellular responses. A phase II human clinical trial of CpG 7909 in HIV-infected subjects, including prior hepatitis B vaccine nonresponders, showed a striking increase in the magnitude of antibody responses in the subjects receiving CpG as a vaccine adjuvant. CpG ODN are produced by cGMP chemical synthesis, have excellent stability and solubility, and show an excellent safety profile, suggesting wide utility as vaccine adjuvants for veterinary applications.

December 3, 2003

Alan Sher¹

C. Feng², C. Collazo¹, and G.A. Taylor³

¹Laboratory of Parasitic Diseases, NIAID, NIH, Bethesda, Maryland

²Center for Biologics Evaluation and Research, FDA, Bethesda, Maryland

³Dept. of Medicine and Immunology, Duke University Medical Center, Durham, North Carolina

The p47 GTPases: a family of IFN induced proteins that regulate host resistance to intracellular pathogens

Activation of the immune system by interferon- γ is critical for host resistance to many infectious agents. This lymphokine induces the expression of a wide range of effectors that undermine the ability of pathogens to survive in host cells. Among these is a newly discovered family of 47 kDa GTPases that provide resistance to intracellular protozoa, bacteria, and viruses. Elimination of different p47 GTPases in mice through gene targeting severely cripples IFN- γ -

regulated defense against *Toxoplasma gondii*, *Listeria monocytogenes*, *Mycobacterium sp.*, and other pathogens. Interestingly, different members of this gene family appear to effect resistance to different groups of pathogens. Current evidence suggests that p47 GTPases function against pathogens at two different levels: by facilitating intracellular killing of pathogens and by promoting survival of effector T lymphocytes.

Plenary Sessions

December 2, 2003

Innate Immunity

Role of toll-like receptors in the host response against mycobacterial infection

Matthew J. Fenton

Director of Research, Division of Pulmonary and Critical Care Medicine; Professor of Medicine, Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, Maryland

Toll-like receptor (TLR) proteins mediate cellular activation by microbes and microbial products. To delineate the role of TLR proteins in the development of host immune responses against mycobacteria, wild-type and TLR-deficient mice were infected with non-pathogenic *Mycobacterium bovis* BCG (BCG). Two weeks after intraperitoneal challenge with BCG, few bacilli were present in the lungs of wild-type and TLR4^{-/-} mice, whereas bacterial loads were 10-fold higher in the lungs of infected TLR2^{-/-} mice. BCG challenge in vitro strongly induced pro-inflammatory cytokine secretion by macrophages from wild-type and TLR4^{-/-} mice, but not by TLR2^{-/-} macrophages. In contrast, intracellular uptake, intracellular bacterial growth, and suppression of intracellular bacterial growth in vitro by IFN- γ were similar in macrophages from all three mouse strains, suggesting that BCG growth in the lungs of TLR2^{-/-} mice was a consequence of defective adaptive immunity. Antigenic stimulation of splenocytes from infected wild-type and TLR4^{-/-} mice induced T cell proliferation in vitro, whereas T cells from TLR2^{-/-} mice failed to proliferate. This lack of T cell proliferation in TLR2^{-/-} mice correlated with an inability of CD11c⁺ dendritic cells to be activated in vivo following BCG challenge. Unexpectedly, activated CD4⁺ T cells from both TLR-deficient mouse strains secreted little IFN- γ in vitro,

compared with control T cells. A role for TLR4 in the control of bacterial growth and IFN- γ production in vivo was observed only when mice were infected with higher numbers of BCG. Thus, TLR2 and TLR4 appear to regulate distinct aspects of the host immune response against BCG.

Toll-like receptor and acute phase cytokine expression in neonatal dairy calves

S.D. Eicher¹, T.A. Johnson², K.A. McMunn¹, and T.R. Johnson²

¹ USDA-ARS, Livestock Behavior Research Unit, West Lafayette, Indiana; ²Purdue University, West Lafayette, Indiana

The well-being of farm animals is dependent on many factors, so that a multi-disciplinary approach is essential. The Livestock Behavior Research Unit uses behavior, physiology, and immunology measures to answer well-being questions for poultry, swine, and dairy cattle. The innate immune system is frequently the first immune responder during many stressors and acute phase cytokines affect behavior and learning. Toll-like receptors are the pathogen recognition molecules that initiate the cascade that increases the acute phase cytokines. The effect of stressors on the toll-like receptors of farm animals is not known. Studies have been completed that examine the effect of stimulants on RNA expression of toll-like receptors in young dairy calves. The objective of study one was to assess the optimum concentration and times of incubation for expression of TLR2 and TLR4 after in-vitro stimulation of bovine monocyte derived macrophages with *Saccharomyces cerevisiae* (yeast cell-wall) beta-glucan (BG), *Escherichia coli* lipopolysaccharide (LPS), and *Staphylococcus aureus* peptidoglycan (PGN). Monocyte derived

macrophages from 10 donor calves (5 to 6 wk-of-age) were first stimulated with LPS or PGN at 1, 5, or 25 µg/ml or BG at 20, 40, or 80 µg/ml for one hour at 37°C. Then the optimum concentrations were used to incubate cells at 30, 60, or 90 min. Real-time RT-PCR was used to quantify RNA expression of TLR2 and TLR4. RNA from monocyte derived macrophage expressed more bovine TLR4 than TLR2. Sixty minutes appears to be an adequate time of stimulation to change TLR4 expression, but at least 90 minutes are needed to attain maximum changes in TLR2 expression. The objective of study two was to determine if in vitro stimulation of whole blood leukocytes with beta-glucan plus ascorbic acid (a potential immunomodulator) was dependent on the age of the calf. Blood samples were taken from 12 non-transported Holstein calves at 1, 3, 7, 10, 14, 18, 21, 24, and 28 d-of-age. Leukocytes were stimulated with ascorbic acid (0.3 µg/ml) plus beta-glucan (0.4 µg/ml) or media only for 1 h and then RNA was extracted. The RNA was subjected to real-time RT-PCR for quantification of the expression of interleukin-1 (IL-1) and its receptor antagonist (IL-1Ra), and toll-like receptors 2 and 4 (TLR2 and TLR4). TLR2 and TLR4 had treatment effects ($P<.05$), but not day or treatment by day interactions. TLR2 was greater ($P<.05$) for treated cells on day 7, 14, and 24 and tended ($P<.10$) to be greater on day 10. In contrast, TLR4 was only greater for treated cells on day 7 ($P<.05$) and tended to be ($P<.10$) on day 24. IL-1 had a treatment main effect and a treatment by day interaction ($P<.05$), but IL-1Ra had main effects for treatment and day ($P<.05$), but only a trend ($P<.10$) for a treatment by day interaction. IL-1 was greater for treated cells ($P<.05$) on all but day 4. IL-1Ra was greater ($P<.05$) for treated cells only on days 1, 7, 10, and 24. Only IL-1 and its receptor antagonist expression were stimulated on day 1. On days 7 and 24 all tested receptors and cytokines had increased RNA expression. So, it appears that there are periods during which the blood leukocytes may be refractory for increased RNA expression of cytokines and toll-like receptors in response to beta-glucan and ascorbic acid stimulus. These data enhance our understanding of the biological kinetics of the developing neonatal calf immune system, so that

management procedures can occur at the best possible times.

Oligodeoxynucleotides (ODN)-induced activation of macrophage-mediated innate immunity against intracellular pathogens, *Eimeria* and *Salmonella*, and the development of novel immunomodulatory strategies using embryo injection of CpG-ODNs in poultry.

Rami Dalloul^{1,2}, Hyun S. Lillehoj¹, Hang Xie², Robert A Heckert², Uma Babu³, Richard Raybourne³, and Dennis M. Klinman⁴

¹ USDA, ARS, Animal Parasitic Diseases Laboratory, Beltsville, Maryland; ² Virginia-Maryland Regional College of Veterinary Medicine, University of Maryland, College Park, Maryland; ³ Center for Food Safety & Applied Nutrition, US FDA, Laurel, Maryland; ⁴ Section of Retroviral Immunology, Center for Biologics Evaluation and Research, US FDA, Bethesda, Maryland

The immunostimulatory properties of synthetic CpG oligodeoxynucleotides (ODNs) have been well studied in various mammalian models including humans and mice. However, little was known about effects of CpG-ODNs on immune responses of chickens, a common avian species with important economical value in the poultry industry. ODN 2006 which has been reported to be an optimal stimulatory sequence for humans, showed strong immunomodulatory effects on HD11 cells. Whereas ODN 1826, a murine optimal CpG-ODN had weak influences on HD11 cells. We also observed that 2006 induced strong IL-6 and nitric oxide secretion by HD11 cells in both dose- and time-dependent manners. 2006 induced activation also resulted in increased intracellular killing of *S. enteritidis* (SE) by HD11 cells. Furthermore, 2006 activated HD11 cells had reduced proliferation and underwent apoptosis after CpG stimulation. Further studies were carried out to test several short synthetic CpG ODNs for their immunomodulatory effects on disease susceptibility in *Eimeria acervulina*. On day 18 of incubation, specific pathogen-free embryos were injected (air cell) with either 25 µg

or 50 µg of CpG ODN (CpG-1, CpG-2, CpG-3 and CpG-4), a non CpG-containing control (CpG-0), or a negative control (PBS, phosphate buffered saline). At one week of age, birds were orally inoculated with 10^4 *E. acervulina* oocysts. Body weights and body weight gains were not affected by any of the treatments. However, among CpG ODN treatments, a differential antibody response was observed with the lowest response in CpG-2 birds and highest in CpG-4 group. This study demonstrates that CpG ODN were effective immunoprotective agents in chickens, and the potentials of CpG-ODNs as immunomodulators in poultry need to be further explored.

Analysis of swine skin-derived dendritic cells in the induction of innate immune responses to Foot-and-mouth disease virus (FMDV)

Elida M. Bautista, Geoffrey S. Ferman, Douglas Gregg, Mario C. Brum, Marvin Grubman, and William T. Golde.

USDA, ARS, Plum Island Animal Disease Center, Greenport, New York

To understand the role of dendritic cells (DC) in the initiation of immune responses against FMDV and to determine whether FMDV infection may alter this function, we analyzed the interaction of FMDV with skin-derived DC of swine. Innate responses of DC to in vitro FMDV exposure with infectious or attenuated leaderless (LL) viruses was determined by measuring cytokine mRNA

and protein expression using Real-time PCR analysis and ELISA, respectively. Activation state of these DC was determined by measuring surface expression of SLAII (MHC Class II) and CD80/86. SLAII was minimally affected and CD80/CD86 slightly reduced in DC exposed to either virus. DC consistently responded to both viruses with a rapid increase in IFN-beta and TNF-alpha mRNA expression that peaked at 4-8 h post-exposure. IFN mRNA was only moderately increased over the constitutive levels. Analysis of supernatants (SN) of virus exposed DC show secretion of IFN and IFN by protein assay (ELISA), but neither protein was detected in the control culture SN with no virus. Biological function of IFN was confirmed only in the virus-exposed cell culture SN by antiviral assay. Consistent with the antiviral effect of IFNs, no productive infection by FMDV was detected in DC. Immunocytochemistry and confocal microscopy of skin sections and freshly isolated DC demonstrated these cells constitutively express IFN. These results show that DC can mount an innate type I IFN response to FMDV, which renders them resistant to productive viral replication. This study also revealed an important property of porcine skin DC in that these cells constitutively express and accumulate IFN protein in the cytoplasm, and respond to virus exposure by secreting IFN and by activating the IFN gene and secreting this cytokine as well. Together, our data show that skin DC have an important role in the innate immune response of swine to FMDV.

Immune Regulation

Regulatory role of gamma/delta T cells in allergic airway hyperreactivity

Willi Born

National Jewish Center, Denver, Colorado.

Airway hyperresponsiveness to cholinergic agonists (AHR) is a life-threatening condition associated with many diseases of the vertebrate lung. Gamma/delta T cells exert a strong regulatory effect on AHR. We have used a mouse model of allergic airway disease to investigate the underlying mechanisms. Features include regulation during antigen priming and challenge, systemically and within the lung, and small populations of gamma/delta T cells with opposed regulatory influences on AHR. As regulatory functions co-segregate with T cell receptor Vgamma expression, specific ligand recognition may play a role, and ligands may be expressed on dendritic cells (DC). Gamma/delta T cells appear to monitor DC in the lung, and early signals from DC might induce their regulatory effect on the developing antigen-specific immune response and airway smooth muscle contraction.

The contribution of memory T cells subsets to antiviral immunity in the lung

David L. Woodland

Trudeau Institute, Saranac Lake, New York

Respiratory virus infections elicit memory CD4⁺ and CD8⁺ T cell populations that persist for the life of the individual and are able to mediate accelerated clearance of secondary virus infections. Given the protective capacity of these cells, there has been substantial interest in developing vaccines that promote cellular immunity in the lung. However, we have little understanding of how T cell memory is established, maintained, and recalled in peripheral organs. To begin to address these issues, we have been analyzing memory T cell populations elicited by influenza and parainfluenza virus infections in a mouse model. We show that memory T cells are extremely heterogeneous in terms of their

anatomical distribution, phenotype, function, and longevity. Moreover, distinct subpopulations vary greatly in their capacity to respond to infection, depending on their initial phenotype and anatomical location. These observations have significant implications for the development of vaccines designed to promote cellular immunity in the lung.

Regulation of the CD4⁺ T lymphocyte response by *Anaplasma marginale*.

Wendy C. Brown, Guy H. Palmer, and Jeffrey R. Abbott

Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, Washington

MSP2 is serologically immunodominant for individuals infected with *Anaplasma*. In cattle protected by immunization with *A. marginale* outer membranes, CD4⁺ T cell and IgG responses are also predominantly directed against MSP2 and MSP3, which share the conserved carboxy region. In fact, MSP2-specific T cells also recognize MSP3. Antigenic variation in MSP2 occurs throughout the lifespan of persistently infected but clinically healthy cattle, and sequentially emerging variants are controlled by the immune response. Both variant-specific CD4⁺ T cells and antibody are believed to be important for this control. We hypothesized that immunization with multiple MSP2 variants should provide protection against challenge with organisms expressing the same MSP2 variants. To test this, eight calves were immunized with native MSP2 using either IL-12 or CpG DNA adjuvants that stimulated IgG and type 1 T cell responses, and together with four control animals, were challenged five months later with homologous organisms that expressed the same major MSP2 variants. In spite of developing significant and strong MSP2-specific CD4⁺ T cell proliferative, IFN-γ, and IgG responses, vaccinates were not protected from challenge. Importantly, acute infection was not due to the emergence of novel MSP2 variants. To elucidate the reasons for

the lack of protection in the vaccinates, we compared antigen-specific responses of CD4⁺ T-lymphocytes from MSP2 vaccinates and control cattle cryopreserved before and after *A. marginale* challenge using IFN- γ ELISPOT and proliferation assays. Reproducibly strong MSP2-specific PBMC responses were detected from immunized animals prior to challenge and for two weeks after challenge, which was prior to peak rickettsemia. Surprisingly, MSP2-specific responses of PBMC collected during the peak of rickettsemia were undetectable in all animals. This severely depressed MSP2-specific T cell response was still observed two-four months later, and was also apparent when using *A. marginale* homogenate. In contrast, the response to unrelated *Clostridium* vaccine antigens was not ablated during acute infection. By four to seven months after peak rickettsemia, MSP2-specific T cell responses were detected in several vaccinates; however, these remained significantly depressed. Our data suggest that in addition to immune evasion via antigenic variation in surface proteins, *A. marginale* induced unresponsiveness may facilitate organism persistence. Potential mechanisms include induction of regulatory T cells, induction of T cell anergy, and exhaustion of MSP2-specific T cells. Supported by USDA NRI grants 99-35204-8368 and 02-35204-12352 and NIH NIAID awards K08-AI49276 and R01-AI44005.

Bovine T cell response to antigens of *Neospora caninum*.

Wenbin Tuo¹, Ray Fetterer², Pat Boyd¹, Lou Gasbarre³, and J.P. Dubey².

¹ USDA, ARS, Immunology and Disease Resistance Lab, Beltsville, Maryland; ²Animal Parasitic Diseases Lab, Beltsville, Maryland; ³Bovine Functional Genomics Lab, ANRI, Beltsville, Maryland

Neospora caninum is an apicomplexan parasite that infects a number of cell types, causing neosporosis in several animal species including cattle. In the past decade, neosporosis has been recognized as a primary cause of abortion in cattle worldwide. It was estimated that approximately 20% of all bovine abortions are attributed to neosporosis; *Neospora*-related abortion costs the

cattle industry in excess of \$35 million a year in the state of California alone. In infected cows, *N. caninum* elicits a detectable antibody (Ab) response as shown by extensive studies, whereas *N. caninum*-induced cell-mediated immunity (CMI) is poorly understood. Previous exposure to, or infection by, the parasite does not provide protection against abortion or vertical transmission, in spite of the presence of *Neospora*-specific Ab in maternal circulation. The objective of this study was to define *Neospora*-specific CMI in cows infected with *N. caninum* and to identify immunodominant T helper cell Ag that may be used as vaccine candidates. Total *Neospora* Ag was prepared by freezing and thawing or sonication of *N. caninum* tachyzoites grown in a monkey kidney or a bovine macrophage cell line. Peripheral blood lymphocytes (PBL) from *Neospora* infected cows were stimulated with total soluble *Neospora* Ag weekly in the presence of irradiated PBL as a source of Ag presenting cells. Flow cytometry analysis showed 90-98% of these *Neospora* Ag specific T cells (week 3-8) were CD2⁺CD3⁺CD4⁺. A T cell proliferation assay was established and used to determine Ag specificity of Ag-specific T helper cells and antigenic activity of *Neospora* Ag. A bovine specific IFN- γ ELISA was used to determine IFN- γ content in supernatants of T cells stimulated with *Neospora* Ag. The total *Neospora* tachyzoite Ag was separated into 5 fractions by anion exchange using HPLC. Results indicate that high levels of IFN- γ were produced by both PBL and T cell lines specific to *Neospora* Ag. Using the T cell proliferation assay, we showed that proteins from HPLC Fractions 1 and 2 had low or undetectable T cell-stimulating activity, whereas proteins from Fractions 4-6 had higher stimulatory activity. However, Fraction 5 contained the highest antigenic activity in all T cell lines assayed. Western blot analysis showed that Fraction 5 contained 8-10 distinctive *Neospora* protein bands that were defined by a rabbit antiserum raised against to *Neospora* tachyzoites (NC-1 strain). These studies suggest that Ag-specific T helper cells (CD4⁺ CD3⁺) may be used to screen for immunodominant *Neospora* Ag. *Neospora* proteins that specifically stimulate T helper cell proliferation and cytokine production identified

using our T cell proliferation and cytokine production assays may be used as vaccine candidates. This research will enhance our understanding of bovine CMI to *Neospora* infection and may facilitate the development of a vaccine against bovine neosporosis.

Progress towards developing effective subunit vaccine responses by CD8+ memory and effector T cells in a bovine model of tuberculosis.

D. Mark Estes

University of Texas Medical Branch, Department of Pediatrics and Pathology, The Sealy Center for Vaccine Development, Galveston, Texas

Cell-mediated immunity (CMI) is a key component of the protective immune response for many intracellular pathogens (viruses, bacteria and parasites). CD8+ cytotoxic T cells (hereafter referred to as CTL) are a primary effector cell type in the CMI response to intracellular pathogens via their ability to lyse infected target cells and produce key regulatory and effector cytokines. CD8+ T cells recognize relatively short peptides (8-11 amino acids in length) presented in the context of MHC class I molecules on the surface of an antigen presenting cell (APC). APC processing of polypeptides from both inside and outside of the cell is the first step in the formation of MHC class I-peptide complexes and the generation of T cell responses. Rational design of vaccines for intracellular pathogens requires knowledge of both this process and the means by which costimulatory signals are delivered to naïve T cells by APC. Costimulatory signals are required in addition to T cell receptor (TCR) engagement to drive the response. The requirements for accessory molecule costimulation (collectively, signal 2) in the activation of naïve CD8+ T cells, in addition to the primary signal (signal 1) generated via the T cell receptor-antigen-MHC complex, are not well understood, particularly in cattle. Knowledge of this important issue is essential to the development of viable vaccine strategies that involve elicitation of effective CD8+ T cell responses.

Effects of the mammary gland on the composition and function of peripheral blood mononuclear leukocyte populations in periparturient dairy cows.

J.P. Goff, K. Kimura, and B.J. Nonnecke

USDA, ARS, Periparturient Disease of Cattle Research Unit, National Animal Disease Center, Ames, Iowa

During the periparturient period, the dairy cow experiences a state of natural immunosuppression that is associated with heightened susceptibility to infectious disease. During this period, alterations in T cell percentages in the circulation parallel a reduction in functional capacities of peripheral blood mononuclear cell (PBMC) population. Hormonal and metabolic demands of pregnancy and lactation as well as the physiologic stress of calving likely contribute to periparturient immunosuppression. Objectives of this study were to evaluate the phenotype and functional capacities of PBMC populations in periparturient cows, and secondly, determine whether changes in composition and function could be, in part, attributable to the presence of a functional mammary gland. Ten mastectomized and eight intact multiparous Jersey cows (six for each for functional study) were used. Using monoclonal antibodies against T-cell subsets, the B-cell, and the monocyte, the composition of PBMC populations was characterized by flow cytometry. For the analyses, cells were obtained several times a week from -4 wk prepartum to 4 wk postpartum. Functionality of PBMC populations was assessed by assays measuring mitogen-induced lymphocyte proliferation, IgM secretion, and IFN- γ secretion. Cells used in these assays were collected twice weekly from -3 wk prepartum to 3 wk postpartum. Percentages of CD3+, CD4+, CD8+, and $\gamma\delta$ TCR+ cells in intact cows decreased progressively with the approach of parturition, while monocyte percentages increased. Synthesis of DNA and secretion of IFN- γ and IgM by mitogen-stimulated intact-cow PBMC were lowest around the time of calving. In contrast, compositional and functional changes in PBMC populations from mastectomized cows were minimal or absent at these times. Regression analysis revealed significant associations between

composition and functional capacity of the PBMC population. These results indicate that the mammary gland and metabolic stresses associated with lactation contribute to the generalized immunosuppression affecting dairy cows during the periparturient period.

Immunogenetics

Accelerating vaccine design: new in silico epitope mapping methods

Annie De Groot

CEO, EpiVax, Associate Professor, Brown Medical School, Providence, Rhode Island

T cell responses play an important role in immunity to parasites and other microbial agents of infectious diseases, therefore a number of T cell-directed vaccines are in development. Computer-driven algorithms that facilitate the discovery of T cell epitopes from protein and genome sequences are now being used to accelerate the development of human vaccines. Similar tools are now becoming available for animal vaccines. My group recently developed new in silico methods for predicting epitopes for cattle and swine; the development process and validation studies will be described in this talk.

Epitope-mapping algorithms offer a significant advantage over other methods of epitope selection, such as the screening of synthetic overlapping peptides, because high throughput screening can be performed in silico, followed by ex vivo confirmatory studies. Using epitope-mapping algorithms, putative T cell epitopes can be derived directly from genomic sequences, allowing researchers to circumvent labor-intensive cloning steps in the genome-to-vaccine discovery pathway.

Marek's disease and the MHC: How viruses subvert the immune system for their advantage.

H. Hunt, H. Cheng, and H. Zhang
USDA, ARS, Avian Diseases and Oncology Laboratory, East Lansing, Michigan

Immune response genes of the chicken play a critical role in the resistance or susceptibility to Marek's disease (MD), a T-cell tumor induced by a herpesvirus named MD virus (MDV). The major histocompatibility complex (MHC) is one component of the immune response that has a

major influence on the outcome of MD as well as the immune protection afforded by MDV vaccines. The USDA-ARS Avian Disease and Oncology Laboratory (ADOL) in East Lansing, MI is investigating the influence of the chicken MHC and other genetic systems involved in the immune response to MD. ADOL's approach is multidisciplinary incorporating traditional genetics, functional genomics/proteomic and immunologic analysis to dissect the host-pathogen interactions responsible for resistance or susceptibility to MDV.

The traditional genetics approach involves the generation of recombinant congenic chicken strains (RCS). ADOL White Leghorn lines 6 and 7 are resistant or susceptible to MDV, respectively, and differ in immune response characteristics and lymphoid organ size. The two lines were intermated and full-sib mated to generate 19 inbred sub-lines designated 6C.7A through X. Each 6C.7 sub-line consists of approximately 12% of the susceptible line 7 genome inserted into the resistant line 6 genomic background. The immune response, disease resistance and genetic phenotypes of these 6C.7 RCS are currently under investigation at ADOL using gene mapping and immune assay approaches.

MDV, like other herpesviruses, has the ability to alter the expression of antigen presenting glycoproteins encoded by the MHC. MDV inhibits the cell surface expression of MHC class I glycoproteins but, unique to MDV, induces the expression of MHC class II glycoproteins. ADOL's multidisciplinary approach incorporating traditional immunologic assays coupled with genomics and proteomics is beginning to unravel the host pathogen interactions responsible for these observations.

Expression of specific MHC Class II DRB1 alleles associates with sheep persistently infected with ovine progressive pneumonia virus (OPPV).

L.M. Herrmann¹, W.P. Cheevers², and D.P. Knowles^{1,2}

¹USDA, ARS, Animal Disease Research Unit, Pullman, Washington; ² Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, Washington

During OPPV infection in a flock, there is a clear dichotomy where a few sheep produce clinical signs and most become persistently infected without clinical signs. In long-term persistent infections, cell-adaptive immunity plays the significant role in controlling virus replication. In other small ruminant lentiviruses, CD4 positive and not CD8 positive T lymphocytes are required for adequate proliferation upon exposure to viral antigens (Perry et al., 1995). This along with two other studies indicating that the immunogenetics of the host dictate the course of infection and the antibody response to OPPV (de la Concha-Bermejillo et al., 1995; Cheevers et al., 1999), support the idea that MHC Class II immunogenetics play a significant role in directing disease outcome. Since ovine MHC Class II DRB1 encodes and expresses the antigen presentation domain on the antigen-presenting cell, we evaluated the allelic expression of DRB1 in peripheral blood mononuclear cells (PBMC) from 10 sheep persistently infected with OPPV. Five of these sheep have high OPPV loads, and five have low OPPV loads by real time PCR of integrated DNA in PBMC. The relationship between viral load and specific DRB1 allele expression will be discussed. In addition, the relationship of viral genetics in terms of the surface envelope glycoprotein (SU) and specific DRB1 allele expression will also be discussed.

From basic porcine immunology to practical application

John E. Butler

University of Iowa, Iowa City, Iowa

Characterizing the elements that comprises the

immune system is the first step needed to provide the information and tools for ultimately determining the role of the immune system in health and disease. Described here is a 12-year journey of discovery in a farm animal with a pattern of passive immunity and a precocial nature that makes it an ideal model for providing insight into immunological development, receptor repertoires and the immunopathology of an important viral disease.

Basic investigations revealed the swine immune system is anything but a carbon copy of the human and mouse systems. For example, swine rely on only 10 combinatorial rearrangements, not 9,000, to develop their heavy chain antibody repertoire. Equally efficient is the development of the V-kappa repertoire that primarily uses one V kappa gene family and one J kappa gene segment. Also in contrast to mice and humans, early B cell development is adult-like, and surprisingly involves the thymus beginning at midgestation, where switch recombination is pronounced *in utero* in the absence of environmental antigen. Naïve B cells migrate to the ileal Peyer's patches, each follicle containing the diversified descendents of a single B cell clone.

While B cell development occurs rapidly in late gestation, the ability of the newborn to mount an immune response requires exposure to colonizing bacteria or infective virus that produce the ligands for the Toll-like receptors (TLRs).

While the mechanism of antibody repertoire development in swine distinguishes it from rodents and primates, T-cell development follows a conserved evolutionary pattern that parallels that in other mammals. The TCR beta repertoire strongly resembles that of humans and mice in both genome organization and the presence of many homologous V-beta families. The V-epsilon repertoire is also conserved and an invariant V epsilon transcript occurs early in fetal life reminiscent of that in mice.

We were able to use the tools of discovery and knowledge we had gained to employ the isolator piglet model to show that PRRSV directly causes

a B cell lymphoproliferative disorder characterized by immune dysfunction, autoimmunity, immune complex deposition and kidney and vascular damage. We hope as the

journey continues, we shall soon show how PRRSV alters the T cells repertoire and how specific bacterial- and viral-derived TLR ligands allow the adaptive immune system to develop.

December 3, 2003

Immunity and Nutrition

The effect of human-derived probiotic bacteria on the immune and intestinal function of pigs

Gloria Solano-Aguilar*¹, Harry Dawson¹, Tonya Ledbetter¹, Terez Shea-Donohue¹, Norberta Schoene¹, Jeff Call², Ethiopia Beshah³, William Hare³, and Joseph Urban¹

USDA, ARS, ¹Beltsville Human Nutrition Research Center, Beltsville, Maryland; ² Eastern Regional Research Center, Wyndmoor, Pennsylvania; ³Animal and Natural Resources Institute, Beltsville, Maryland

Probiotics are specific microorganisms that can establish and grow in a compartment of an exposed host and provide some positive health benefit; strains of *Lactobacilli*, *Bifidobacteria* and *Enterococcus* are commonly used probiotics. The benefits claimed vary from protection against pathogenic microorganisms and viruses, stimulation of the intestinal immune function and enhanced disease resistance. However, the basis of these claims is often confounded by a lack of demonstrable growth and function of the probiotic strain in the gut. Our objective was to test if dietary probiotics can establish in the pig and affect immunity to an infectious agent.

Experiment 1:

A pilot study using eight 12-week-old pigs was performed to determine if animals fed with a daily dose of 10 billion colony forming units (cfu) of freeze-dried *Lactobacillus rhamnosus* GG (LGG; Culturelle, Conagra, NE) would show some change in their immune response. Whole blood for isolation of peripheral blood mononuclear cells (PBMC) and fecal samples for DNA extraction were collected throughout the experiment at days 4, 7, 12, 18 and 25 after initiation of dietary treatment. At the end of the experiment (day 26) pigs were euthanized and tissues from different mucosal sites were collected for: 1) cell immune

phenotype analysis by flow cytometry; 2) DNA isolation and identification of LGG strain in intestinal mucosa and fecal samples, 3) measurement of IFN- γ production in vitro after mitogen stimulation, and 4) mRNA expression of a panel of immune markers by real-time PCR. PBMC isolated from LGG-exposed pigs produced IFN- γ in response to Con A stimulation, while IFN- γ was barely detectable (assay range in ng/ml) in culture supernatant of PBMC isolated from unexposed pigs. LGG-specific fluorogenic probes were able to detect LGG DNA in fecal samples of treated animals as early as 4 days post-treatment. Mucosal tissue samples and fecal contents taken from intestinal sites of LGG treated animals indicated that LGG DNA was detected at 4-fold higher levels in the proximal colon compared to distal colon and cecum. No signal was detected in jejunum or ileum of any animals or in samples from non-treated animals. Real time PCR analysis indicated a selective low level increase in gene expression for Th1-derived type 1 immune cytokines in localized mucosal sites after probiotic delivery.

Experiment 2:

To test the effect of early colonization of probiotic bacteria on the immune response to a nematode parasite infection, *Bifidobacterium lactis* (Bb12)(Mak Wood Inc, WI) (10×10^{10} cfu) was administered to 2 sows starting at the last third of pregnancy through weaning of their piglets at day 28; two other sows and their litters were left as untreated controls. After birth, Bb12 capsules were also administered to pigs born to treated sows. Pigs born from untreated sows were maintained as untreated controls. All piglets were inoculated with *Ascaris suum* eggs at 6 weeks of age and tissue samples were collected for functional gut in Ussing chambers 3 weeks later. Probiotic treatment did not alter normal absorption of glucose in the small intestine, but did attenuate

A. suum-induced inhibition of glucose absorption that correlates with expulsion of the worm from the pig intestine.

Conclusions:

Human derived probiotics can modulate immune function and selectively affect local responses to parasitic infection while promoting swine health. This model can be extended to assess the activity of selected probiotics on pig responses to other infectious agents and allergens that negatively affect pig production.

The care and feeding of an immune system: How much is needed and what are the priorities?

Kirk C. Klasing

Department of Animal Science, University of California, Davis California

How big is the immune system and how good is it at competing for scarce nutrients? Immunologists have neglected to measure the size of the immune system. Nutritionists have neglected to determine the nutritional requirements of the processes that we call 'immunity'. Currently, these first principles of nutrition and immunity can only be estimated from a synthesis of experiments that were designed for other reasons. We need to know: the costs of developing an immune system that is sufficiently robust to be dependably protective; the costs of maintaining this system of leukocytes in large enough copy number to be effective; and the cost of using this system when challenged the host is challenged by a proficient pathogen. A preliminary analysis of these costs suggests that development of innate immunity is cheap but development of adaptive systems (lymphocytes) is relatively expensive. The reverse is true for maintenance. Following a challenge, the production of immunoglobulin and the clonal expansion needed for protective T-cell responses are cheap to generate but can cause immunopathology in defending against some pathogens. The systemic sequelae of an acute phase response are especially expensive and markedly impact the nutritional state and productivity of an animal. The types of nutrient transporters utilized by leukocytes determine their

priorities for nutrients. Macrophages have a high priority, B-lymphocytes are intermediate, and T-lymphocytes are inept.

Aging, immune response, and infectious disease: molecular mechanisms and reversal by nutrient intervention.

Simin Nikbin Meydani, Dayong Wu, Sung Nim Han, and Raina T. Gay

USDA, ARS, Nutritional Immunology Laboratory, Jean Mayer Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts

The incidence of neoplastic and infectious diseases is increased in the elderly, as is the resulting morbidity and mortality. The well-documented age-associated dysregulation of immune function is an important contributor to the increased incidence of mortality from these diseases. Accumulating evidence indicates that T cell-mediated immune responses decline with aging. We, as well as others, have demonstrated that in addition to intrinsic changes in T cells, increased macrophage (M ϕ)-derived prostaglandin E₂ (PGE₂) production contributes to the age-associated decline in T cell function [1]. We have previously shown that vitamin E supplementation of aged mice and elderly humans significantly improves T cell mediated functions including their response to vaccination [2-4]. Vitamin E exerted its immunostimulatory effect by two distinct mechanisms: 1) by reducing M ϕ PGE₂ production, and 2) by a direct effect on T cell function. We further showed that vitamin E enhances T cell function directly by enhancing the cell dividing and IL-2 producing capacity of naïve T cells from old mice, with no effect on memory T cells [5].

To determine the clinical significance of vitamin E-induced immune enhancement in the aged we investigated the effect of vitamin E supplementation on viral and bacterial infection in the aged. Vitamin E supplementation of aged mice significantly reduced lung viral titer in mice infected with influenza virus.[6, 7]. On the other hand, old mice fed a diet supplemented with vitamin E (500 ppm) compared to those fed a diet

adequate in vitamin E, vitamin E supplementation did not protect against primary pulmonary *Staphylococcus aureus* (*S. aureus*) infection, whereas it was protective against secondary pulmonary *S. aureus* infection following influenza infection. This was indicated by a significantly lower bacterial count in the lungs of vitamin E-supplemented mice compared to that in control mice infected with *S. aureus* following influenza infection. In a recently completed double-blind, and placebo controlled study, we determined the effect of vitamin E on respiratory infection in nursing home residents. The results of this study as well as others will be discussed.

This work was supported in part by Federal funds from USDA, Agriculture Research Service under contract number 58-1950-9-00, and from the National Institute on Aging, grant AG 09140-09.

References:

1. Beharka, A.A., et al. 1997. Macrophage PGE₂ production contributes to the age-associated decrease in T cell function which is reversed by dietary antioxidants. *Mechanisms of Ageing and Development* 93: 59-77.
2. Meydani, S.N., et al. 1986. Vitamin E supplementation suppresses prostaglandine E₂ synthesis and enhances the immune response of aged mice. *Mechanisms of Ageing and Development* 34: 191-201.
3. Meydani, S.N., et al. 1990. Vitamin E supplementation enhances cell-mediated immunity in healthy elderly subjects. *American Journal of Clinical Nutrition* 52: 557-563.
4. Meydani, S.N., et al. 1997. Vitamin E supplementation and in vivo immune response in healthy elderly subjects. A randomized controlled trial. *JAMA* 277: 1380-1386.
5. Adolfsson, O., B.T. Huber, and S.N. Meydani. 2001. Vitamin E-enhanced IL-2 production in old mice: naive but not memory T cells show increased cell division cycling and IL-2-producing capacity. *Journal of Immunology* 167(7): 3809-17.
6. Han, S.N. and S.N. Meydani. 2000. Antioxidants, cytokines, and influenza infection in aged mice and elderly humans. *Journal of Infectious Disease* 182: 574-581.
7. Hayek, M.G., et al. 1997. Vitamin E supplementation decreases lung virus titers in mice infected with influenza. *Journal of Infectious Disease* 176: 273-276.

Nutritional plane affects antigen-induced proliferation of lymphocyte subsets in milk replacer-fed calves vaccinated with *M. bovis* bacillus Calmette-Guerin (BCG).

B.J. Nonnecke¹, M.R. Foote², M.A. Fowler³, B.L. Miller³, and W.R. Waters⁴

¹ USDA, ARS, Periparturient Diseases of Cattle Research Unit, National Animal Disease Center, Ames, Iowa; ²Iowa State University, Ames, Iowa;

³Land O' Lakes, Inc., Webster City, Iowa;

⁴USDA, ARS, Bacterial Diseases of Livestock Research Unit, NADC, Ames, Iowa

National Research Council recommended nutrient requirements for the neonatal dairy calf do not support optimal growth, and may compromise immune function. In a preliminary study, dietary energy and protein were shown to modulate in vitro production of interferon-gamma and nitric oxide by polyclonally stimulated blood lymphocytes from milk replacer-fed calves. We also have shown that adaptive (i.e. Th1-dependent) immune responses of milk replacer-fed neonatal calves vaccinated with *Mycobacterium bovis* bacillus Calmette Guerin (BCG) and challenged with purified protein derivative (PPD) are comparable in magnitude to adult responses. Using this sensitization/challenge model, effects of nutritional plane on antigen-induced proliferation of T cells from milk replacer-fed calves were evaluated. Holstein bull calves were fed a standard (ST, 0.57 kg/d of a 22% crude

protein, 20 % fat, n=4) milk replacer or an intensified (INT, 1.14 kg/d of a 28% crude protein, 20% fat milk replacer, n=4) milk replacer for 7 wk. Calves and four juvenile steers (5-6 m of age) were vaccinated with *M. bovis* BCG at the initiation of the study. During the experimental period, average daily weight gain was greater for INT than ST calves (0.68 and 0.32 kg/d, respectively). Proliferation of CD4⁺, CD8⁺, and $\gamma\delta$ TCR⁺ T cell subsets in PPD- and mitogen [pokeweed mitogen (PWM) and concanavalin A (ConA)]-stimulated cultures of peripheral blood mononuclear cells (PBMC) acquired before and 7 wk after vaccination were evaluated using flow cytometry. Recall responses to antigen were absent in all treatment groups prior to vaccination. Seven weeks after vaccination, PPD-induced proliferation of CD4⁺ and $\gamma\delta$ TCR⁺ cells from vaccinated calves exceeded responses of T cell subsets from vaccinated steers. In addition, antigen-induced proliferative responses CD8⁺ cells from INT calves exceeded those of ST calves. Lymphocytes from young adults were generally more responsive to mitogenic (i.e. ConA and PWM) stimulation than were cells from neonates. Overall, these results suggest that the proliferation of T cell subsets essential in CMI is affected by nutritional plane during the neonatal period.

Real-time PCR arrays can delineate immunological and nutrition-related gene expression in swine.

Harry Dawson

USDA, ARS, Nutrient Requirements & Functions Laboratory, Beltsville Human Nutrition Research Center, Beltsville, Maryland

Real-time PCR is commonly used to verify micro array expression data. We have designed over 300 human and swine real-time PCR assays for large-scale primary analysis of genes involved in immunological or nutritional function. These assays have a greater linearity, specificity and sensitivity when compared to micro array. The immunological foci of these assays are on pathways involved in apoptosis, toll receptor signaling and Th1/Th2 differentiation. The nutritional foci are on genes involved in iron, selenium, vitamin A and vitamin D metabolism. In conjunction with these assays, we have developed an annotated database for data mining using micro array analysis software packages. Using subsets of these assays, we profiled pig responses to *Toxoplasma gondii*, *Ascaris suum*, *Oesophagostomum dentatum*, *Salmonella typhimurium*, and *Campylobacter jejuni*. *Toxoplasma* elicited a generalized Th1 response typified by high-level induction of IFN- γ , TNF- α and iNOS. *Trichuris* induced a limited Th2-associated response including localized increased IgG, IgG1, IgG4, IL-13, IL-13Ra2, SOCS3, and CHIA expression. Resistant animals had higher GATA-3, IgE, IL-10, CD3e, CD40L, CD86, and MCP-2 expression, and lower ARG1 and MUC5AC expression. The impact of this tool lies in the expanding access to genetic information between mammalian species allowing hypothesis testing of infection resistance mechanisms using quantitative expression analysis of large numbers of genes at multiple tissue sites. There is also potential application to the measurement of therapeutic efficacy for a variety of immunological and nutritional interventions against human and swine diseases. Ongoing work includes studies with vitamin A and its derivatives.

Immunity and Disease Resistance

Mechanisms of innate and adaptive immunity in swine

Michael Murtaugh, Zhengguo Xiao, and Mark Rutherford

University of Minnesota, College of Veterinary Medicine, Veterinary Pathobiology, St. Paul, Minnesota

Mouse models of immune mechanisms of infectious disease resistance may be unsuitable for livestock species. The porcine immune system, for example, differs in immune tissue architecture, T cell phenotypes, cytokine expression patterns, cytokine activities, and possibly gene expression profiles. Thus, genomic, molecular, cell biologic, and other direct approaches will be required to elucidate porcine protective immune responses to infectious agents. Studies of immunity to porcine reproductive and respiratory syndrome virus (PRRSV), currently the most important pathogen of swine worldwide, illustrates the current limitations in the field. PRRSV exclusively infects macrophages of swine and causes a prolonged acute and persistent infection of tissues rich in macrophages. Detailed analysis of nonspecific innate and specific acquired immunity to PRRSV is beginning to reveal heretofore unsuspected mechanisms for control and elimination of viral infection in animals. PRRSV appears to subvert host defense by infection of lung macrophages without provoking the characteristic innate interferon α/β response to RNA viruses. The appearance of neutralizing antibodies does not occur until about three weeks after infection; acute infection manifested as viremia lasts for 4-5 weeks, and infectious virions are detected in lymphoid tissues for 3-4 months thereafter. While infection lingers for a prolonged time, viral load in lymphoid tissues plummets dramatically in the early phases of persistent infection. During this period, the PRRSV-specific T cell response at sites of infection is weak, transient, variable, and independent of local viral load. Surprisingly, pigs are largely resistant to reinfection with homologous or heterologous viral isolates even though neither humoral nor T-cell anamnestic

responses are observed. We hypothesize that PRRSV control in swine is due primarily to a reduction in permissive macrophages, and secondarily to specific humoral and cellular factors. Further investigation of immunological mechanisms that might provide insights to development of effective vaccines and immunotherapeutic strategies will require reagents that stimulate antigen-specific T cells and B cells and assays that are both specific and functionally relevant. Achieving these goals will require systems for expression and presentation of individual proteins or polypeptide fragments, tetramers, cytokine-specific ELISA, ELISPOT and FACS assays, and a better understanding of basic immunology of livestock species. Genomic identification of the gene repertoire involved in innate and adaptive immunity to specific pathogens and of the allelic variation in key regulatory and effector genes will be required to fully characterize basic immunological processes. Our initial foray into this area, identification of genes differentially expressed in porcine Peyer's patch, revealed that 45% of expressed sequences have no known function or are novel genes. Genes of unknown function also are differentially expressed in macrophages infected with PRRSV. These results indicate that the biochemical and cell biologic mechanisms responsible for immunity in swine, to a substantial degree, remain to be discovered. A better grasp of this knowledge will be required for rational design and genetic selection of disease resistant animals.

Genetic and immunologic basis for resistance to GI nematode infections

Louis C. Gasbarre, Tad Sonstegard, Ricardo Araujo, and Curt Van Tassell
USDA, ARS, Bovine Functional Genomics Laboratory, ANRI, BARC, Beltsville, Maryland

Gastrointestinal nematode infections of cattle are a constraint on the efficient raising of cattle throughout the world. Most of the nematodes species found in cattle stimulate a level of protective immunity that reduces the number of

worms that develop after secondary challenge, but cattle remain susceptible to reinfection by *Ostertagia* for a prolonged period of time (i.e. 2 years or more). This prolonged susceptibility to reinfection is a major reason that this parasite species remains the most economically important GI nematode in temperate regions throughout the world. Although, individual animals remain susceptible to reinfection for a prolonged period of time, there are a number of manifestations of the immune response that result in an enhanced level of herd immunity, and the overall result of these manifestations of immunity is a reduction in parasite transmission within the cattle herd. In cattle, the immune response seen in the abomasum shows high levels of expression of IL4 in the draining lymph nodes and in lymphocytes isolated from the mucosa, but effector cell populations in the tissues surrounding the parasites, are not typical, inferring that *Ostertagia* has evolved means to suppress or evade protective immune mechanisms, including the synthesis and release of immunosuppressive substances. The immune responses elicited by GI nematode infections are moderately influenced by host genetics, and a small percentage of individual animals in a given herd are responsible for the majority of parasite transmission. In general, cattle can be separated into three types: 1) those which never demonstrate high parasite numbers, 2) those that show increasing signs of infection during the first 2-3 months on pasture and then develop responses that regulate parasite numbers and or parasite fecundity, and 3) those which continue to maintain high parasite burdens. Studies are in progress to identify the genes controlling resistance. Two approaches are being utilized. The first involves mapping of quantitative trait loci (QTL) that influence the level of parasites infection. For these studies a population of cattle has been developed in which careful phenotypic characterization of calves has been combined with a complete genome scan to identify polymorphic markers that segregate with either increased or diminished resistance to infection. Using this approach QTL that affect parasite resistance have been demonstrated on 8 different bovine chromosomes. The second approach utilizes microarray technology to assess differential gene expression

in resistant versus susceptible cattle. We have developed a 400+ candidate gene array and have used it to look at gene expression in the local tissues of infected cattle. Results to date have indicated marked differences in gene expression patterns in genes associated with both adaptive and innate immune responses. Current work is focused on validation of these results by real time PCR, the development of global arrays for a more thorough look at the gene expression patterns, and integration of the QTL and array technologies for identification of important genes.

Heterophils isolated from chickens resistant to extraintestinal *Salmonella enteritidis* infection express higher levels of pro-inflammatory cytokine mRNA following infection than heterophils from susceptible chickens

Pamela J. Ferro¹, Christina L. Swaggerty², Pete Kaiser³, Igal Y. Pevzner⁴ and Michael H. Kogut²
¹Texas A&M University, Poultry Science, College Station, Texas; ²USDA, ARS, SPARC, College Station, Texas; ³Institute for Animal Health, Compton, Berkshire RG20 7NN, UK; ⁴Cobb-Vantress Inc., Siloam Springs, Arkansas

Previous studies in our laboratory have shown differences in *in vitro* heterophil function between two pure lines of broiler chicks and their F1 reciprocal crosses. The objectives of the current study were to 1) determine if the previously reported *in vitro* differences in heterophil function between the lines translated to an increase in *in vivo* resistance to extraintestinal infection with *Salmonella enteritidis* (SE) and 2) to quantitate pro-inflammatory cytokine mRNA expression levels in heterophils isolated from SE-infected chicks from different lines. Day-old chicks from four different lines (pure lines=A and B; line C = male B × female A; line D = male A × female B) were challenged orally with SE. Twenty-four hours post-challenge, livers and spleens were aseptically removed from each chick and incubated overnight in tetrathionate enrichment broth. Enrichment cultures were plated and subsequently analyzed for the presence of non-lactose fermenting *Salmonella* colonies. The two lines reported to have more functionally efficient

heterophils (A and D) were less susceptible to extraintestinal SE infections whereas the two lines reported to have functionally less efficient heterophils (B and C) were more susceptible to SE organ invasion. Four hours post-challenge peripheral blood heterophils were isolated from SE-infected and control chicks, total RNA extracted, and then analyzed for pro-inflammatory cytokine (IL-1 β , IL-6, IL-8) expression using real-time quantitative RT-PCR. Pro-inflammatory cytokine mRNA expression was significantly up-regulated in heterophils isolated from chicks that were more resistant to SE organ invasion (A and D) when compared to expression in heterophils isolated from the more susceptible lines (B and C). To our knowledge, this is the first report to quantitate IL-1 β , IL-6, and IL-8 mRNA expression levels in heterophils isolated from day-old chicks orally challenged with SE. These results show a relationship between in vitro heterophil function and resistance to organ invasion by SE. These data also indicate that pro-inflammatory cytokine mRNA expression by heterophils contributes, at least in part, to resistance/susceptibility of day-old chicks to extraintestinal infection by SE.

High-throughput molecular approaches to identify coccidiosis disease resistance genes and protective immunological mechanisms in avian coccidiosis.

Hyun S. Lillehoj¹, Wongi Min¹, Benjamin F. Matthews², and Marlene Emara³

¹USDA, ARS, Animal Parasitic Diseases Laboratory, ANRI, Beltsville, Maryland;

²Soybean Genomics Improvement Laboratory, Beltsville, Maryland; ³Department of Animal and Food Sciences, University of Delaware, Newark, Delaware

Avian coccidiosis is caused by several *Eimeria* strains which infect different regions of the

intestine inducing a strain-specific immunity. Coccidiosis usually stimulate a number of immunological defense mechanisms, namely both antibody- and cell-mediated. Recent technological advances in high-throughput molecular approaches to identify disease resistance genes and molecular/cell biological pathways associated with complex biological phenomenon now enable an alternative strategy to combat coccidiosis. We have hypothesized that chicken genes influencing resistance to coccidiosis can be identified through DNA marker technology. Supporting this hypothesis, we present data identifying a quantitative trait locus (QTL) on chromosome 1 that significantly affects *Eimeria* oocyst shedding and three QTL that influence body weight of chickens during coccidiosis. One hundred and nineteen microsatellite markers, covering 80 % of the chicken genome with an average marker interval of 25 cM, were used for genotyping of F1 parents and F2 offspring from commercial broilers. The genetic mechanism of this locus appeared additive. The genomic scan also identified three potential growth QTL on Chromosomes 1, 6, and 8. These results provide the foundation for further investigation to validate the QTL. In addition, our recent studies on local host immunity to *Eimeria* clearly indicate that intricate and complex interactions of host local cell-mediated immunity and parasites determine the outcome of host response to coccidiosis. High-throughput gene expression profiling and real-time PCR have been applied to analyze underlying immune mechanisms controlling disease resistance/susceptibility to coccidiosis. The role of various cytokines whose expression increase and decrease in response to coccidia invasion have been studied. Recently, we have developed the first chicken intestinal ETS database which will be used to investigate innate and acquired immunities to enteric pathogens such as coccidia and *Salmonella*.

Immunointervention

Protective killed *Leptospira* vaccine induces potent type 1 (Th1) immunity comprising responses by CD4 and WC1+ $\gamma\delta$ T lymphocytes

C.L. Baldwin

University of Massachusetts, Department of Veterinary and Animal Sciences, Amherst, Massachusetts

<http://www.umass.edu/vasci/faculty/baldwin.html>

Leptospira borgpetersenii serovar *hardjo* is a common cause of bovine leptospirosis and also causes zoonotic infections of humans. A protective killed vaccine against serovar *hardjo* was shown to induce strong antigen-specific proliferative responses by peripheral blood mononuclear cells (PBMC) from vaccinated cattle but not from non-vaccinated control cattle in three studies conducted. By 7 days, up to one-third of the cells in cultures of sonicated *leptospira* and PBMC from vaccinated animals produced interferon- γ (IFN- γ) as ascertained by flow cytometric analysis. The IFN- γ -producing cells included both WC1+ $\gamma\delta$ T cells and CD4⁺ T cells. Analysis of carboxyfluorescein succinimidyl ester loaded cells also showed that a similar proportion of these 2 subpopulations were found among the dividing cells in antigen-stimulated cultures. Infected non-vaccinated cattle also had an IFN- γ -based response but less than that by vaccinated animals and while naïve/challenged and vaccinated/challenged animals had similar levels of antigen-specific IgG1 following challenge, vaccinated animals had 1.4 fold more IgG2. In conclusion, while infection may induce a type 1 response we suggest it is too weak to prevent establishment of chronic infection. Overall the significance of these studies is the very potent type 1 (Th1) immune response induced and sustained following administration of a killed bacterial vaccine adjuvanted with aluminum hydroxide and the involvement of WC1 $\gamma\delta$ T cells in the response. The response by $\gamma\delta$ T cells is suggestive of a memory response since it has never occurred

with cells from a large number of naïve cattle tested. Moreover, recent results suggest it is a specific subpopulation of the WC1+ cells and we hypothesize that the response by these cells may result in establishment of a milieu that supports development of a type 1 response by CD4 T cells. Finally, induction of this cellular immune response is associated with the protection afforded by the bovine leptospiral vaccine against *L. borgpetersenii* serovar *hardjo* although this is an extracellular pathogen. Similar results have been found with a second monovalent vaccine made with *L. interrogans* serovar *hardjo* and adjuvanted with alum while we have found that a traditional pentavalent vaccine containing 5 serovars of *Leptospira* does not induce this response.

Development of an oral vaccine for *Escherichia coli* O157:H7

Evelyn A. Dean-Nystrom

USDA, ARS, National Animal Disease Center, Ames, Iowa

Enterohemorrhagic *Escherichia coli* (EHEC) O157:H7, a subset of Shiga toxin-producing *E. coli* (STEC), are the most common infectious cause of bloody diarrhea in the United States. A sequela of this infection, the hemolytic uremic syndrome (HUS), is the primary cause of acute kidney failure in U.S. children. Cattle are major sources of EHEC O157:H7 and other STEC pathogens associated with human infections. Our long-term goal is to identify ways to reduce EHEC infections in cattle, and thus reduce the risk of EHEC disease in humans. One approach, which is the objective of an ongoing collaboration with Alison O'Brien (USUHS, Bethesda, Maryland), is to develop an inexpensive, effective vaccine that will prevent cattle from becoming infected with EHEC. Intimin is an outer membrane protein of EHEC O157:H7 that is required for colonization of neonatal calves. We hypothesized that an intimin-based vaccination strategy in calves might reduce colonization of cattle with EHEC O157:H7. The demonstration that antibodies

against intimin interfere with colonization and intestinal damage in neonatal pigs supports this hypothesis. We transferred the *eae* gene that encodes for intimin into the tobacco NT-1 cell line and demonstrated that the transformed plant cells produce intimin, that transgenic intimin is immunogenic in mice, and that feeding mice transgenic intimin reduced the length of time these animals shed *EHEC O157:H7* following experimental challenge. Our collaborator, Wayne Curtis, recently scaled up the production of intimin-expressing NT-1 cells and the immunogenicity of transgenic intimin in cattle is being tested. The bacterial DNA that encodes for intimin is being modified so that it can be transferred into an edible plant that can be fed to calves and used to test the efficacy of transgenic intimin as an oral vaccine for reducing EHEC infections in cattle. Such a vaccine that reduces O157:H7 levels in cattle would help reduce EHEC infections in humans.

Adjuvant effect of IL-1 β , IL-2, IL-8, IL-15, IFN- α , IFN- γ , TGF- β 4 and lymphotactin on DNA vaccination against *Eimeria acervulina*.

Wongi Min¹, Hyun S. Lillehoj¹, and Peter Staeheli².

¹ USDA, ARS, Animal Parasitic Diseases Laboratory, Beltsville, Maryland; ²Department of Virology, University of Freiburg, D-79008, Freiburg, Germany

Avian coccidiosis, an economically important disease for the poultry industry, is caused by intracellular protozoa belonging to several species of the genus *Eimeria*. Coccidiosis results in extensive destruction of the gut epithelium accompanied by severe depression in body weight gain, reduced feed efficiency and intestinal shedding of parasite oocysts. Although coccidiosis is mainly controlled by the use of chemotherapeutic agents, alternative control strategies are needed due to the increasing emergence of drug-resistant parasite strains and impending withdrawal of drug use in animal farming. Recently, novel vaccination strategies using antigen-encoding DNA plasmids have been shown to successfully induce protective cellular

and humoral immune responses against a variety of infectious diseases. Since the emergence of gene-based immunization methodologies, various strategies have been used to increase the efficiency of DNA vaccination. For example, immune responses induced by DNA vaccination can be enhanced by cytokines. This study reports the first evidence of adjuvant effect of various chicken cytokines when given with coccidia DNA vaccine. Eight chicken cytokine genes (IL-1 β , IL-2, IL-8, IL-15, IFN- α , IFN- γ , TGF- β 4, lymphotactin) were evaluated for their adjuvant effect on a suboptimal dose of an *Eimeria* DNA vaccine carrying the 3-1E coccidia gene (pcDNA3-1E). One or 10 μ g of IFN- α or 10 μ g of lymphotactin expressing plasmids, when given simultaneously with the pcDNA3-1E vaccine, significantly protected against body weight loss induced by *E. acervulina*. Parasite replication was significantly reduced in chickens given the pcDNA3-1E vaccine along with 1.0 μ g of the IL-15, IFN- γ , TGF- β 4 or lymphotactin plasmids or 10 μ g of the IL-1 β , IL-8, IL-15 or TGF- β 4 plasmids compared with chickens given the pcDNA3-1E vaccine alone. In duodenum, chickens that received the pcDNA3-1E vaccine simultaneously with the IL-8 or IL-15 genes had significantly increased CD3+ cells compared with vaccination using pcDNA3-1E alone or in combination with the other cytokine genes tested. These results indicate that the type and the dose of cytokine genes co-injected with a DNA vaccine influence the quality of the local immune response to DNA vaccination against coccidiosis.

Stimulation of chicken leukocytes and reduction of *Salmonella* organ invasion in neonatal chicken by immunostimulatory CpG-oligodeoxynucleotide.

Haiqi He and Michael H. Kogut
USDA, ARS, Southern Plains Agricultural Research Center, College Station, Texas

Unmethylated CpG oligodinucleotide, flanked by particular base contents, are known to stimulate innate immune responses. Immune stimulatory activities of these CpG molecules have been studied mostly in immune cells from mammalian,

particularly human and mice. Recently, studies have emerged to show similar immune stimulatory activities of CpG oligodinucleotide in other species such as chicken and fish. Here, we present results from our studies on immune stimulatory activities of CpG oligodinucleotide in chicken. Our results show that synthetic oligodeoxynucleotides (ODN) containing CpG motif (CpG-ODN) activated the avian macrophage cell line and induced NO production. The most active CpG-ODN motif for NO induction was GTCGTT. Increasing the number of GTCGTT motif in the CpG-ODN significantly enhanced the stimulatory effect. This CpG-ODN also stimulated IL-1 β gene expression in the macrophage HD11 cells. CpG-ODN stimulated inducible NO synthase (iNOS) to produce NO through a clathrin-dependent endocytosis and subsequent endosomal maturation signaling pathways, involving activation of protein kinase C (PKC), mitogen-activated protein kinases (p38 MAPK and MEK1/2), and transcription factor NF- κ B. Moreover, we have discovered that CpG-ODN induces degranulation in chicken heterophils. This heterophil degranulation activity induced by CpG-ODN was absolutely serum-dependent and also receptor mediated. Finally, we evaluated synthetic CpG-ODNs for their *in vivo* protection against organ invasion by *Salmonella enteritidis* (SE) in neonatal chickens. Day-old chickens were i.p. injected with synthetic CpG-ODN and then orally challenged with live SE. Organs (liver and spleen) were removed from experimental chickens 24 hours after the challenge and cultured for SE. The results showed a significant reduction of organ invasion by SE ($p < 0.05$) in chickens pretreated with CpG-ODN containing the immunostimulatory GTCGTT motif.

Regulation of memory CD8 T cell generation by cytokines

M. Suresh

University of Wisconsin, School of Veterinary Medicine, Madison, Wisconsin

<http://www.pathology.wisc.edu/gradprogram/trainers/msuresh.html>

CD8 T cells play a critical role in defense against viral, intracellular bacterial, and protozoan

infections. Therefore, vaccines against these viruses need to engender potent CD8 T cell memory. Although generation of T cell memory is the basis of vaccinations, the mechanism(s) of induction of CD8 T cell memory are not well understood. Typically, CD8 T cell responses can be divided into three distinct phases: (1) the expansion phase when naive T cells undergo antigen-driven proliferation and differentiation into effector cells, (2) the contraction phase when 90-95% of the expanded T cells are eliminated, and (3) the memory phase when the remaining ~5% of the expanded CD8 T cells survive for extended periods as memory T cells. Upon infection, these memory CD8 T cells respond expeditiously and clear the infection by rapidly differentiating into effector T cells. The accelerated response that typifies immune memory can be attributed to both quantitative and qualitative changes in CD8 T cells: (1) Firstly, in an immune animal, there are more antigen-specific T cells, as compared to an immunologically naive animal; (2) Secondly, in striking contrast to naive T cells, the memory CD8 T cells in immune animals are hyper reactive to antigenic stimulation and capable of efficient trafficking into peripheral tissues. The number of memory CD8 T cells generated is dependent upon the extent of expansion and the magnitude of contraction during the T cell response. Since protective immunity is dependent upon the induction of a threshold number of memory CD8 T cells, it is critical to gain a thorough understanding of the mechanisms that regulate the expansion and contraction phases of the T cell response. Our studies using the well-characterized lymphocytic choriomeningitis virus (LCMV) model in mice have shown that signaling via IFN γ receptor (IFN γ R) and TNF receptors (TNFR) play important roles in limiting the number of antigen-specific memory CD8 T cells that survive the contraction phase of the T cell response. In mice, loss of IFN γ R reduced the expansion of LCMV-specific CD8 T cells, but remarkably ablated the contraction phase of the CD8 T cell response; IFN γ R-deficient mice contained substantially more virus-specific memory CD8 T cells, as compared to control wild type mice. Loss of either TNF or TNF receptors did not affect the expansion phase of the CD8 T

cell response to LCMV, but enhanced the survival of LCMV-specific memory CD8 T cells in both lymphoid and non-lymphoid organs. These

findings have implications in rational vaccine design and therapy of immune-mediated diseases.

Poster Sessions

December 2, 2003

The first author listed presented the poster unless another name is listed in **bold**.
Posters are numbered as indicated.

Innate Immunity

Analysis of swine skin-derived dendritic cells in the induction of innate immune responses to Foot-and-Mouth Disease virus (FMDV)

Elida M. Bautista, Geoffrey S. Ferman, Douglas Gregg, Mario C. Brum, Marvin Grubman and William T. Golde
USDA, ARS, Plum Island Animal Disease Center, ARS, USDA, Greenport, New York
See abstract, on page 16.

The inhibition of nitric oxide production in chicken phagocytes.

Tawni L. Crippen, Haiqi He and Michael H. Kogut
USDA, ARS, Southern Plains Area Research Center, College Station, Texas

The production of reactive nitrogen, nitric oxide (NO), has previously been demonstrated to be a major mechanism by which the innate immune system defends against microbial invasion. The purpose of this study was to characterize components of the signal transduction pathway induced by bacterial stimulation of NO for antimicrobial defense by chicken phagocytes. We quantified the production of nitrite by chicken macrophages (HD11 cells) which had been exposed to specific signal transduction inhibitors prior to stimulation by formalin-fixed *Enterococcus gallinarum* (EG) or *Klebsiella pneumoniae* (KP). The induction of many antimicrobial peptide genes is known to be regulated by factors of the NF- κ B family in a toll-like-receptor dependent or independent manner. NF- κ B is in turn regulated by I κ B kinase. We

found that NO production induced by both EG and KP was reduced in a dose dependent manner by specific inhibitors of NF- κ B, I κ B and JAK 3.

Oligodeoxynucleotides (ODN)-induced activation of macrophage-mediated innate immunity against intracellular pathogens, *Eimeria* and *Salmonella*, and the development of novel immunomodulatory strategies using embryo injection of CpG-ODNs in poultry.

Rami Dalloul^{1,2}, Hyun S. Lillehoj¹ Hang Xie², Robert A. Heckert², Uma Babu³, Richard Raybourne³, and Dennis M. Klinman⁴
¹USDA, ARS, Animal Parasitic Diseases Laboratory, Beltsville, Maryland; ²Virginia-Maryland Regional College of Veterinary Medicine, University of Maryland, College Park, Maryland; ³Center for Food Safety & Applied Nutrition, US FDA, Laurel, Maryland; ⁴Section of Retroviral Immunology, Center for Biologics Evaluation & Research, US FDA, Bethesda, Maryland

See abstract on page 15.

Toll-like receptor and acute phase cytokine expression in genetically selected lines of layers following an LPS challenge

S. D. Eicher and H. W. Cheng
USDA, ARS, Livestock Behavior Research Unit, West Lafayette, Indiana
A multi-disciplinary approach is essential to ascertain the well-being of farm animals. The Livestock Behavior Research Unit uses behavioral, physiological, and immunological

measures to answer well-being questions for poultry, swine, and dairy cattle. The innate immune system is frequently the first immune responder during many stressors and the resulting acute phase cytokines affect behavior and learning. Toll-like receptors are the pathogen recognition molecules that initiate the cascade that increases the acute phase cytokines. The effect of pathogens and stressors on the toll-like receptors of farm animals is not known. A study has been completed that examines the effect of lipopolysaccharide (LPS) on RNA toll-like receptor expression in chickens. The objective of this study was to determine expression of TLR2, TLR4, and two acute phase cytokines (interleukin-1 and tumor necrosis factor- α) following an LPS challenge of two genetic lines of layers divergently selected for aggression (aggressive, MBB and passive, KGB) and one commercial genetic line (COMM). The 3 lines of layers were given LPS (5.0 mg/kg of body weight), then organs (heart, liver, and spleen) were weighed and lung, liver, and spleen tissues were collected at 0, 24, and 72 hours following a single LPS i.v. injection. RNA was extracted from the tissues and subjected to real-time RT-PCR. Heart and spleen weights had genetic line ($P < 0.01$) and time effects ($P < 0.05$ and 0.01 , for heart and spleen respectively). Liver weights were different for genetic line ($P < 0.01$) but not time. Liver IL-1 and TLR2 had time effects ($P < 0.01$) but were not different among treatments. Liver TNF and TLR4 were not different over time or among treatments. Lung IL-1, TLR2, TLR4, and TNF were different over time ($P < 0.05$ for TLR4 and $P < 0.01$ for IL-1, TLR2, and TNF) but not among treatments. Spleen IL-1, TLR2, TLR4, and TNF were different over time but not among treatments ($P < 0.05$ for IL-1, $P < 0.01$ for TLR2, TLR4, and TNF). No treatment by time interactions were detected for any of these variables ($P > 0.05$). The only effects of genetic line were liver and spleen weight. Toll-like receptors 2 and 4 and cytokine changes were not influenced by genetic lines, only by time. Toll-like receptor expression was dissimilar for the two receptors that we investigated. Toll-like receptor 2 had effects of time in all three tissues, while TLR4 was influenced by time only in lung and spleen. Toll-

like receptor 2 increase expression in the liver and spleen by 24 h, but did not increase in the lung until 72 h post-LPS injection. Liver TLR2 remained elevated through the 72 h sample. Therefore, this genetic selection did not alter RNA expression of these innate immune receptors and cytokines. Since toll-like receptors recognize heat-shock proteins, further exploration of toll-like receptor expression during heat stress in chickens is in progress.

Toll-like receptor and acute phase cytokine expression in neonatal dairy calves

S. D. Eicher¹, T. A. Johnson², K. A. McMunn¹, and T. R. Johnson²

¹USDA, ARS, Livestock Behavior Research Unit, West Lafayette, Indiana; ²Purdue University, West Lafayette, Indiana

See abstract on page 14.

Spleen-dependent innate immunity in calves to *Babesia bovis* infection is characterized as a type-1 response involving the proliferation of NK-like cells

W.L. Goff*¹, W.C. Johnson¹, G.M. Barrington², W. C. Davis³ and D.P. Knowles¹

¹USDA, ARS, Animal Disease Research Unit, Pullman, Washington; ²Department of Clinical Medicine and Surgery, and ³Department of Microbiology and Pathology, Washington State University, Pullman, Washington

Innate immunity is important for control of primary infections and for induction of acquired responses. Young calves possess a strong spleen-dependent innate immunity to the intraerythrocytic protozoan, *Babesia bovis*. Prominent cellular components in innate immunity are dendritic cells, macrophages, and NK-cells. The study reported here demonstrates the importance of a type-1 response involving the early induction of IL-12 and IFN- γ and proliferation of NK-like cells in the spleen of calves during a primary tick-transmitted *B. bovis* infection. Unlike gdT-cells, CD3⁺, CD2⁺, CD8⁺ NK-like cells proliferated in the spleen, increasing almost three-fold over pre-infection levels. A model and hypothesis are presented for macrophage and/or dendritic cell participation

with NK cells in the induction of a type-1 protective response against this parasite.

Interferon- γ upregulates the expression of proinflammatory and Th1 cytokine mRNA in chicken heterophils during receptor-mediated phagocytosis of opsonized and nonopsonized *Salmonella enteritidis*.

Michael Kogut, Lisa Rothwell, Pete Kaiser
USDA, ARS, Southern Plains Area Research Center, and Institute for Animal Health, College Station, Texas; Compton, Berkshire, UK

The immediate response to invasive pathogens, clearance via the inflammatory response, and activation of the appropriate acquired responses are all coordinated and orchestrated by the innate host defenses. Recognition of microbes is accompanied by the induction of multiple cell processes including the production of pro- and anti-inflammatory cytokines. Polymorphonuclear leukocytes (PMNs) are vital cellular components of innate immunity, and function by killing pathogenic microbes following phagocytosis. The primary PMN in poultry is the heterophil. Priming is the potentiation of the phagocyte activation process by previous exposure to a priming agent. IFN- γ is a pleiotropic cytokine involved in basically all phases of immune and inflammatory responses that has been shown to prime heterophil functional activities. In the present experiments, using real-time quantitative RT-PCR, we evaluated the role of rChIFN- γ as a priming mediator to control heterophil responses at the level of gene transcription and expression of the mRNA for proinflammatory (IL-1b, IL-6, IL-8) and Th1 (IL-18 and IFN- γ) cytokine genes following stimulation with phagocytosis agonists opsonized and nonopsonized *Salmonella enteritidis*. RChIFN- γ primed the heterophils for an increase in transcription of pro-inflammatory cytokines induced by phagocytic agonists, but also upregulated expression of Th1 cytokine (IL-18 and IFN- γ) mRNA. Although rChIFN- γ priming modulated the expression of cytokine mRNA in heterophils stimulated by different phagocytic agonists, the rChIFN- γ by itself did not directly induce gene expression of either the proinflammatory or Th1

cytokines. The enhanced expression of cytokine mRNA does not appear to be differentially expressed depending on the receptor activated during phagocytosis. The results from the present experiments suggest that rChIFN- γ may play a significant role in avian innate immunity against *Salmonella* infection and may offer an adjunct use in the prevention and treatment of salmonellae infections in newly hatched chickens.

Recombinant bovine soluble CD14 reduces fatality of endotoxin challenged mice and reduces severity of intramammary infection by *Escherichia coli* in dairy cows.

Max J. Paape, Jai-Wei Lee and Xin Zhao
USDA, ARS, Beltsville, Maryland; McGill University, Ste-Anne-de-Bellevue, Quebec, Canada

Three million cases of clinical mastitis caused by Gram-negative bacteria occur every year in the U.S. A significant number of these cases will result in acute endotoxin shock and death. Three hundred thousand dairy cows are removed from herds or die annually because of acute coliform mastitis. Standard therapy for treating Gram-negative bacterial sepsis and shock, including administration of potent antibiotics, aggressive fluid resuscitation, and metabolic support, has not been successful in reducing mortality. Soluble CD14 (sCD14) neutralizes endotoxin and prevents it from binding and activating macrophages. Further, LPS/CD14 complexes bind to epithelial cells and results in rapid recruitment of neutrophils to the infection site. Novel reagents such as host defense proteins like sCD14 have a role as a potential prophylactic against serious Gram-negative bacterial infections. In the present study, the full length and the truncated forms of bovine sCD14 were cloned into a transfer vector and expressed by recombinant baculovirus that infect insect cell line sf-9. The protective effect of the recombinant bovine sCD14 (rbosCD14) was evaluated in mice and lactating dairy cows. Eighty-one female mice were randomly assigned to two groups and injected intraperitoneally with either LPS (8 μ g/g BW, n = 41) or LPS plus rbosCD14 (6.8 μ g/g BW, n = 40). Survival rate for LPS and LPS plus rbosCD14 injected mice at

24 h was 30 and 72%. Two mammary quarters of each of 9 cows were challenged with either 50 CFU of *Escherichia coli* plus saline or 50 CFU of *E. coli* plus 100 µg rboCD14. Quarters challenged with *E. coli* plus rboCD14 had a more rapid recruitment of neutrophils, which was accompanied by a faster clearance of bacteria and reduced clinical symptoms than challenged quarters injected with saline. At 24 h after bacterial challenge, only two (22%) of nine quarters challenged with *E. coli* and injected with

rboCD14 remained infected. On nine challenged quarters injected with saline, six (67%) remained infected. Results indicated that rboCD14 was able to decrease the fatality of LPS challenged mice and reduced incidence of infection in dairy cows following experimental challenge with *E. coli*. Use of rboCD14 in the treatment of and prevention of acute coliform mastitis has the potential of saving US dairymen 1.4 billion dollars annually.

Immune Regulation

Systemic cytokine and antibody responses following vesicular stomatitis virus infection of naïve and vaccinated cattle.

José Barrera^{1,2}, William T. Golde¹, D. Mark Estes² and Luis L. Rodriguez¹

¹USDA, ARS, Plum Island Animal Disease Center, Greenport, New York; ² University of Missouri. College of Veterinary Medicine. Department of Veterinary Pathobiology, Columbia, Missouri

Vesicular stomatitis virus (VSV) causes disease of varying severity in cattle, swine, and horses. The mechanisms mediating protection from clinical disease remain unknown. This study was carried out to gain a better understanding of the systemic T-helper cell balance mediating the immune response against VSV in cattle. We compared two cattle vaccinated with a commercial oil adjuvanted bivalent vaccine and four naïve animals, all challenged with a virulent strain of VSV New Jersey (VSNJV). Vaccinated animals developed high titers of neutralizing antibodies against VSNJV and VSV Indiana serotypes (VSIV) (range from 2.5 to 4.6 log) and showed no disease following challenge. In contrast, non-vaccinated naïve cattle developed fever and lesions. VSV was recovered from epithelial tissue and from esophageous-pharyngeal and vesicular fluids. VSNJV - specific IgM antibodies were detected in naïve infected cattle at 4 days post-infection (dpi) and class switch to IgG1 and IgG2 occurred by 14 dpi and 14 days post-vaccination (dpv) in naïve

and vaccinated cattle respectively. Titers of IgG2 isotype were higher than those of IgG1 isotype (IgG2/IgG1 ratio ranged from 1.2 to 6.5) both in naïve and vaccinated animals. IFN-α, IFN-β, IFN-γ, TNF-α, IL-1b, IL-2, IL-12p40, IL-4, IL-10 and IL-13 were measured by real-time reverse transcription-PCR (RT-PCR) in fresh peripheral blood mononuclear cells (PBMC) or PBMC stimulated with VSNJV antigen included. Transcripts for IFN-α, IFN-γ, and IL-12p40 were more abundant than those for IL-4, IL-10, and IL-13 in PBMC cultures -stimulated with VSNJV. Taken together, these results suggest that VSV antigen induces a T-helper Type 1 response in both infected and vaccinated cows. We propose that analysis of cellular responses at the sites of lesion and in draining lymph nodes, would give a more accurate picture of the events associated with development of immunity to this localized infection in domestic animals.

Mycoplasma gallisepticum: competitive exclusion or immune base

Stephanie D. Collier

USDA, ARS, Mid-South Area, Poultry Research Unit, Mississippi State, Mississippi

<http://msa.ars.usda.gov/ms/msstate/csrl/collier.htm>

Much confusion exists concerning the mechanism(s) by which the available *Mycoplasma gallisepticum* (MG) vaccines confer protection in layer chickens. One of the major goals of this laboratory is to discern the role of competitive

exclusion and/or immune based mechanisms following vaccination with MG. The parameters to be examined in the present study include T-cell subpopulations (CD4 and CD8) and cytokines levels (IFN-gamma, IL-6 and IL-8) immediately before and following MG vaccination *via* flow cytometry and real-time qRT-PCR.

Development of specific immunoglobulin IgGa and IgGb antibodies correlate with control of parasitemia in *Babesia equi* infection: IgG isotype responses in *B. equi* infection.

Cristina W. Da Cunha^{1,2}, Travis C. McGuire¹, Lowell S. Kappmeyer³, Stephen A. Hines¹, Alicia M. Lopez¹, Odir A. Dellagostin², AND Donald P. Knowles^{1,3*}

¹Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, Washington; ²Center for Biotechnology, Federal University of Pelotas, RS, Brazil; ³USDA, ARS, Animal Disease Research Unit, Pullman, Washington

Following infection with *Babesia equi* horses often survive acute disease but remain life-long carriers. Acquired immune responses are necessary to control replication while adaptive immune responses are not required for parasite-mediated erythrocyte lysis. Following lysis of infected erythrocytes, parasites are accessible to antibody and associated mechanisms of antibody-mediated control. To determine if IgG isotypes to *B. equi* erythrocyte stage antigens correlate with parasite control, antibody isotypes were measured following infection by parenteral inoculation or by the tick vector *Boophilus microplus*. Isotypes IgGa, IgGb, and IgG(T) were titrated in *B. equi* infected horse serum by ELISA using *B. equi* erythrocyte stage extract as antigen and monoclonal antibodies specific for each equine subisotype. IgGa and IgGb antibodies to *B. equi* developed during acute infection, but IgG(T) antibodies were detected only after resolution of acute parasitemia. During chronic infection, antibodies of all three isotypes were present at titers of 10³ to 10⁴ in naturally and experimentally infected horses. To test whether immunization induced IgGa and IgGb antibodies and not IgG(T)

antibodies similar to those during acute infection with *B. equi*, recombinant erythrocyte stage surface protein EMA-1 in the adjuvant saponin was tested. Four immunized horses developed IgGa and IgGb antibodies to *B. equi*, but no measurable IgG(T) antibodies. These results demonstrated that development of IgGa and IgGb antibodies to *B. equi* erythrocyte stage antigens correlated with control of *B. equi* parasitemia, and that this IgG isotype profile was induced by subunit immunization.

Analysis of lymphocytes isolated from white-tailed deer (*Odocoileus virginianus*) fawns

Sasha J. Fach¹; W. Ray Waters², Mitchell V. Palmer², William C. Davis³, **Randy E. Sacco²**

¹ Immunobiology Graduate Program, Iowa State University, Ames, Iowa; ² USDA, ARS, National Animal Disease Center, Ames, Iowa; ³ Department of Veterinary Microbiology and Pathology, College of Veterinary Medicine, Washington State University, Pullman, Washington

Peripheral blood mononuclear cells (PBMC) were isolated from ten female white-tailed deer fawns at 48 hours of age and every two weeks thereafter until the fawns were three months of age. Bone marrow aspirates and peripheral blood samples were taken on the same fawns at ten months of age (yearlings). In a separate experiment, mesenteric lymph node (MLN), spleen, thymus, bone marrow (BM) and peripheral blood (PB) were collected from euthanized female fetuses at approximately 190 days gestation (n=4) and from fawns at 24 hour (n=2), two weeks (n=2) and four weeks (n=2) of age. Mononuclear cells were isolated from pooled, homogenized tissues and individual peripheral blood samples at each of the four time points. Lymphocytes were phenotyped to examine the expression of specific surface receptors as the fawns aged. Three-color flow cytometric analysis of leukocytes was performed using a panel of monoclonal antibodies (mAb). Adult deer were also phenotyped using the same monoclonal antibody panel. Included in the panel were mAbs recognizing WC1, the α chain and β chains, CD4, CD8, CD62L, CD44, CD21, IL-2R, MHC Class II, B-B1, B-B2 and B-B4. Analysis revealed

dynamic changes in the expression of specific surface receptors associated with development and maturation of lymphocyte subpopulations. WC1⁺ gamma delta T cells were predominant in the PB of fetal and neonatal fawns and decreased with age. In contrast, percentages of CD4⁺ and CD8⁺ populations were observed to increase over time in the PB. T cell surface antigen expression in the PB and tissues was consistent with observations made in other ruminants. B cell surface IgM expression was heterogeneous at two days, but became more discrete as the fawns matured. Interestingly, B-B2, a putative B cell lineage marker expressed on lymphocytes in the PB at 24-48 hours was down modulated by two weeks. However, expression of another B cell lineage marker, B-B4, was consistently expressed throughout the fawns' development in the PB, MLN, and spleen. Mononuclear cells isolated from bone marrow aspirates revealed phenotypically distinct expression of surface receptors as compared to PBMCs. Mononuclear cells co-expressing B-B2 and surface IgM were a unique population found in the bone marrow and not in the PB. The sIgM⁺B-B2⁺ population was observed in the fetal spleen and MLN. B-B4 was expressed on a small percentage of bone marrow mononuclear cells. Proliferative responses of the isolated PBMC's to pokeweed mitogen (PWM), Concanavalin A (ConA) and *Mannheimia (Pasteurella) haemolytica* LPS were also examined. Cells isolated from fawns proliferated in response to PWM and ConA, but not in response to LPS stimulation. Data obtained in the present study provides baseline information regarding lymphocyte subpopulations in white-tailed deer fawns.

Effects of the mammary gland on the composition and function of peripheral blood mononuclear leukocyte populations in periparturient dairy cows.

J.P. Goff, K. Kimura, and B.J. Nonnecke
Periparturient Disease of Cattle Research Unit
USDA, ARS, National Animal Disease Center,
Ames, Iowa

See abstract on page 19.

Comparison of adhesion molecules on milk and blood lymphocytes from periparturient dairy cattle.

James A. Harp*, Theresa E. Rahner, and Jesse P. Goff.

USDA, ARS, Periparturient Diseases of Cattle Research Unit, National Animal Disease Center, Ames, Iowa

Fifteen Holstein dairy cows were monitored for expression of adhesion molecules on lymphocytes in blood and milk at parturition and at intervals up to 21 days post partum. Using flow cytometric analysis, we examined expression of CD62L, CD11a, CD44, and $\alpha_4\beta_7$ on T (CD4⁺, CD8⁺, $\gamma\delta$ +) and B lymphocytes. CD62L was expressed on 20-70% of T cell subsets in blood and 35-80% in milk from normal cows. CD62L expression on B cells was 20-25% in blood and 60-80% in milk. CD11a was found on 10-25% of T cells in blood and 35-70% in milk. CD11a was found on 10-15% of B cells in blood and 45-75% in milk. $\alpha_4\beta_7$ was expressed on 5-10% of T cells in blood and 10-35% in milk. $\alpha_4\beta_7$ was expressed on 5-10% of B cells in blood and 20-40% in milk. CD44 was found on 5-15% of T cells in blood and 20-55% in milk. CD44 was found on 5-10% of B cells in blood and 30-55% in milk. There were consistently more adhesion molecules, at a statistically significant level, on milk lymphocytes compared with blood lymphocytes. These differences may reflect increased recruitment of lymphocytes into the bovine mammary gland, which could then modulate the inflammatory response to pathogens during infection. Since CD62L and $\alpha_4\beta_7$ are peripheral and mucosal homing receptors, respectively, these data suggest increased recruitment of both pools of lymphocytes into the bovine mammary gland.

Evaluation of the chicken crop (ingluvies) as a model system for following mucosal immunity in chickens.

Peter S. Holt

USDA, ARS, Southeast Poultry Research Laboratory, Athens, Georgia

The crop (ingluvies) is an enlargement or out pouching of the esophagus proximal to the proventriculus, or glandular stomach and functions primarily in a food storage role for avian species. While recent work has shown the crop can be colonized with pathogens such as *Salmonella*, little information is available regarding the presence of any immune response within this organ. During a recent study investigating mucosal immunity to *Salmonella enteritidis* (SE) in the alimentary tract of White Leghorn chickens, we discovered the presence of substantial titers of IgA specific for SE in samples from crops removed from SE-infected, but not from noninfected, birds. Histological examination of the crop tissues found a paucity of lymphocytes in noninfected birds but lymphocyte aggregates were detected in the crop mucosa by 2 weeks post infection which increased in size and number thereafter. Because of the easy access to this organ orally, a crop lavage system was developed where a narrow diameter plastic tubing attached to a syringe was inserted down the esophagus into the crop. A flush solution was administered into the crop and immediately aspirated back into the syringe, with minimal discomfort to the bird and allowing multiple samplings of the same live birds over time. While specific IgA responses were the most robust, high levels of SE-specific IgG were also detected. Comparing IgA titers in crop lavage samples vs intestinal flushes from individual SE-infected birds, a very good correlation between titers of the two sample types was obtained, indicating that crop immunity may be used as an easy model for following mucosal immunity in a bird following infection. Studies are currently underway to determine crop immune responses in chickens infected with other pathogens, the prevalence of lymphoid tissues in crops from different avian species, the phenotypes of lymphocyte populations that develop in the chicken crop post infection and the importance of mucosal invasion in driving the development of this lymphoid tissue.

Neutralization of bluetongue virus in vertebrate and invertebrate cells.

James O. Mecham

USDA, ARS, Arthropod-borne Animal Diseases Research Laboratory, Laramie, Wyoming

Bluetongue virus (BTV) is an orbivirus that infects sheep, cattle and wild ruminants. It is transmitted to susceptible vertebrate hosts by biting midges in the genus *Culicoides*. Following infection, the vertebrate host produces neutralizing antibodies that are responsible for virus clearance and conferring immunity to subsequent virus challenge. These neutralizing antibodies are directed against the virus outer capsid, which is also responsible for initiating infection in the vertebrate host. Infection of the insect vector is believed to be mediated primarily by the inner capsid, which is not the target of neutralizing antibody. Therefore, virus that is neutralized for infection of vertebrate cells may still be infectious for invertebrate cells. This may have important implications in the natural epidemiology of BTV infections. To investigate this further, the mechanisms of neutralization of BTV in both vertebrate and invertebrate cells was investigated. Antibodies that neutralize BTV infection in vertebrate cells failed to neutralize viral infection in a *Culicoides* cell line. However, non-neutralizing antibodies that react with a virus core protein inhibited infection and/or replication of BTV in the *Culicoides* cell line. Since this virus core protein is highly conserved between BTV serotypes, and appears to play an important role in the initiation of BTV infection in insect cells, it may represent a novel target for control of infection caused by these viruses.

Comparison of the adaptive immune response in neonatal and young adult dairy cattle utilizing a *Mycobacterium bovis* BCG sensitization/PPDb challenge model.

B. J. Nonnecke¹, M. R. Foote², B. L. Miller³, M. A. Fowler³, W. R. Waters⁴

¹USDA, ARS, Periparturient Diseases of Cattle Research Unit, National Animal Disease Center, Ames, Iowa; ²Iowa State University, Ames, Iowa; ³Land O'Lakes, Inc., Webster City, Iowa; ⁴USDA, ARS, Bacterial Diseases of Livestock Research Unit, National Animal Disease Center, Ames, Iowa.

Acquisition and modulation of immune competency in calves during the neonatal and postnatal period are not well described. The purpose of this study was to characterize age-related changes in the functional capacities of peripheral blood mononuclear cell (PBMC) and neutrophil populations from antigen-sensitized calves. Milk replacer-fed Holstein bull calves were nonvaccinated (n = 6, vaccination controls) or vaccinated subQ (n = 6) with an attenuated strain of *Mycobacterium bovis* [bacille Calmette Guerin (BCG)] at 1 and 7 wk of age. Yearling heifers (n=4, adult controls) were vaccinated and sampled concurrently with the calves. Functional capacities of PBMC (i.e. mitogen and antigen-induced DNA-synthesis, T cell subset proliferation, cytokine and NO production, IgG secretion) and neutrophil (iodination and cytochrome C reduction) populations collected at 0 (pre-vaccination), 2, 5, 6 (boosted), 7, 8 and 11 wk were evaluated *in vitro*. The phenotype of PBMC populations was characterized by flow cytometry. The responsiveness of PBMC from nonvaccinated and vaccinated calves to polyclonal stimulation frequently was lower than adult PBMC. Responses of PBMC from vaccinated calves to recall antigen (PPDb) were evident at >2 wk (3 wk of age) after primary vaccination. With the exception of antigen-induced DNA synthesis, most functions (T cell subset proliferation, TNF- α , IFN- γ and NO secretion) of PBMC from vaccinated calves were comparable to or exceeded those of vaccinated adults. Changes in cervical skin-fold thickness after intradermal administration of PPD (wk 11) were significant and comparable in vaccinated calves and adults confirming the development of a robust cell-mediated immune response in vaccinated calves. In contrast, antigen-specific IgG responses (i.e. PPD-specific IgG in sera and supernatants from PPD-stimulated cultures) of vaccinated calves were minimal or nonexistent relative to adult responses suggesting the contribution of the B cell to the immune response is limited in young cattle. Nonvaccinated calves did not respond to recall antigen *in vivo* (i.e. cutaneous delayed-type hypersensitivity) or *in vitro*. Neutrophil functions were affected by age but not vaccination.

Molecular characterization of a putative Tumor Necrosis Factor superfamily 13b (TNFSF 13b) homologue from rainbow trout (*Oncorhynchus mykiss*).

Gregory D. Wiens*, Katherine Hovatter, Yniv Palti, Carid Rexroad III, Timothy Welch.

*USDA, ARS National Center for Cool and Cold Water Aquaculture, Kearneysville, West Virginia

Little is known about the immune system function of economically important cool and cold water fish species. Such knowledge is important for developing successful vaccines and disease resistant fish. In mammals, members of the tumor necrosis factor superfamily (TNFSF) play important roles in both innate and adaptive immune responses. In order to determine which TNFSF members are conserved and expressed in rainbow trout, we have searched a computer database of ~50,000 expressed sequence tags generated at the National Center for Cool and Cold Water Aquaculture. We have identified 17 cDNAs that have homology to known TNFSF ligands or receptors. Here, we describe one cDNA, 1RT126I22, which has homology to TNFSF 13b. TNFSF 13b is also known as B-cell activating factor (BAFF) or B-lymphocyte stimulating factor (BLYS). In mice and humans, TNFSF 13b is produced by myeloid cells and is critical for mature B cell survival and antibody production. The rainbow trout cDNA, 1RT126I22, encodes a predicted 141 amino acid protein, which shares 54% amino acid identity with human TNFSF13b, 51% identity with mouse TNFSF13b, and 54% identity with chicken TNFSF13b. The rainbow trout TNFSF 13b shares 100% identity with a 99 amino acid protein predicted from a truncated Atlantic salmon cDNA, and 78% identify with puffer fish TNFSF13b protein predicted from genomic DNA sequence. The rainbow trout TNFSF 13b is unusual as it does not possess a transmembrane domain or a furin-type protease cleavage site characteristic of the human, mouse, and chicken TNFSF 13b molecules. Examination of rainbow trout TNFSF 13b gene expression during vaccination and bacterial challenge is underway. To our knowledge, this is the first identification and characterization of fish TNFSF 13b homologues.

Immunogenetics

Expression of specific MHC Class II DRB1 alleles associates with sheep persistently infected with ovine progressive pneumonia virus (OPPV).

L.M. Herrmann¹, W.P. Cheevers², D.P. Knowles^{1,2}

¹USDA, ARS, Animal Disease Research Unit, Pullman, Washington; ² Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, Washington

See abstract on page 22.

Marek's disease and the MHC: How viruses subvert the immune system for their advantage.

H. Hunt, H. Cheng, and H. Zhang
USDA, ARS, Avian Diseases and Oncology Laboratory, East Lansing, Michigan

See abstract on page 21.

Physical and genetic mapping of the rainbow trout major histocompatibility regions: evidence for duplication of the class I region

Ruth Phillips¹, Ana Zimmerman², Marc Noakes¹, Yniv Palti³, Matt Morash¹, Lisa Eiben⁴, Sandra S. Ristow², Gary H. Thorgaard² and John D. Hansen⁴.

¹School of Biological Sciences, Washington State University, Vancouver, Washington; ²School of

Biological Sciences, Washington State University, Pullman, Washington; ³USDA, ARS-NCCCWA, Kearneysville, West Virginia; ⁴Center of Marine Biotechnology, University of Maryland Biotechnology Institutes, Baltimore, Maryland

Previous segregation analysis in trout suggested that the class I and II regions reside on completely different chromosomes. To learn more about MHC genomics in trout, we have isolated BAC clones harboring class IA and IB loci, a single BAC clone containing an MH class II gene (DAB) as well as BAC clones containing the ABCB2 gene. Upon PCR and sequence confirmation, BAC clones were labeled and used as probes for *in situ* hybridization on rainbow trout metaphase chromosomes for determination of the physical locations of the trout MH regions. Finally, SNPs, RFLPs and micro-satellites found within the BAC clones allowed for these regions to be assigned to specific linkage groups on the OSUxHotcreek and OSUxArlee genetic linkage maps. Our data demonstrates that the trout MH regions are located on at least four different chromosomes and the corresponding linkage groups while also providing direct evidence for the partial duplication of the MH class I region in trout.

Immunity and the Pathogenesis of Disease

A dual role for FLICE-like inhibitory protein (FLIP) in mediating both bacterial lipopolysaccharide-induced apoptosis and NF- κ B activation.

Douglas D. Bannerman¹, Kristine T. Eiting², Robert K. Winn², and John M. Harlan²

¹ USDA, ARS, Immunology and Disease Resistance Laboratory, Beltsville, Maryland;

² University of Washington School of Medicine, Seattle, Washington

Bacterial lipopolysaccharide (LPS) contributes to much of the vascular injury/dysfunction associated with Gram-negative sepsis. We have previously shown that inhibition of *de novo* gene expression with cycloheximide sensitizes vascular endothelial cells (EC) to LPS-induced apoptosis and that the onset of apoptosis correlates with decreased expression of FLICE-like inhibitory protein (FLIP), an anti-apoptotic protein. We now have data that conclusively establish a role for FLIP in protecting EC against LPS-induced apoptosis. Downregulation of FLIP using anti-sense oligonucleotides sensitized EC to direct LPS killing, whereas, overexpression of FLIP protected against LPS-induced apoptosis in the presence of cycloheximide. Interestingly, FLIP overexpression completely blocked NF- κ B activation induced by LPS, but not by phorbol ester, suggesting a specific role for FLIP in mediating LPS activation. Together, these data support a dual role for FLIP in mediating LPS-induced apoptosis and NF- κ B activation.

Gene expression profiling of bovine macrophages in response to *Escherichia coli* O157:H7 lipopolysaccharide

Carol G. Chitko-McKown*, James M. Fox, Laura C. Miller, Michael P. Heaton, James L. Bono, James E. Keen, William M. Grosse¹, William W. Laegreid
USDA, ARS, Roman L. Hruska U.S. Meat Animal Research Center, Clay Center, Nebraska

The aim of this study was to identify changes in bovine macrophage gene expression in response to treatment with *Escherichia coli* O157:H7 lipopolysaccharide (LPS), utilizing a human gene microarray. Bovine cDNA from control and LPS-treated primary macrophages hybridized to greater than 5,644 (79.8%) of the non-control gene targets on a commercially available microarray containing greater than 7,075 targets (Incyte Genomics, St. Louis, MO). Of these target sequences, 44 were differentially expressed upon exposure to LPS, including 18 genes not previously reported to exist in cattle. These included a pentaxin-related gene, CASP8, TNF-induced genes, interferon-induced genes, and inhibitors of apoptosis. Using the human microarray, cDNA from bovine LPS-treated and control macrophages consistently hybridized to targets known to be expressed constitutively by macrophages, as expected given the predicted cDNA sequence homology. That this human system was accurately estimating levels of bovine transcripts was further verified by real-time quantitative reverse transcriptase polymerase chain reaction (RTQ-PCR) using bovine-specific primers. This first report of bovine-human cross-species expression profiling by microarray hybridization demonstrates the utility of this technique in bovine gene expression and discovery.

Evaluation of immune response induced by attenuated and wild-type PRRS virus

J Janutenaite¹, K M Lager², T Stabel², B Pesch², S L Brockmeier²

¹National Veterinary Laboratory, Vilnius, Lithuania; ²USDA, ARS, National Animal Disease Center, Ames, Iowa

Introduction and Objectives

Porcine reproductive and respiratory syndrome virus (PRRSV) modified-live vaccines (MLV) have been developed by attenuation in cell culture. Under experimental conditions the vaccines are efficacious; however, under field conditions

vaccine efficacy reports have ranged from excellent to poor. One possible reason for questionable efficacy is that MLV may not induce a similar immune response when compared to wild-type PRRSV infection. The goal of the study reported here was to compare the cellular immune response in pigs following exposure to an attenuated PRRSV or its wild-type parent virus.

Materials and Methods

Two experiments were conducted using 52 4-5 week-old conventionally-raised pigs free of PRRSV and PRRSV-specific antibody. PRRSV strain NADC-8 (3rd passage) and NADC-8 (251st passage) were wild-type parent virus and attenuated virus, respectively. Based on sequence of ORFs 2-7 (3,184 nucleotides) the attenuated virus was 99.4% homologous with its parent virus. A different wild-type PRRSV strain, SDSU 73 (3rd passage) was used in the 4th group in experiment 2. Each experiment had a similar design: a sham-inoculated control group (group 1), an attenuated virus-inoculated group (group 2) and a wild-type parent virus-inoculated group (group 3). In experiment 2 there was a 4th group inoculated with SDSU 73. There were 13, 15, 15 and 8 pigs in groups 1, 2, 3 and 4, respectively. Pigs were each inoculated intramuscularly with 2 ml of their respective virus inoculum (about 2×10^5 TCID₅₀ virus/ml) day 0, pigs in group 1 received 2 ml of sham inoculum. Blood samples were collected on days -7, -4, 0, 3, 6, 9, 13 and 20 (21 for experiment 2). Serum was tested for PRRSV specific antibodies and used for PRRSV virus isolation. A rapid flow cytometric method (6) and a panel of monoclonal antibodies were used to analyze porcine peripheral blood leukocytes by two-color flow cytometric analysis. Statistical analysis was performed using GraphPad InStat software.

Results and Discussion

Results represent the combined data from the two experiments. Statistically significant changes ($P < 0.05$) in lymphocyte subpopulations percentage were observed in groups 3 and 4; whereas no changes were observed in group 2. CD8⁺ T cells increased on days 6, 9 and 13 for group 3 and on days 9 and 13 for group 4. CD4⁺ T

cells increased on day 9 for group 4. CD3⁺ T cells increased on day 9 for groups 3 and 4. Cells expressing Class II marker decreased on day 3 in group 3 and on days 3 and 6 in group 4; Class II cells increased in the both groups on days 13 and 20(21). CD45⁺ cells increased on days 9 and 13 for groups 3 and 4. B cells decreased on days 3, 6 and 9 for group 3 and on days 3 and 6 in group 4. Total WBC counts decreased on days 6 and 9 in group 3 and on days 3, 6 and 9 in group 4. Virus was isolated from all wild-type PRRSV-infected pigs (groups 3 and 4), but no virus was isolated from the sham or attenuated-virus inoculated pigs (groups 1 and 2). All wild-type PRRSV-inoculated pigs and 6 of the 15 attenuated-virus inoculated pigs (group 2) developed PRRSV-specific antibody. No sham-inoculated pigs developed antibody. Group 2 data were further analyzed by dividing pigs in seropositive and seronegative subgroups. No statistical differences in percentage of lymphocyte populations were found.

The percentage lymphocyte values determined in this study for the naïve animals represent normal values for conventionally raised pigs and are in agreement with previously reported data (6). This agreement of data suggests the methodology used in these experiments is reproducible. The peripheral blood cellular immune response characterized in the pigs infected with wild-type PRRSV appears similar to what has been reported by others. Following experimental infection with wild-type PRRSV there may be a transitory decrease in CD4⁺ and CD8⁺ T cells within the first few days. This is followed by a return of CD4⁺ cells to normal levels and an elevated level of CD8⁺ cells that may last for at least several weeks (1-5) Collectively, these studies in conjunction with the present study suggest T cells play an important role in the PRRSV immune response following wild-type virus infection. No significant changes in blood cell populations were noted following infection with the attenuated PRRSV (group2) when compared to negative control group. Perhaps this observation may account for the lack of vaccine efficacy that has been described under field conditions.

The study reported here indicates that the magnitude of the PRRSV-specific immune response may be related to PRRSV virulence. Additional studies are needed to confirm this hypothesis and what, if any, relevance this may have on vaccine development.

References:

Albina, E., et al. 1998. *Veterinary Immunology and Immunopathology* 61: 49-66.

Meier, W., et al. 1999. *Proceedings of the PRRS [Porcine Reproductive and Respiratory Syndrome] and Aujeszky's Disease Symposium*, p. 71-74.

Nielsen, J., and A. Botner. 1997. *Veterinary Microbiology* 55: 289-294.

Samson, J.N., et al. 1999. *Proceedings of the PRRS [Porcine Reproductive and Respiratory Syndrome] and Aujeszky's Disease Symposium*, p. 81.

Shimizu, M., et al. 1996. *Veterinary Immunology and Immunopathology* 50: 19-27.

Stabel, T., et al. 2000. *Journal of Immunological Methods* 245: 147-152.

Decreased neutrophil function as a cause of retained placenta in dairy cattle

Kayoko Kimura, Jesse P. Goff, Marcus E Kehrli, Jr., and Timothy A. Reinhardt
USDA, ARS, National Animal Disease Center,
Periparturient Disease of Cattle Research Unit
Ames, Iowa

It is unclear why some cows fail to expel the placenta following calving. One theory suggests the fetal placenta must be recognized as "foreign" tissue and rejected by the immune system after parturition to cause expulsion of the placenta. We hypothesized that impaired neutrophil function causes retained placenta (RP). We examined the ability of neutrophils to recognize fetal cotyledon

tissue as assessed by a chemotaxis assay. Neutrophil killing ability was also estimated by determining myeloperoxidase activity in isolated neutrophils. Blood samples were obtained from 142 periparturient dairy cattle in 2 herds. Twenty cattle developed RP (14.1 %). Neutrophils isolated from blood of cows with RP had significantly lower neutrophil function in both assays prior to calving and this impaired function lasted for 1-2 wk after parturition. Addition of antibody directed against interleukin-8 (IL-8) to the cotyledon preparation used as a chemoattractant inhibited chemotaxis by 41% suggesting one of the chemoattractants present in the cotyledon at parturition is IL-8. At calving plasma IL-8 concentration was lower in RP cows (51 ± 12 pg/ml) than in cows expelling the placenta normally (134 ± 11 pg/ml). From these data, we suggest that neutrophil function is a determining factor for the development of RP in dairy cattle. Also, depressed production of IL-8 may be a factor affecting neutrophil function in cows developing RP.

Comparative proteomic analysis of normal and periparturient bovine neutrophils demonstrating a reduction of myeloperoxidase during transition.

John D. Lippolis, Jesse P. Goff and Timothy Reinhardt
USDA, ARS, National Animal Disease Center,
Ames Iowa

Neutrophils constitute a primary effector of the host immune response to an infectious pathogen. Normally low numbers of neutrophils reside in a health mammary gland, however, infection and subsequent release of chemoattractants results in the rapid influx of activated neutrophils into the site of infection. Periparturient dairy cows exhibit signs of immunosuppression that correlate with a higher incidence of mastitis. Furthermore, periparturient immunosuppression includes a significant reduction of neutrophil function. To better understand the nature of the periparturient immunosuppression, neutrophils from dairy cows were isolated greater than 21 days prior to calving (normal function) and shortly after calving (immunosuppressed) and compared using

differential mass spectrometry. Proteomes of the immunosuppressed and normal neutrophils were labeled with isotope-coded affinity tags (ICAT), digested and subjected to nano-flow LC-MS/MS analysis. Neutrophil proteins were identified and quantitated, interestingly myeloperoxidase was shown to be reduced three-fold around parturition whereas cathelicidin 1 and beta defensin were unaffected. Myeloperoxidase is a strongly bactericidal neutrophil enzyme that generates reactive oxidants and is considered an important component of neutrophil's ability to destroy bacteria.

The effects of *Pasteurella multocida* toxin (PMT) on early murine B-lineage cells

Crystal L. Loving¹ and Randy E. Sacco²

¹Immunobiology Graduate Program, Iowa State University; ²USDA, ARS, National Animal Disease Center, Ames, Iowa

Pasteurella multocida toxin (PMT) induces pre-osteoclast proliferation and osteoclast differentiation that leads to bone destruction. Developing B-cells and osteoclasts share the same microenvironment; therefore, it is hypothesized that the toxin may elicit alterations in B-cell progenitors in the bone marrow. To test this, mice were exposed to the toxin for 24 hours; peripheral blood, spleen, and bone marrow cells isolated, counted, and developing B-cell populations analyzed by flow cytometry. Toxin exposure resulted in a 5% increase in the percentage of B220⁺CD19⁺IgM⁻ cells in the bone marrow, as well as a 5% increase in the percentage of B220⁺CD19⁺IgM⁺ cells in the peripheral blood. Conversely, percentages of B-cell subsets in the spleen were similar in toxin-treated and control mice. The increase in bone marrow B220⁺CD19⁺IgM⁻ cells was not due to proliferation as determined by *in vivo* administration of bromodeoxyuridine (BrdU) during PMT exposure. Furthermore, PMT did not alter the expression of the anti-apoptotic Bcl-2 as measured by intracellular staining. As B220⁺ cells have been shown to serve as osteoclast precursors, we are currently investigating whether PMT induces osteoclastogenesis of bone marrow B220⁺CD19⁺IgM⁻ cells.

Functional genomics analysis of infection by bovine viral diarrhea virus Type 2 in a bovine B-cell lymphosarcoma cell line

John D. Neill and Julia F. Ridpath

USDA, ARS, National Animal Disease Center, Ames, Iowa

Bovine viral diarrhea virus (BVDV), a major viral pathogen of cattle in the US, causes disease characterized by immunosuppression and lymphocytopenia. In addition, infection of a fetus in the first 150 days of pregnancy can give rise to a persistently infected calf that sheds the virus for life. This is a major means of dissemination of the virus to naïve animals and herds. BVDV is lymphotropic and acute infection results in depletion of lymphoid tissues, even with BVDV strains of low virulence. The limited *in vitro* studies of host cell response done to date were performed in epithelial cells and compared cytopathic and noncytopathic strains without regard to virulence *in vivo*. Our research has shown that cytopathology *in vitro* does not correlate with virulence *in vivo*. We are taking a functional genomics approach, using serial analysis of gene expression (SAGE), to determine the molecular events that take place in the infected lymphoid cells that contribute to the disease syndromes observed *in vivo*. SAGE is a powerful technology that allows quantitation of transcripts in cells and direct comparisons between libraries and experiments. For these studies, we used the B-cell lymphosarcoma cell line BL-3 and compared virus strains of differing virulence (virulence was based on previous *in vivo* studies). To date, we have constructed and sequenced SAGE libraries from mRNA derived from non-infected cells, as well as cells infected with high and low virulence BVDV strains. Data analysis is ongoing. Preliminary results from the non-infected cells showed the normal metabolism of these cells with corresponding gene expression levels. When compared to results from BVDV-infected cells, a remodeling of cellular functions was observed that reflected changes that, overall, benefited the replication of the virus. These effects included changes in protein translation, transport and ER processing, intracellular signaling, and some important housekeeping functions. Also observed

was a decline in expression of proteins that are involved in detoxification, particularly reactive oxygen species. Some of these changes have suggested new lines of research that should lead to elucidation of the mechanism(s) behind immunosuppression and viral persistence.

Immune responses controlling *Toxoplasma gondii* infection in pigs.

S.M. Nishi¹, H. Dawson², J.P. Dubey³, J.F. Urban² and J.K. Lunney¹.

¹USDA, ARS, Immunology and Disease Resistance Lab, Beltsville, Maryland; ²Nutrient Requirements and Functions Lab, BHNRC, and ³Parasite Biology, Epidemiology and Systematics Lab, ANRI, Beltsville, Maryland

Toxoplasma gondii (Tg) infection induces a strong interferon-gamma (IFN γ) dominated, T helper-1 (Th1) response in pigs with immune pathology that quickly resolves. Pigs were infected with 4.5×10^6 Tg oocysts and various tissues were collected at 2, 4, 7 and 14 days after infection (DAI). Tg organisms were detected in mesenteric lymph nodes (MLN) at 2DAI and, at 4 DAI, in liver, spleen, lung and ileum, by IHC and Tg specific Real-Time PCR. At the same infection times changes in inflammatory and Th1 immune mRNA gene expression were observed by Real-Time RT-PCR. Increased IFN γ mRNA and protein expression were observed in MLN and hepato-splenic (HS) LN cells. Changes were observed in IL1b, IL-6, NRAMP1, IFN γ , INDO and STAT-1 expression at 2DAI. Responses were more intense at 4DAI and also included TNF α , Arginase, IRF-1, IL-10, SOCS-1, SOCS-3, and IFN α activation. Liver showed the most intense changes in gene expression in response to Tg infection; PMBC had the lowest changes. High serum haptoglobin and NO levels at 4DAI ($p < 0.01$) confirms inflammatory and macrophage activation. Up-regulation of inflammatory and Th1-dominated markers indicates mechanisms involved in the host defense against acute toxoplasmosis. Supported by ARS funds, and National Pork Board and BRDC grants.

Acute phase proteins in relation to disease and immunity: an avian paradigm

Narayan C. Rath

USDA, ARS, PPPSRU, Fayetteville, Arkansas

Acute phase proteins (APPs) are serum proteins elevated in response to a variety of physiological injuries including infection and inflammation. These are pathogen nonspecific proteins predominantly synthesized in the liver and other tissues such as endothelium, and modulate both innate and acquired immune responses. However, the physiological significance of different APPs is not well understood. In monitoring veterinary health the APPs are used as diagnostic and prognostic indicators. But their use to improve immunity has not been explored. The avian APP response and their significance are much less understood compared to mammals. The object of our study was to understand avian APP response and their possible significance. We induced experimental inflammation in young broiler chickens, by the use of intramuscular injection either of bacterial lipopolysaccharides (LPS), gram positive bacterial cell walls, or croton oil, and compared the serum proteins profiles with sham injected control sera using polyacrylamide gel electrophoresis. Our results showed quantitative changes in several protein bands of which a band corresponding to MW of 65-66 kD was significantly elevated in response to inflammation (1-4). Further analyses using two-dimensional gel electrophoresis and electroblotting onto PVDF membrane, the spots equivalent to 65 kD MW was N-terminal sequenced to yield a molecular homology to ovotransferrin (OVT) (3,4). Using an anti-serum transferrin antibody and western blot, the results further proved that the serum protein was the same as the egg white transferrin. We developed an enzyme-immunoassay to measure OVT concentrations and showed that the serum transferrin levels are elevated in response to a variety of bacterial and viral infections (5). To understand the possible significance of elevated OVT in relation to immunity, we studied its *in vitro* effects on macrophages and heterophil functions under non-stimulated and endotoxin stimulated conditions. Our results show that OVT augmented the functional activity of both of these

cell types as measured using nitric oxide, interleukin-6, matrix metalloproteinase, reactive oxygen production as indicators (6). Additionally, OVT also stimulated heterophil degranulation. These studies provide a paradigm of the possible usefulness of monitoring serum OVT to identify the presence of avian health problems and its function as an immunity modifying protein in poultry.

References:

Xie, H., N.C. Rath, G.R. Huff, W.E. Huff, and J.M. Balog. 2000. Effects of *Salmonella typhimurium* polysaccharide on broiler chicken. Poultry Science 79: 33-40.

Xie, H., N.C. Rath, G.R. Huff, J.M. Balog, and W.E. Huff. 2001. Inflammation-induced changes in serum modulate chicken macrophage function. Veterinary Immunology and Immunopathology 80: 225-235.

Xie, H., N.C. Rath, L. Newberry, G.R. Huff, W.E. Huff, and J.M. Balog. 2001. Ovotransferrin, a major acute phase protein induced by inflammation. In K.A. Schat, editor, Current Progress On Avian Immunology Research: Proceedings of the 6th Avian Immunology Research Group Meeting, p. 302-307.

Xie, H., G.R. Huff, W.E. Huff, J.M. Balog, P. Holt, N.C. Rath. 2002. Identification of ovotransferrin as an acute phase protein in chickens. Poultry Science 81: 112-120.

Xie, H., L. Newberry, F.D. Clark, W.E. Huff, G.R. Huff, J.M. Balog, and N.C. Rath. 2002. Changes in serum ovotransferrin levels in chickens with experimentally induced inflammation and diseases. Avian Diseases 46: 122-131.

Xie, H., G.R. Huff, W.E. Huff, J.M. Balog, and N.C. Rath. 2002. Effects of ovotransferrin on chicken macrophages and heterophil-granulocytes. Developmental and Comparative Immunology 26: 805-815.

In vitro studies of BVDV strains in lymphoid cells lines reveal a third biotype

Julia F. Ridpath and John D. Neill

USDA, ARS, National Animal Disease Center, Ames, Iowa

While bovine viral diarrhea viruses (BVDV) are highly lymphotropic in vivo, in vitro they are studied in epithelial cells. BVDV are classified as cytopathic (cp) or noncytopathic (ncp) based on their activity in cultured epithelial cells. This does not correlate with virulence in acute infections, as the most highly virulent BVDV are ncp. The purpose of this study was to determine if virulence in vivo could be correlated with growth characteristics in cultured lymphoid cells. A BL-3 cell line (derived from a B cell lymphosarcoma) free of BVDV and bovine leukemia virus was used. Cultures were infected with the highly virulent (hv) ncp BVDV2 strain 1373, the low virulent (lv) ncp BVDV2 strain 28508-5 or the lv cp BVDV2 strain 296c. Cell growth, proliferative state, ratio of live to dead cells and virus replication were monitored. Cell morphology was observed by light and electron microscopy. No significant differences were seen between noninfected cells and 28508-5 infected cells. Cytopathic effect (cpe), as defined by cell death, was observed in cells infected with 1373 and 296c. However, the cpe observed with the two strains were different. The cpe associated with 296c infection was apparent within the first 48 hours after infection and was associated with the condensation and fragmentation of most nuclei. Cpe associated with 1373 was not apparent until 96 hours after infection. Cells were non-proliferative but nuclei appeared morphologically normal. The cell cytoplasm became less dense and disorganization of mitochondria was observed. Nuclear condensation and fragmentation became common only in cells in the late stages of cpe. These results suggests that different mechanisms are involved in the cell death associated with lv cp strains and hv ncp virulent strains. Based on these findings BVDV may be grouped into three biotypes, cp (based on cpe in epithelial cells and cultured lymphoid cells within the first 48 hours after infection), ncp (based on lack of cpe in both epithelial and lymphoid cells lines and

lymphocidal (based on cell death in lymphoid cell lines 96 hrs post infection).

Local infection of VSV: pathogenesis, role of inflammatory cells and cytokines

Charles F.C. Scherer^{1,2}, William T. Golde¹, Douglas Gregg¹, D. Mark Estes², and Luis L. Rodriguez¹

¹ USDA, ARS, Plum Island Animal Disease Center, Greenport, New York; ² College of Veterinary Medicine, University of Missouri, Columbia, Missouri

Vesicular Stomatitis Virus (VSV) is negative-sense RNA virus that causes vesicular disease in cattle, horses and pigs. Little is known about the pathogenesis of this virus in livestock species. In order to better understand the pathogenesis, virus distribution and local immune response to VSV infection, 3 bovines were inoculated with the New Jersey strain of vesicular stomatitis virus (VSV-NJ), utilizing a coronary-band scarification procedure. Punch biopsies were obtained at 20 minutes, 6, 12, 24, and 48 hours post-inoculation (hpi). Plasma and esophageal-pharyngeal fluid (OPF) were obtained prior to inoculation and at 24, 48 and 72 hpi. At 72 hours animals were euthanized and various tissues were collected including all major internal organs and various lymphoid tissues. Samples were assayed for virus isolation, presence of viral RNA by real-time RT-PCR and viral antigen identification by immunohistochemistry (IHC). Levels of mRNA for twelve cytokines were measured by real-time RT-PCR in the coronary band. Virus inoculation resulted in fever and vesicular lesions in the coronary bands by 48 hpi in all animals, with virus titers increasing over time from 1 log at 20 minutes to 6 logs TCID₅₀ by 72 hpi. Virus was isolated only from lymph nodes draining the coronary bands (prescapular and popliteal). Virus was also isolated from other lymphoid tissues: tonsil, retropharyngeal, and parotid lymph nodes in one animal and tonsil and axilar lymph node in a second. Virus was never found in blood, OPF samples or in any other lymph node or internal organ tested. Viral antigens were detected predominantly in the stratum spinosum of the coronary band, in a few cells of the basal layer,

and in the afferent and efferent sections of the local lymph nodes, with minimal involvement of germinal centers. Target cells in lymph nodes were morphologically compatible with monocytes. Local cytokine responses were dominated by an IFN alpha, beta, gamma and IL6, all showing increases of 3 to 4 fold by 6 hpi and 15-20 fold by 72 hpi. Levels of IL1 β , IL2, IL4, IL10, IL12p40, IL13, IL15 and TNF alpha maintained basal levels up to 24 hpi but increased 3-25 fold in all animals thereafter. Results demonstrate that infection with VSV-NJ was restricted to the local infected tissue and draining lymphoid tissue. A viral growth curve was observed in coronary band biopsies with increasing titers 12 to 72 hpi. These results provide an important contribution for understanding VSV pathogenesis and transmission in cattle.

Characterizing nematode-induced Th2-dependent changes in intestinal function.

Terez Shea-Donohue¹, Aiping Zhao¹, Kathleen Madden², Motoko Morimoto², William Gause², Dante Zarlenga¹ and Joseph Urban¹

¹USDA, ARS, Nutrient Requirements & Functions Laboratory, Beltsville Human Nutrition Research Center, Beltsville, Maryland; ²US Department of Defense, Uniformed Service University of the Health Services, Bethesda, Maryland

The Th2 cytokines IL-4 and IL-13 are induced by parasitic worm infection and contribute to their expulsion through IL-4 receptor (R)-induced Stat6 activation. IL-4 binds the type 1 and type 2 IL-4R that contain the IL-4R α chain and the common γ chain or the IL-13R α 1 chain, respectively; IL-13 binds to the type 2 IL-4R and to the IL-13R α 2 chain that is not linked to Stat6.

Aim: To determine the effects of intestinal nematodes on intestinal function.

Methods Stripped mucosae were mounted in Ussing chambers to measure tissue permeability and concentration-dependent changes in short circuit current in response to glucose or various secretagogues, and strips of smooth muscle were suspended in organ baths and subjected to stimulation of enteric nerves. Expression of IL-

4R α , IL-13R α 1, and IL-13R α 2 mRNA was analyzed by real time PCR in specific mucosal cells collected by laser capture microdissection.

Results: Infection of mice with 3 different nematodes (*Heligmosomoides polygyrus*, *Nippostrongylus brasiliensis*, and *Trichinella spiralis*) elicits a stereotypic, Stat6-dependent inhibition of glucose absorption and increased mucosal permeability leading to an elevation of fluid in the small intestine; these effects are mimicked by treatment with IL-4 and IL-13 and are Stat6-dependent. Mast cells play a key role in the pro secretory effects of IL-4 that are not shared by IL-13. Nematodes uniformly induces hyper contractility of intestinal smooth muscle by mechanisms that involve both Stat6-dependent and Stat6-independent pathways; IL-13 has more potent effects on contractility than IL-4. Expression of IL-4R α , IL-13R α 1 and IL-13R α 2 mRNA was evident in epithelial and smooth muscle cells. Infection of pigs with the nematode *Ascaris suum* revealed similar Th2-driven changes in intestinal function associated with expulsion of 4th stage larvae.

Conclusion: Nematode infection of the small intestine elicits a characteristic "weep and sweep" response that facilitates worm clearance. IL-4 and IL-13 play a critical role in the host response to nematode infection by altering gut function through receptor-mediated activation of Stat6 pathways. Currently, cloned soluble IL-4R α , IL-13R α 1, and IL-13R α 2 are being used in swine as molecular scavengers to 1) delineate the role of IL-4 and IL-13 in worm clearance and intestinal function; 2) as antagonists and/or agonists of the classic Th2 response and; 3) to mediate IL-13 induced as a model of human allergic disease.

Modulatory effects of IFN- γ , IL-10 and TGF- β in cattle infected with *Mycobacterium paratuberculosis*

J. R. Stabel and M. S. Khalifeh
USDA, ARS, National Animal Disease Center,
Ames, Iowa

Johne's disease is an important disease that results in great economic losses for both dairy and beef production in the United States and worldwide. The disease progresses through distinct stages, a subclinical stage where there are no clinical signs and a clinical stage that is characterized by progressive symptoms associated with chronic shedding of high levels of bacteria in the feces along with severe diarrhea and concomitant weight loss. The host immune response to *M. paratuberculosis* is paradoxical, with strong cell-mediated immune responses during subclinical stages and strong humoral responses during clinical stages of the disease. It is possible that immune modulation of an effective cell-mediated immune response may play an important role in disease progression. We hypothesized that the clinical stage of Johne's disease is mediated by production of cytokines such as TGF- β and IL-10 that downregulate IFN- γ production and interfere with an effective cell-mediated immune response. Therefore, ileum, ileal cecal junction, ileocecal lymph node and mesenteric lymph node tissues from the healthy, subclinical or clinical animals were collected and analyzed for the presence of TGF- β , IL-10 and IFN- γ mRNA by quantitative RT-cPCR. The results show that TGF- β and IL-10 mRNA levels in animals that have progressed to the clinical stage of the disease is higher than that found in subclinical and healthy animals, whereas IFN- γ is higher in subclinical animals. Interferon- γ plays a significant role in the control of mycobacterial infections, including *Mycobacterium paratuberculosis*. However, the contribution of other immunoregulatory cytokines such as IL-10 and TGF- β in Johne's disease has not been investigated as yet. In the present study, we examined the effects of in vivo and in vitro infection with *M. paratuberculosis* on the production of IFN- γ , IL-10 and TGF- β by peripheral blood mononuclear cells (PBMC). We also examined the effects of exogenous IFN- γ , IL-10 and TGF- β on *M. paratuberculosis* survival in the cell cultures. PBMC obtained from naturally infected cows, regardless of their disease status, specifically upregulated IL-10 and TGF- β in culture supernatants in response to stimulation with live *M. paratuberculosis*. Non-stimulated PBMC recovered from subclinically infected

animals secreted the lowest levels of TGF- β but after stimulation with live *M. paratuberculosis* TGF- β levels in the culture supernatants increased to levels similar to that produced by PBMC from healthy animals. Compared to healthy cows, naturally infected animals had higher numbers of viable *M. paratuberculosis* recovered from their cultures after in vitro infection with *M. paratuberculosis*. The addition of exogenous IL-10 and TGF- β to PBMC isolated from healthy cows inhibited the bactericidal activity of these cells as evidenced by the increased number of viable *M. paratuberculosis* recovered from those cultures compared to cell cultures containing medium alone. These data suggest an important immune regulatory role of IL-10 and TGF- β during infection with *M. paratuberculosis* that may be directly related to their effects on macrophage activation and killing of *M. paratuberculosis*.

Bovine T cell response to antigens of *Neospora caninum*.

Wenbin Tuo¹, Ray Fetterer², Pat Boyd¹, Lou Gasbarre³, JP Dubey².

¹ USDA, ARS, Immunology and Disease Resistance Lab, Beltsville, Maryland; ² Animal Parasitic Diseases Lab, Beltsville, Maryland;

³ Bovine Functional Genomics Lab, ANRI, Beltsville, Maryland

See abstract on page 18.

***Ascaris* Infection as a Marker of Local Mucosal Immunity in Pigs.**

Joseph Urban, Ethiopia Beshah, Motoko Morimoto, Gloria Solano-Aguilar, Norberta Schoene, Dante Zarlenga, Ben Mathews, Terez Shea-Donohue, Harry Dawson
USDA, ARS, Beltsville Human Nutrition Research Center, Beltsville, Maryland

Ascaris spp. are ubiquitous in swine with significant economic consequences to the industry, and are the most prevalent helminth infection in man worldwide. The migration of this nematode parasite from the cecum to the liver, lung and small intestine along with its ability to induce immunity and inflammation provides a tool for analysis of local responses at several mucosal surfaces. The Th2-like response to

A. suum in pigs that is characterized by mucosal mast cell and goblet cell hyperplasia, eosinophilia and reagenic antibody production was supported by the demonstration of a pattern of cytokine gene expression consistent with a polarized Th2-related immune response in the lymph nodes draining the sites of infection using a series of 24 gene probes for Th1 and Th2-associated products. Alveolar macrophages obtained from the lungs of *A. suum*-infected pigs had reduced capacity to phagocytise killed *Staphylococcus aureus* but did release enhanced levels of super oxide dismutase indicating a functional skewing of immunity in the lung. Likewise, expulsion of the migrating 4th stage larvae from the jejunum of infected pigs was associated with a "weep and sweep" phenomenon in the small intestine characterized by increased fluid accumulation in the lumen and enhanced smooth muscle contractility. Micro array analysis of parasites that survived expulsion revealed a more robust expression of transcripts for glyceraldehydes 3 phosphate dehydrogenase, succinate dehydrogenase and muscle proteins. Application of sensitive new technologies for evaluation of gene expression of the host and parasite have made the pig a more accurate predictor of immunity in this species and as an attractive model of immune function in humans.

Experimental *Mycobacterium avium* subsp. *paratuberculosis* infection of calves: early induction of a concurrent humoral and cellular immune response.

W. R. Waters^{1*}, J. M. Miller¹, M. V. Palmer¹, J. R. Stabel¹, D. E. Jones², K. A. Koistinen², E. M. Steadham², M. J. Hamilton³, W. C. Davis³, and J. P. Bannantine¹

¹USDA, ARS, National Animal Disease Center, Ames, Iowa; ²Iowa State University, Ames, Iowa;

³Washington State University, Pullman, Washington

Johne's disease (paratuberculosis) of cattle is widespread and causes significant economic losses for producers due to decreased production and poor health of affected animals. The chronic nature of the disease and the lack of a reproducible model of infection hinder research efforts. In the present study, instillation of *Mycobacterium avium* subsp. *paratuberculosis* into the tonsillar crypts of

neonatal calves resulted in chronic infection with peripheral colonization detected by ante-mortem culture of feces and post-mortem (320 d postchallenge) culture of intestinal tissues. Antigen-specific blastogenic, IFN- γ , and nitric oxide responses by blood mononuclear cells from infected calves exceeded prechallenge responses beginning 194 d postchallenge. Upon in vitro stimulation with paratuberculosis antigens, CD4⁺ cells from infected calves proliferated, produced IFN- γ , and increased expression of CD26 and CD45RO (i.e., indicative of an activated memory phenotype). Utilizing a lipoarabinomannan-based ELISA, specific serum immunoglobulin was detected as early as 134 d postchallenge and generally increased after this time point. Two antigens of ~50-kDa and ~60-kDa were particularly immunodominant early in infection as shown by immunoblot with serum collected within 2 weeks postchallenge. Findings indicate that the intratonsillar inoculation route will prove useful as an experimental model for paratuberculosis pathogenesis, diagnosis, and vaccine efficacy studies. Additionally, this study confirms that mycobacteria-specific antibody is detectable early in the course of experimental Johne's disease, even preceding the development of specific cell-mediated responses.

Mechanism for age-associated upregulation in COX-2 expression: role of NF κ B.

Wu D, M. Marko, KJ Claycombe, Eric Paulson, SN Meydani
USDA, ARS, Nutritional Immunology
Laboratory, Boston, Massachusetts

Increased prostaglandin E₂ (PGE₂) production has been shown to contribute to age-associated decline in T cell function as well as dysregulated inflammatory response. We previously demonstrated that macrophages (M ϕ) from old mice have significantly higher cyclooxygenase (COX) activity and COX-2 expression compared to young mice. We further showed that the higher COX-2 expression was due to transcriptional upregulation rather than a change in COX-2 degradation with aging. The purpose of the present study was to determine the mechanism of age-related upregulation in COX-2. We hypothesized that the activation of transcription factors may undergo a change with aging, leading to altered COX-2 transcription. The results demonstrate that LPS-initiated activation of NF κ B, but not AP-1 and CREB, is higher in old than in young mice. No significant age difference was observed in the activity of MAPK P38, JNK, or ERK. Inhibition of NF κ B activation decreased PGE₂ production, COX activity, COX-2 protein and mRNA expression. These reductions were more dramatic in old M ϕ compared to that of young M ϕ . We further showed that old M ϕ have higher I κ B degradation in cytoplasm and p65 translocation to nucleus compared to young M ϕ . Thus the age-associated up regulation of COX-2 transcription in M ϕ is mediated through higher NF κ B activation, which in turn is due to increase in I κ B degradation. Supported by NIA grant # RO1AG09140-09

December 3, 2003

Immunity and Nutrition

Real-time PCR arrays can delineate immunological and nutrition-related gene expression in swine.

Harry Dawson

USDA, ARS, Nutrient Requirements & Functions Laboratory, Beltsville Human Nutrition Research Center, Beltsville, Maryland

See abstract on page 27.

Dietary modulation of host mucosal immunity against avian coccidiosis.

Dalloul¹, R.A., H.S. Lillehoj², and John A. Doerr¹.

¹Department of Animal and Avian Sciences, University of Maryland, College Park, Maryland; ² USDA, ARS, Animal Parasitic Diseases Laboratory, ANRI, Beltsville, Maryland

Infectious diseases constitute one of the greatest threats to the viability of the food animal industry. Commercial broilers, in particular, have a higher risk of acquiring contagious diseases than other livestock due to intensive collective farming practices. For example, avian coccidiosis is a major parasitic disease of significant economic importance and the U.S. industry spends greater than \$800 million annually in medication. In the absence of efficient vaccines to control this disease and the emergence of new antigenic variants of *Eimeria*, the broiler industry has relied upon prophylactic medication. However, anti-coccidial drugs are expensive and their effectiveness is hindered by widespread parasite drug resistance and the high cost of new drug development. Moreover, recent banning of growth promoters and anti-coccidial drugs in commercial animal production in Europe and parts of US will eventually force the industry to eliminate this practice and will require the development of non-chemical methods for coccidiosis control. Thus, probiotics represent an important alternative for antibiotic use in poultry production. However, the exact modes of action in which probiotics achieve such protective roles need to be better elucidated

to improve effectiveness. Previously we have shown that resistance in broilers to *Eimeria acervulina* (EA) was significantly reduced by vitamin A deficiency and enhanced by a probiotic (*Lactobacillus*-based). In the present 2 x 2 factorial study, a broiler starter ration was amended for vitamin A (control, C, or deficient, A) and probiotic status (-, P) to investigate their modulatory effects on the host immune system. Birds were inoculated orally with EA oocysts and disease susceptibility was evaluated by assessing fecal oocyst shedding. Humoral and local cellular mediated immunity were assessed by evaluating antibody and cytokine (IFN- γ and IL-2) levels in sera and intestinal secretions on a 3-day interval following inoculation. Fecal oocyst shedding was highest ($P < 0.05$) in A- birds, followed by AP, C- and CP. Feeding the probiotic reduced shed oocysts by 20% in A fed birds and by 26% in C fed birds. Intestinal IFN- γ was relatively constant in all treatment groups except for A- where it declined steadily and was lower ($P < 0.05$) from day 6 on. Serum IFN- γ levels fluctuated within each treatment and over time were not revealing. Intestinal IL-2 was highest in CP birds at 3 and 9 dpi, and lowest in A- birds at 3, 9, and 12 dpi ($P < 0.05$); no difference between treatments was found at 6 dpi ($P > 0.05$). *Eimeria*-specific intestinal Ab level was constant ($P > 0.05$) in C- birds, but increased with time ($P < 0.05$) in A-, AP and CP birds. Serum Ab levels were also constant in A- and CP, but increased ($P < 0.05$) in C- and AP after 6 dpi. The data demonstrate for the first time a probiotic-enhanced immunity in vitamin A-deficient birds. It is also the first study to demonstrate the probiotic effect on local cell-mediated immunity of chickens, best manifested by apparent lower intestinal invasion and development by EA, based on higher IL-2 secretion and lower EA oocyst production. The new information concerning the immunological mechanism involved in the probiotic-induced immune enhancement would promote the

development of novel ways to modulate the immune response to enhance the health and productivity of poultry.

The effect of vitamin E on secondary bacterial infection following influenza infection in young and old mice

Raina Gay¹, Sung Nim Han¹, Melissa Marko¹, Sarah Belisle¹, Roderick Bronson², Simin N Meydani¹.

¹ USDA, ARS, JMHNRCA, Boston, Massachusetts; ² Tufts University School of Veterinary Medicine, North Grafton, Massachusetts

In the U.S. alone, influenza accounts for over 10,000 deaths annually and over 40,000 deaths during epidemic years. For example, individuals 65 years and older accounted for 89% of all influenza associated deaths in 1992. These deaths were not typically due to influenza infection alone, but from the development of complications due to influenza infection. *Staphylococcus aureus* is commonly associated with influenza infection and can cause sudden and serious illness, especially in the elderly. Previously, we showed that vitamin E supplementation significantly improves the antioxidant status and immune response in aged mice and humans. We further showed that vitamin E supplementation reduces influenza viral infection and improves antioxidant status in aged mice. However, little is known about the potentially deadly bacterial infections that commonly follow influenza infection in the aged. The goal of this study was to determine the efficacy of vitamin E supplementation on secondary bacterial infection following influenza infection in young and old mice. This was tested by feeding young and old C57BL/6 mice semi-synthetic diets containing adequate (30 ppm) or high (500 ppm) levels of vitamin E and determining viral and bacterial titers after primary influenza A or *S. aureus* infection, or following a secondary *S. aureus* infection subsequent to an influenza infection. In this animal model, age did not have a significant effect on *S. aureus* infection alone or *S. aureus* infection following influenza infection. Supplementation with vitamin E did not have a significant effect on *S. aureus* infection

alone. Priming with influenza infection significantly increased *S. aureus* colony counts in the lungs of infected mice fed the control diet. Vitamin E supplementation successfully abolished the priming effect of influenza on *S. aureus* infection. Future studies will determine the mechanism of vitamin E's protective effect.

Supported by a 2002 Glenn/AFAR Scholarship for Research in the Biology of Aging by the American Federation for Aging Research and USDA contract 58-1950-9-001.

Changes in gene expression profiles of T cells by age and vitamin E.

Sung Nim Han¹, Oskar Adolfsson¹, Cheol-Koo Lee², Tomas A Prolla², Jose Ordovas¹, Simin N Meydani¹.

¹USDA, ARS, JMHNRCA, Boston, Massachusetts; ² University of Wisconsin, Madison, Wisconsin

Aging is associated with dysregulation of immune cells particularly that of T cells. Previous studies indicated that vitamin E improves T cell function in part by directly affecting T cells. We studied the gene expression profile of T cells to better understand the underlying mechanisms of age and E-induced changes in T cell functions. Young and old C57BL mice were fed diets containing 30 ppm (control) or 500 ppm (E) of vitamin E for 4 weeks. T cells were purified from splenocytes by negative selection using magnetic beads (anti-Mac-1 and anti-MHC class II), then cultured with media (unstimulated) or plate-bound anti-CD3 and soluble anti-CD28 (stimulated) for 2 hours. Gene expression profile was assessed using microarray analysis. Genes showing more than 2 fold changes, $p < 0.05$, and with at least 1 present call were selected. In unstimulated T cells, 43 genes were expressed significantly higher in old, mainly immunoglobulin genes, while 22 genes were expressed significantly higher in young including T cell receptor related genes. Upon stimulation, 103 genes were upregulated in at least one age or diet group (16 genes in all age and diet groups). Response to stimulation was significantly affected by age (40 genes) and vitamin E (24 genes). Significantly lower expression of T receptor alpha chain and factors involved in the signal

transduction pathways were observed with aging. Significantly higher expression of suppressor of cytokine signaling (SOCS) 3 and lower expression of growth factor independent (gfi) 1 were observed with aging. Overexpression of SOCS3 has been shown to suppress T cell proliferation and IL-2 production and gfi1 has been reported to down regulate SOCS3. Expressions of IL-2 increased significantly in both young and old T cells while that of IL-4 decreased in old T cells by vitamin E. These findings suggest that: 1) aging has significant effects on signal transduction pathways in T cells, and 2) vitamin E has significant effect on Th1/2 balance.

Supported by NIA grant #RO1AG09140-09, USDA contract 58-1950-9-001, and gifts from Roche, Inc.

Stress-related chronic disease in turkeys and the effects of nutritional and environmental immunomodulation.

Gerry Huff, Bill Huff, Narayan Rath, and Janice Balog

USDA, ARS, Poultry Production and Products Safety Research, Fayetteville, Arkansas

Our research using the synthetic glucocorticoid, dexamethasone (DEX), to suppress the immune response of turkeys to low doses of opportunistic bacteria has led to the development of a number of nutritional and environmental strategies for decreasing the impact of stress-induced immunosuppression. The immunosuppressive effects of stress appear to be greater in male turkeys than in females. We have found that female turkeys are more resistant to colibacillosis and respiratory disease in a DEX-*E.coli* challenge infection than are males. The anti-bacterial activity of monocytes from male and female turkeys treated with 0.5 or 2.0 mg/Kg DEX was measured by monitoring the percentage of infected cells over time. Antibacterial activity was significantly decreased at both concentrations of DEX at both 8 and 16 hours post-infection in both sexes combined and was significantly lower in males as compared to females, suggesting that stress can reduce the ability of turkey mononuclear cells to kill bacteria, and that the difference in this function may be relevant to the

sex-related difference in disease resistance. Water supplementation with vitamin D₃, vitamin E, sodium salicylate, and β -1,3/1,6-glucan, a helical polysaccharide derived from the cell wall of *Saccharomyces cerevisiae*, have all shown immunomodulatory effects on male turkeys challenged with *E. coli* in our model. We have shown that excess handling or environmental enrichment during the first 2 weeks after hatch can lead to decreased resistance later in life. Our model has further suggested that a small population of male turkeys is highly susceptible to the immunosuppressive effects of stress, and that these individuals can be identified by their performance in a behavioral test utilizing a T-maze at 3 days of age. These individuals are also more susceptible to the immunosuppressive effects of excess environmental stimulation during the first two weeks after hatch. These studies illustrate the dramatic effects that stress can have on both production values and resistance to opportunistic infection, and suggest that a combination of genetic selection, nutritional immunomodulation, and environmental strategies to reduce stress may lead to healthier birds and a safer and a more wholesome product.

Purified β -glucan as an abiotic feed additive up-regulates heterophil function in immature chickens.

V.K. Lowry^{1*}, M.B. Farnell², P.J. Ferro¹, A. Bahl³, and M.H. Kogut⁴

¹Texas A&M University, Departments of Anatomy and Public Health & Poultry Science, respectively, College Station, Texas; ²Institutes for Animal Health, Compton, Berkshire RG20, 7NN, UK; ³ImmuDyne, Inc., Houston, Texas; ⁴USDA, ARS,SPARC, College Station, Texas

Functionally, the innate immune system of immature chickens is inefficient during the first week post-hatch. Pathogens, such as *Salmonella enterica* serovar *Enteritidis* (SE), are able to invade various tissues due to this immunological inefficiency of the immature chicken. The objective of the present study was to evaluate the effect of purified β -glucan as an immunomodulator of the innate immune response, specifically the heterophil. The functional

efficiency of heterophils isolated from immature chickens fed a β -glucan ration was significantly ($P<0.05$) up-regulated when compared to heterophils isolated from chickens fed a control ration as determined with an array of functional assays. Phagocytosis, bactericidal killing and oxidative burst were significantly increased in heterophils isolated from chickens fed the purified β -glucan ration ($P<0.05$). To our knowledge this is the first report of a purified β -glucan feed additive significantly up-regulating the functional abilities of heterophils isolated from immature chickens against an invading pathogen, SE.

Altered structure, function and allergenicity of peanut proteins due to processing

Soheila J. Maleki

USDA, ARS, Food Processing and Sensory Quality Research Unit, Southern Regional Research Center, New Orleans, Louisiana
sjmaleki@srrc.ars.usda.gov

There has been a dramatic increase in the prevalence of IgE mediated allergy over the past decade, which is one of the reasons that it is viewed as an area of increasing importance. During this time a significant, growing, and costly, portion of the FDA's food recalls have been due to mislabeled products that contain food allergens. Meanwhile, little is known about the reason why certain foods contain allergens and others are considered hypoallergenic. Some classic characteristics associated with food allergens are resistance to enzymatic digestion and heat, stimulation of T cell proliferation, IgE binding and abundance in food. Certain enzymatic functions have also been associated with allergenicity of proteins. However, not much is known about what happens to the allergenicity of food products after processing. In this study, roasted whole peanut proteins were shown to be less soluble, more resistant to digestive enzymes and bound higher levels of IgE than raw whole peanut proteins. Mice were more likely to be sensitized when treated with roasted peanuts than with raw peanuts and had higher vascular permeability when skin tested with roasted peanuts. The major peanut allergens Ara h 1 and Ara h 2 were then purified

from both raw and roasted peanuts and examined at a molecular level for alterations that may explain the *in vivo* observations. Following roasting, Ara h 1 was shown to form irreversible trimers that are extremely resistant to digestion and bind higher levels of IgE than Ara h 1 from raw peanuts. The amino acid sequence of a second major peanut allergen, Ara h 2 was found to be homologous to the family of trypsin/alpha-amylase inhibitors. Most significantly, while the secondary structure remained unaltered, Ara h 2, purified from roasted peanuts was several fold more active as a trypsin inhibitor and bound higher levels of IgE than the Ara h 2 purified from raw peanuts. Our findings suggested that the structural and functional changes due to food processing contribute to increased allergenic properties of peanut proteins and may indeed enhance the possibility of becoming sensitized to peanuts.

Interpretive summary

It is not known why certain foods are allergenic and others are not. Some classic characteristics associated with food allergens are resistance to digestion and heat, stimulation of T cell proliferation, IgE binding and abundance in a food. Certain enzymatic functions have also been associated with allergenicity of proteins. However, not much is known about what happens to the allergenicity of food products after processing. Here the consequences of roasting on the structural, functional, immunological and allergenic properties of peanut proteins were examined. Roasting was found to alter the structure and function of peanut allergens Ara h 1 and Ara h 2. In addition, whole peanut proteins from roasted peanuts were more resistant to degradation by digestive enzymes and more likely to immunize mice than raw peanut proteins. Our findings suggested that the structural and functional changes caused by food processing contribute to increased allergenic properties of peanut proteins and may indeed enhance the possibility of becoming sensitized to peanuts.

Aging, immune response, and infectious disease: molecular mechanisms and reversal by nutrient intervention.

Simin Nikbin Meydani, Dayong Wu, Sung Nim Han, Raina T. Gay

USDA, ARS, Nutritional Immunology Laboratory, Jean Mayer Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts

See abstract on page 25.

Nutritional plane affects antigen-induced proliferation of lymphocyte subsets in milk replacer-fed calves vaccinated with *M. bovis* bacillus Calmette-Guerin (BCG).

B.J. Nonnecke¹, M.R. Foote², M.A. Fowler³, B.L. Miller³, and W.R. Waters⁴

¹USDA, ARS, Periparturient Diseases of Cattle Research Unit, NADC, Ames, Iowa; ²Iowa State University, Ames, Iowa; ³Land O' Lakes, Inc., Webster City, Iowa; ⁴USDA, ARS, Bacterial Diseases of Livestock Research Unit, NADC, Ames, Iowa

See abstract on page 26.

Leukocyte migration and metabolism in adipose tissue of leukocyte adhesion receptor deficient mice.

Rebecca Robker, Robert Collins, Nancy Butte, Harry Mersmann, and C. Wayne Smith
Section of Leukocyte Biology, Children's Nutrition Research Center, Department of Pediatrics, Baylor College of Medicine, Houston, Texas

ICAM-1 (CD54) and Mac-1 (CD11b) mediate the migration of leukocytes into tissues where they influence tissue function. Our studies sought to determine whether ICAM-1 and Mac-1 have an intrinsic role within adipose tissue by 1) analyzing the expression of ICAM-1 in adipose tissue, 2) identifying leukocyte populations within adipose tissue, and 3) determining whether ICAM-1 and Mac-1 mutant mice exhibit abnormal numbers of adipose tissue leukocytes or altered metabolism. In wildtype mice ICAM-1 was expressed in adipose tissue and localized to the vascular endothelium. ICAM-1 mRNA was detected in

isolated adipocytes, but surface staining of the adipocyte was not evident, though cultured adipocyte supernate was positive for soluble ICAM-1. ICAM-1 and Mac-1 do not appear to be essential, however, for adipose tissue homeostasis since ICAM-1 and Mac-1 mutant mice exhibited no alterations in body fat deposition or adipocyte metabolism. Macrophages were observed to be a prevalent cell type within the stromal-vascular cell fraction of adipose tissue, and adipose tissue and plasma from obese mice exhibited significant elevations in MCP-1. Macrophage numbers in adhesion molecule deficient mice did not differ from controls. However, gender-specific differences were observed, with adipose tissue from female mice containing significantly more macrophages than males. Our results are in contract to a previous report of increased obesity in ICAM-1 and Mac-1 mutant mice, but demonstrate leukocyte populations within adipose tissue which must be considered in light of increasing evidence of immune-modulation of body weight.

The effect of human-derived probiotic bacteria on the immune and intestinal function of pigs

Gloria Solano-Aguilar*¹, Harry Dawson¹, Tonya Ledbetter¹, Terez Shea-Donohue¹, Norberta Schoene¹, Jeff Call², Ethiopia Beshah³, William Hare³ and Joseph Urban¹.

¹USDA, ARS, Beltsville Human Nutrition Research Center, Beltsville Maryland; ²USDA, ARS, Eastern Regional Research Center, Wyndmoor, Pennsylvania; ³USDA, ARS, Animal & Natural Resources Institute, Beltsville, Maryland
See abstract on page 24.

Vitamin A deficiency decreases development of Th1 memory cells following adoptive transfer of TCR-transgenic T cells.

Charles B. Stephensen¹, Xiaowen Jiang¹, and Tammy Freytag
¹USDA, ARS, Western Human Nutrition Research Center, Davis, California

Vitamin A deficiency enhances production of Th1 cytokines, particularly IFN-(, during the memory and effector phases of the immune response. This

had led to the presumption that vitamin A deficiency biases T-helper cell development in the same direction. To test this hypothesis we adoptively transferred naïve T cells from DO11.10 TCR-transgenic mice to vitamin A-deficient or control BALB/c recipients. Recipients were then immunized with the cognate peptide antigen for the TCR-transgenic T cells (OVA323-339) and the Th1/Th2 phenotype of the transferred cells characterized by intracellular cytokine staining following in vitro stimulation with phorbol myristate acetate and ionomycin. In contrast to expectations, the percentages of IFN- γ -positive cells ($9.0 \pm 0.5\%$; mean \pm standard error) and IL-2-positive cells ($39.2 \pm 2.3\%$) in draining lymph

nodes of deficient mice were both lower than in control mice ($11.3 \pm 0.4\%$, $p = 0.002$ and $46.4 \pm 2.1\%$, $p = 0.028$, respectively). In addition, the percentage of IL-10 positive cells was higher in vitamin A-deficient mice ($3.71 \pm 0.24\%$) than it was in control mice ($1.74 \pm 0.22\%$; $p < 0.001$). IFN- γ and IL-2 concentrations were also lower in supernatants from splenocyte cultures restimulated with OVA323-339 peptide antigen. Thus vitamin A deficiency at the time of initial antigen exposure diminishes the development of Th1 memory cells and enhances the development of IL-10-producing Th2 memory cells.

Immunity and Disease Resistance

Relationship of the probability of disease incidence with immunity

Gamal A. Abdel-Azim^{1*}, Albert E. Freeman¹, M.E. Kehrli, Jr.², S.C. Kelm¹, J.L. Burton³, A.L. Kuck¹, and S. Schnell

¹ Department of Animal Sciences, Iowa State University, Ames, Iowa; ²USDA, ARS, Metabolic Diseases and Immunology Research Unit, National Animal Disease Center, Ames, Iowa; ³ Michigan State University, Department of Animal Science, Immunogenetics Laboratory and Center for Animal Functional Genomics East Lansing, Michigan; * Current address: Cooperative Resources International, Shawano, Wisconsin

Health data in Holstein cows sired by a selected group of sires were analyzed as categorical traits by a threshold model. Data were collected from 174 herds and included incidences on 18,000 daughters of 1365 sires. Breeding values and genetic parameters for disease incidence were estimated. Probability of disease incidence in daughters as values associated with their sires were computed. The immune competence of the sires had been previously tested in a model of glucocorticoid immunosuppression. Statistical relationships between daughter health and immune response traits in sires were investigated. Heritability estimated by the threshold model was 0.277, an estimate sufficiently high to indicate the great potential of selection for disease resistance. Evidence of association between probability of disease incidence and seven immune response traits was found, which motivates and justifies the indirect selection of dairy sires for disease resistance. Indirect selection for disease resistance saves time and money, and could serve as a first guide for what bulls to select for health until young bulls have been progeny tested. Further research is warranted to develop indirect models of selection for disease resistance.

Correlations between disease resistance, pathogen levels, and clearance rate in channel catfish

Lanie Bilodeau¹, Brian Small¹, David Wise², and William Wolters¹

¹USDA-ARS, Catfish Genetics Research Unit, Stoneville, Mississippi; ²Mississippi State University, MAFES, Stoneville, Mississippi

Enteric septicemia of catfish (ESC) is the most prevalent disease affecting commercial catfish farms. The USDA-ARS Catfish Genetic Research Unit has an ongoing program for genetic improvement of disease resistance amongst other economically important traits. Susceptibility to the causative agent of ESC, the bacterium *Edwardsiella ictaluri*, appears to be consistent within each family/spawn of NWAC103 channel catfish.

Real-time PCR technology was utilized to measure differences in bacterial loads and clearance rates for 6 ESC-susceptible and resistant families of channel catfish during immersion challenge. Fish from resistant families had increased survivorship during the challenge when compared to susceptible families. Significant differences ($p < 0.05$) in the quantity of bacterial DNA between resistant and susceptible families were evident for both blood and spleen tissue 5 days following exposure to *E. ictaluri*. Mean quantities of bacterial cell equivalents per 100 μ L of blood at 5 days post-exposure were $2.84 \times 10^5 \pm 143123$ for the susceptible families and 496 ± 455 for the resistant families. Significant differences in spleen tissue also occurred on day 12, with fish from the resistant families having higher ($p < 0.05$) levels of bacteria than fish from the susceptible families. Overall, families that are susceptible to ESC carried higher levels of bacterial DNA in their blood than resistant families.

Pathogen clearance differed between the two sets of families. Clearance was evident by day 12 in

fish from the susceptible families and bacterial levels continued to decrease throughout the remainder of the trial. However, no significant clearance was evident in fish from the resistant families. This may be due to chronic low levels of infection that did not trigger a clearance response from the immune system. The innate immune system may suppress infection only in resistant families, preventing an acute host-response whereas in susceptible families, acute infection occurs and may secondarily be cleared. Clearance rate does not appear to significantly affect mortality.

Heterophils isolated from chickens resistant to extraintestinal *Salmonella enteritidis* infection express higher levels of pro-inflammatory cytokine mRNA following infection than heterophils from susceptible chickens

Pamela J. Ferro¹, Christina L. Swaggerty², Pete Kaiser³, Igal Y. Pevzner⁴, and Michael H. Kogut²
¹Texas A&M University, Poultry Science, College Station, Texas; ²USDA-ARS, SPARC, 2881 F&B Road, College Station, Texas; ³Institute for Animal Health, Compton, Berkshire, UK; ⁴Cobb-Vantress Inc., Siloam Springs, Arkansas
See abstract on page 29.

Genetic and immunologic basis for resistance to GI nematode infections

Louis C. Gasbarre, Tad Sonstegard, Ricardo Araujo, and Curt Van Tassell
USDA, ARS, Bovine Functional Genomics Laboratory, ANRI, Beltsville, Maryland
See abstract on page 28.

High-throughput molecular approaches to identify coccidiosis disease resistance genes and protective immunological mechanisms in avian coccidiosis.

Hyun S. Lillehoj¹, Wongi Min¹, Benjamin F. Matthews², and Marlene Emara³
¹ USDA, ARS, Animal Parasitic Diseases Laboratory, Beltsville, Maryland; ² USDA, ARS, Soybean Genomics Improvement Laboratory, Beltsville, Maryland; ³Department of Animal and

Food Sciences, University of Delaware, Newark, Delaware

See abstract on page 30.

Structural and functional characterization of chicken IL-15, IL-16 and IL-17 cDNAs.

Wongi Min and Hyun S. Lillehoj
USDA, ARS, Animal Parasite Diseases Laboratory, ANRI, Beltsville, Maryland

Limited progress in chicken cytokine immunobiology has slowed down immunological studies in poultry diseases. From our recently developed chicken intestinal gene database which contains >38,000 expressed sequence tags (EST), we searched for novel genes which are associated with innate and acquired immunities. This report describes the cloning and characterization of cDNAs encoding IL-15, IL-16 and IL-17 from our chicken intestinal library. DNA sequence analysis of a chicken IL-15 cDNA identified a 187 amino acid open reading frame encoding a protein with a predicted molecular weight of 21,964, 2 potential N-linked glycosylation sites, 4 highly conserved Cys residues, 2 out-of-frame AUG initiation codons in the 5' untranslated region, and an unusually long (66 amino acid) signal peptide such that the expected size of the mature protein is 14,462 Daltons. Chicken IL-15 and IL-2 were compared with regards to their molecular, cellular, and functional characteristics. The predicted amino acid sequences of both chicken cytokines showed greater homologies with mammalian IL-15s compared with mammalian IL-2s. Northern hybridization and RT-PCR demonstrated chicken IL-15 gene transcripts in a wide variety of tissues and cell types while the chicken IL-2 gene was expressed only in concanavalin A-activated spleen cells. Both recombinant cytokines stimulated the growth of spleen T cells and enhanced the activity of natural killer cells *in vitro*. Subcutaneous injection with an expression plasmid encoding IL-15 increased the percentage of CD3+ spleen T lymphocytes whereas injection of an IL-2 cDNA augmented CD3+, CD4+, CD8+, TCR1+, and TCR2+ T cells. Collectively, these results indicate that chicken IL-15 and IL-2 are T cell growth factors potentially capable of enhancing cell-mediated immunity *in vivo*.

IL-16 and IL-17 are proinflammatory cytokines produced by activated lymphocyte cells. Chicken IL-16 gene contained the entire open reading frame (ORF) of pro-IL-16 and encoded the mature IL-16 protein which consists of 607 amino acids. Chicken IL-16 showed 86% sequence homology to duck pro-IL-16 and 49-52% to various mammalian IL-16s. By Northern blot analysis, IL-16 transcripts showed a restricted expression to the lymphoid tissues. Recombinant chicken IL-16 consisting of 149 C-terminal amino acids of pro-IL-16 was biologically active when expressed in COS-7 cells and showed chemoattractant property for lymphocytes. IL-17 contained a 507 bp open reading frame predicted to encode a protein of 169 amino acids with a molecular mass of 18.9 kDa. Chicken IL-17 shared 37%-46% amino acid sequence identity with mammalian IL-17 and homologous to the ORF 13 of *Herpesvirus saimiri*. By Northern blot analysis, IL-17 transcripts were identified in a reticuloendotheliosis virus-transformed chicken lymphoblast cell line (CU205) and Con A-stimulated splenic lymphocytes, but not Cu91, RP9, RP13 cell lines or kidney, bursa, heart, spleen, cecal tonsils and thymus. Conditioned medium from COS-7 cells transfected with chicken IL-17 cDNA induced the production of IL-6 by embryonic fibroblasts. These studies will facilitate our understanding of poultry immune system.

Differential pro-inflammatory cytokine mRNA expression in heterophils isolated from *Salmonella*-resistant and -susceptible chickens

Christina L. Swaggerty¹, Pamela J. Ferro², Lisa Rothwell³, Igal Y. Pevzner⁴, Michael H. Kogut¹, and Pete Kaiser³

¹ USDA, ARS, SPARC, College Station, Texas; ² Texas A&M University, College Station, Texas; ³

Institute for Animal Health, Compton, Berkshire, UK; ⁴ Cobb – Vantress, Inc., Siloam Springs, Arkansas

We have conducted studies utilizing a parental pair of broiler chickens (lines A and B) and the F1 reciprocal crosses (C and D). Previously we showed increased *in vitro* heterophil functional efficiency translates to increased *in vivo* resistance against a systemic *Salmonella enteritidis* (SE) infection. Heterophils are capable of producing cytokines and are known to modulate acute protection against *Salmonella* in young poultry. We hypothesize that heterophils from SE-resistant chickens (A and D) have an upregulated pro-inflammatory cytokine response compared to heterophils from SE-susceptible chickens (B and C). In this study, heterophils were isolated from day-old chicks, stimulated with SE or SE opsonized with either normal chicken serum or immune serum against SE, and quantitative real-time RT-PCR using TaqMan chemistry was used to ascertain levels of cytokine mRNA expression. Heterophils from SE-resistant chicks (A and D) had significantly higher mRNA expression levels of pro-inflammatory cytokines (IL-6, IL-8, and IL-18) upon treatment with all stimulants compared to heterophils from SE-susceptible lines (B and C). Further, heterophils from SE-resistant chicks had significantly decreased mRNA expression levels of TGF- β 4, an anti-inflammatory cytokine, compared to heterophils from SE-susceptible chicks. These data indicate that chickens with heterophils more able to mount a pro-inflammatory cytokine response may determine if the chickens are going to be resistant or susceptible to *Salmonella* infections. Therefore heterophil functional efficiency and cytokine production may be useful biomarkers for poultry breeders to consider when developing new immunocompetent lines of birds.

Immunointervention Strategies

Immunopotential of a recombinant adenovirus-FMD vaccine by IFN α in swine

Sonia A. Botton, Mario C.S. Brum, Elida M. Bautista, Marla J. Koster, William T. Golde, Marvin J. Grubman*
USDA, ARS, Plum Island Animal Disease Center, Greenport, New York
*e-mail address: mgrubman@piadc.ars.usda.gov

The adjuvant effect of interferon alpha (IFN α) was examined in swine vaccinated with a recombinant replication-defective adenovirus containing foot-and-mouth disease virus (FMDV) capsid (P1-2A) and 3C proteinase coding regions (Ad5-A24). Swine were divided into 5 groups and inoculated with high (5×10^9 PFU) or low (5×10^8 PFU) doses of Ad5-A24 in the presence or absence of porcine IFN α (Ad5pIFN α , 10^9 PFU). Control animals received 6×10^9 PFU of adenovirus containing the glycoprotein from vesicular stomatitis virus (VSV) (Ad5VSV-G). All swine were challenged at 42 days post vaccination (dpv) with FMDV-A24. Prior to challenge, blood samples were examined for IFN production, induction of IFN-induced genes and FMDV-specific neutralizing antibodies (PRN $_{70}$). After challenge a number of disease parameters were analyzed including clinical score, viremia and antibodies against viral structural and nonstructural (NS) proteins.

Preliminary results indicate that both high-dose Ad5-A24 inoculated groups developed significant PRN $_{70}$ by 14-21 dpv which was maintained until the day of challenge. While both low-dose inoculated groups had lower PRN $_{70}$, the IFN group had an enhanced response. After challenge all control animals developed early viremia, vesicular lesions, and antibodies to FMDV NS proteins and high-levels of PRN $_{70}$. Animals receiving low dose Ad5-A24 with no IFN had similar clinical signs, except that fewer animals developed viremia. In contrast pigs inoculated with the same dose plus IFN had a delayed onset of vesicular lesions and only one animal had detectable viremia. Animals

vaccinated with high dose Ad5-A24 without IFN had no viremia, significantly fewer lesions and delayed onset of disease compared to the group given a low dose and no IFN. Four of five pigs vaccinated with high dose plus IFN were completely protected from disease and only one animal in this group had a mild vesicular lesion restricted to the site of inoculation of challenge virus. Our results indicate that IFN α enhances the level of protection induced by the adenovirus-FMD vaccine against homologous FMDV, supporting the use of IFN α as a potential adjuvant in FMD vaccination strategies.

Swine vaccine trials with replication-defective human adenovirus serotype 5 containing the capsid and 3C proteinase coding regions of Foot-and-Mouth Disease Virus O1 Campos in the presence or absence Ad5-porcineGM-CSF

Luizinho Caron, Mario C.S. Brum, Mauro P. Moraes, Clarice W. Arns, William T. Golde and Marvin J. Grubman*.
USDA, ARS, Plum Island Animal Disease Center, Greenport, New York
*e-mail address: mgrubman@piadc.ars.usda.gov

Granulocyte-macrophage colony stimulating factor (GM-CSF) has been widely used to successfully stimulate the immune response in vaccine trials involving HIV, hepatitis C and B, cancer etc. Because of these promising results, we constructed a recombinant-defective human adenovirus 5 (Ad5) containing porcine GM-CSF gene (Ad5-pGMCSF) to examine its effect on an experimental FMDV serotype O1 Campos vaccine that is poorly immunogenic. The recombinant virus was grown in 293 cells, cesium chloride purified and expression verified in IBRS2 cells by radio-immunoprecipitation and Western blotting. The activity of pGM-CSF expressed in IBRS2 cells was demonstrated in TF-1 cells, a cell line dependent on GM-CSF for growth.

Eighteen pigs were divided into 5 groups: Groups 1 and 2, containing 3 animals each, were vaccinated with control virus, Ad5-Blue, or Ad5-pGMCSF and three other groups, 4 animals per group, were vaccinated with Ad5 containing the capsid (P1-2A) and 3C proteinase coding regions of FMDV O1 Campos (Ad5-O1C) in the absence or presence of 2 different amounts of Ad5-pGMCSF. All animals were challenged 21 days post-vaccination with a field strain of homologous virus. The results indicate that the presence of pGMCSF induced a more rapid FMDV-specific neutralizing antibody response but the titers declined faster than in the group given vaccine alone. Furthermore, after challenge, the groups administered both Ad5-pGMCSF and Ad5-O1C developed somewhat more severe disease when compared with the group given only Ad5-O1C.

Development of an oral vaccine for *Escherichia coli* O157:h7

Evelyn A. Dean-Nystrom
USDA, ARS, National Animal Disease Center,
Ames, Iowa
See abstract on page 31.

Oral dosing of mice with *Escherichia coli* expressing recombinant interferon gamma reduces *Cryptosporidium parvum* infection.

James A. Harp, Joan C. Oesper, and Thomas A. Casey
National Animal Disease Center, USDA-ARS,
Ames, Iowa

Cryptosporidium parvum is a protozoan parasite that causes intestinal infection in numerous mammalian species. Previous studies indicate that interferon gamma is critical to resistance to and recovery from *C. parvum* infection. We electroporated a gene for recombinant murine interferon gamma (rIFN) into a nontoxigenic *E. coli* strain that expresses the F41 adhesion pilus, allowing it to adhere in the mammalian small intestine (strain 4907.2.1). Groups of mice were treated at 1 day of age with an oral dose of 4907.2.1 expressing rIFN, either alone, or in combination with *Lactobacillus brevis* and/or killed *C. parvum*. Controls received phosphate buffered saline. All mice were orally inoculated

with 103 viable *C. parvum* oocysts at 7 days of age. Mice were killed at 2 weeks of age, and colon contents examined microscopically for the presence of *C. parvum* oocysts. The number of oocysts seen per microscopic field was compared between groups. Control mice had a mean of 2.1 oocysts per field. Mice receiving *L. brevis*, killed *C. parvum*, and 4907.2.1 had a mean of 1.75 oocysts per field. Mice receiving killed *C. parvum* and 4907.2.1 had a mean of 1.3 oocysts per field. Mice receiving 4907.2.1 alone had a mean of 1 oocyst per field. While these differences were not statistically significant, there was a clear trend that mice receiving *E. coli* expressing rIFN were less heavily infected with *C. parvum*.

Stimulation of chicken leukocytes and reduction of *Salmonella* organ invasion in neonatal chicken by immunostimulatory CpG-oligodeoxynucleotide.

Haiqi He and Michael H. Kogut
USDA, ARS, Southern Plains Agricultural
Research Center, College Station, Texas
See abstract on page 32.

Development of an inactivated Newcastle disease virosome vaccine that protects against challenge from velogenic Texas GB

Darrell R. Kapczynski and Terrence M. Tumpey
USDA-ARS, Southeast Poultry Research
Laboratory, Athens, Georgia

While outbreaks of highly virulent Newcastle disease virus (NDV) are a major concern to the poultry industry internationally, economic losses caused by low virulent (lentogenic) strains continue to result from decreased egg production in layers and airsacculitis in broilers. Since lentogenic strains of NDV are widely used as vaccine strains, it is possible that these viruses, may be directly or indirectly responsible for some of the economic losses. In an effort to examine protection against NDV, we developed a non-replicating virosome vaccine that retains the ability to hemagglutinate chicken red blood cells and fuse with chicken embryo cells. Preliminary vaccine efficacy studies indicate that two intranasal doses containing 10 µg of virosomes

protects chickens against lethal challenge from velogenic Texas GB.

Development of immunotherapeutic recombinant chicken scFv antibodies reactive with an apical antigens of *Eimeria*.

Hyun Lillehoj¹, Wongi Min¹, and Jin K. Kim²

¹USDA, ARS, Animal Parasite Diseases Laboratory, ANRI, Beltsville, Maryland;

²Department of Microbiology, Changwon National University, Changwon, Korea.

Slow progress in the identification of protective *Eimeria* antigens hinders the development of control strategies against avian coccidiosis. In order to identify the epitopes of *Eimeria* which are recognized by chicken B cells, novel immunization strategy was used that led to the development of several chicken monoclonal antibodies which blocked sporozoite invasion of host cells. While most antibodies recognized uncharacterized parasite surface proteins, two were identified with reactivity against apical complex proteins located at the anterior tip of sporozoites. To express high concentration of these antibodies for potential application as immunotherapeutic reagents, a single chain fragment variable (ScFv) region gene was constructed by PCR amplification of the variable heavy (VH) and light (VL) chain genes with a flexible linker. By indirect immunofluorescence, recombinant ScFv antibody expressed in *E. coli* showed identical binding specificity as the original monoclonal antibody against *E. acervulina* and *E. tenella* sporozoites, staining the apical complex and thus demonstrating reactivity against native parasite protein. These results demonstrate that recombinant 6D12 ScFv recognizes the native conoid antigen on *Eimeria*. Potential application of this antibody in prophylactic therapy against avian coccidiosis is being explored using various delivery strategies including embryo injection.

Adjuvant Effect of IL-1 β , IL-2, IL-8, IL-15, IFN- α , IFN- γ , TGF- β 4 and lymphotactin on DNA vaccination against *Eimeria acervulina*.

Wongi Min¹, Hyun S. Lillehoj¹, and Peter

Stacheli².

¹ USDA, ARS, Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Beltsville, Maryland; ²Department of Virology, University of Freiburg, D-79008, Freiburg, Germany.

See abstract on page 32.

Comparison of immunologic responses to *Brucella* and *Mycobacterium* vaccines across species

S.C. Olsen, A. Maue, M.V. Palmer, W.S. Stoffegren, and W.R. Waters.

USDA, ARS, Bacterial Diseases of Livestock Research Unit, National Animal Disease Center, Ames, Iowa

Eradication of brucellosis and tuberculosis from livestock populations is a high priority for regulatory personnel. Persistence of these diseases in wildlife reservoirs pose a substantial risk for transmission to livestock. In a series of studies conducted at the National Animal Disease Center, immunologic responses of wildlife species of interest to brucellosis and/or mycobacterium vaccines were characterized and compared to responses of cattle. As cell-mediated immunity is critical for long-term protection against diseases caused by intracellular bacteria, our studies particularly focused on proliferative responses of peripheral blood mononuclear cells (PBMC), nitric oxide production, γ -interferon production, and flow cytometric analysis of proliferating PBMC subsets. Our data suggest that immunologic responses to *Brucella* or *Mycobacterium* vaccines may differ by species. Species that are considered to be closely related may demonstrate significant differences in immunologic responses to vaccination. Our data also suggest that immunologic responses to vaccination cannot be predicted but must be evaluated in the species of interest. Lastly, some wildlife species may have value as models for characterizing the role of humoral responses in ruminants as their immunologic responses to *Brucella* and *Mycobacterium* vaccines include robust antibody and deficient cell-mediated responses.

Does antigenic drift occur with avian influenza in poultry?

David Suarez and Chang-Won Lee
USDA, ARS, Southeast Poultry Research
Laboratory, Athens, Georgia

Avian influenza in poultry and human influenza are both type A influenza viruses and both originated from an influenza reservoir found in wild birds, including ducks and gulls. Human influenza has been well characterized for antigenic drift, which is the accumulation of point mutations in the viral genome. These changes in the hemagglutinin gene can slowly create antibody escape mutants that make current vaccines less effective. Therefore the strains of virus in these killed vaccines must be updated on a yearly basis to provide optimal protection against the prevailing strains of circulating virus. In general when the predicted circulating strain of virus elicits an antibody titer with a fourfold loss in hemagglutination inhibition (HI) cross neutralization activity to the current vaccine, then the vaccine is changed. The changes in nucleotide similarity between the old and new vaccine strains are often less than 2%. For highly pathogenic avian influenza (HPAI), antigenic drift as it affects vaccine selection has not been a consideration in the past. Vaccine strains with differences of over 15% to the challenge strain can still provide good protection from clinical disease. There is a correlation with nucleotide sequence similarity and viral shed, with the closer a challenge strain is to the vaccine the greater reduction in viral shed. The disconnect between human influenza and HPAI vaccines may be that human influenza is a mucosal infection, but HPAI is a mucosal and a systemic infection. Vaccines for influenza do not prevent infection, but they do reduce viral shed and prevent or lessen clinical signs. Antibodies in the blood stream may effectively prevent the systemic phase of disease with HPAI viruses. However, recent studies of low pathogenic avian influenza from Mexico and the U.S. may present a different view of antigenic drift in poultry. In Mexico H5N2 LPAI has circulated for 9 years and vaccination with two different vaccines have been used widely. When recent isolates are examined by HI cross neutralization to the killed vaccine,

large decreases in titers are observed. However, when examining H7N2 LPAI viruses that have been circulating in the U.S. since 1994, large decreases in HI cross neutralization titers are not observed. In both cases nucleotide sequence differences between 3-5% are observed. One major difference in the two groups of isolates is whether vaccination is used or not. The possible effect that antigenic drift could have on vaccination practices will be discussed.

Recombinant Paramyxovirus type 1-Avian Influenza-H7 virus as a vaccine for protection of chickens against influenza and Newcastle disease

D.E. Swayne¹, D.L. Suarez¹, S. Schultz-Cherry¹, T.M. Tumpey¹, D.J. King¹, T. Nakaya², P. Palese², and A. Garcia-Sastre²

¹USDA, ARS, Southeast Poultry Research Laboratory, Athens, Georgia

²Department of Microbiology, Mount Sinai School of Medicine, New York, New York

Current vaccines to prevent avian influenza rely upon labor-intensive parenteral injection. A more advantageous vaccine would be capable of administration by mass immunization methods such as spray or water vaccination. A recombinant vaccine (rNDV-AIV-H7) was constructed by using a lentogenic Paramyxovirus type 1 vector (Newcastle disease virus [NDV] B1 strain) with insertion of the hemagglutinin (HA) gene from avian influenza virus (AIV) A/chicken/NY/13142-5/94 (H7N2). The recombinant virus had stable insertion and expression of the H7 AIV HA gene as evident by detection of HA expression via immunofluorescence in infected Vero cells. The rNDV-AIV-H7 replicated in 9-10 day embryonating chicken eggs and exhibited hemagglutinating activity from both NDV and AIV proteins that was inhibited by antisera against both NDV and AIV H7. Groups of 2-week-old white Leghorn chickens were vaccinated with transfectant NDV vector (tNDV), rNDV-AIV-H7 or sterile allantoic fluid and were challenged 2 weeks later with viscerotropic velogenic NDV (vvNDV) or highly pathogenic (HP) AIV. The sham-vaccinated birds were not protected from vvNDV or HP AIV challenge. The transfectant

NDV vaccine provided 70% protection for NDV challenge but did not protect against AIV challenge. The rNDV-AIV-H7 vaccine provided partial protection (40%) from vvNDV and HP AIV challenge. The serologic response was examined in chickens that received one or two immunizations of the rNDV-AIV-H7 vaccine. Based on hemagglutination inhibition and ELISA tests, chickens that received a vaccine boost seroconverted to AIV H7, but the serologic response was weak in birds that received only one vaccination. This demonstrates the potential for NDV for use as a vaccine vector in expressing AIV proteins.

Evaluation of a commercial avian influenza (H7N2) vaccine for protection in turkeys against an avian influenza virus (H7N2) isolated from turkeys in Virginia during 2002

Terrence M. Tumpey, Darrell R. Kapczynski, Rashida Horton, and David E. Swayne
Southeast Poultry Research Laboratory, USDA, Agricultural Research Service, Athens, Georgia

During the spring of 2002, a low pathogenic avian influenza (LPAI) A (H7N2) virus caused a major outbreak in commercial poultry in Virginia and adjacent states. The low pathogenic avian influenza (LPAI) virus primarily affected turkey flocks causing respiratory distress and decreased egg production. Experimentally, turkeys were more susceptible than chickens to H7N2 virus infection and higher titers of infectious virus could be recovered from the oropharynx in comparison to the cloaca. The outbreak of H7N2 virus raised concerns regarding the availability of vaccines that could be used for the prevention and control of this virus in poultry. We sought to determine if an existing commercial AI vaccine prepared from a 1997 seed stock virus could provide protection against a 2002 LPAI H7N2 virus isolated from a turkey (A/Turkey/Virginia/158512/02 [TV/02]) in Virginia that was from the same lineage as the vaccine virus. The inactivated AI vaccine, prepared from A/Chicken/Pennsylvania/21342/97 (CP/97) virus, significantly reduced viral shedding from vaccinated turkeys in comparison to sham controls, but did not prevent infection. The

protective effect of vaccination correlated with the level of virus-specific antibody as a second dose of vaccine increased antiviral serum IgG and hemagglutination inhibition (HI) reactivity titers in two different turkey age groups. Serum from CP/97-vaccinated turkeys reacted equally well to CP/97 and TV/02 antigens by HI and ELISA. These results demonstrate the potential benefit of using an antigenically-related 1997 H7N2 virus as a vaccine candidate for protection in poultry against a H7N2 virus isolate from 2002.

Antigen-induced IFN- γ , nitric oxide, and TNF- α production by blood mononuclear cells from *Mycobacterium bovis*-infected cattle: diagnostic implications.

W. R. Waters¹, M. V. Palmer¹, D. L. Whipple¹, M. P. Carlson², and B. J. Nonnecke¹

¹ USDA, ARS, National Animal Disease Center, Ames, Iowa; ² University of Nebraska Veterinary Diagnostic Center, Toxicology Laboratory, Lincoln, Nebraska

Bovine tuberculosis in the United States has proven costly to cattle producers as well as government regulatory agencies. While in vivo responsiveness to mycobacterial antigens is the current standard for the diagnosis of tuberculosis, in vitro assays are gaining acceptance, especially as ancillary or complimentary tests. To evaluate in vitro indices of cellular sensitization, antigen induced interferon (IFN)- γ , nitric oxide (NO), and tumor necrosis factor (TNF)- α responses by blood mononuclear cells from *Mycobacterium bovis*-infected cattle were quantified and compared. Using an aerosol model of infection, two doses each of two strains of *M. bovis* (95-1315 and HC-2045T) were used to induce a range of IFN- γ , NO, and TNF- α responses. Infection specific increases in NO, but not IFN- γ or TNF- α , were detected in nonstimulated cultures at 48h, indicative of nonspecific activation and spontaneous release of NO. The infective dose of *M. bovis* also influenced responses. At 34d postinfection, IFN- γ , NO, and TNF- α responses in antigen-stimulated cells from cattle receiving 10^5 cfu *M. bovis* were greater than responses of cells from cattle infected with 10^3 cfu *M. bovis*. The NO response, but not

IFN- γ and TNF- α responses, was influenced by infective strain of *M. bovis*. TNF- α , NO, and IFN- γ responses followed similar kinetics with strong positive associations between the three readouts.

Overall, these findings indicate that NO and TNF- α , like IFN- γ , may prove useful as indices for the diagnosis of bovine tuberculosis.

Working Group Reports

Immunity and Disease Resistance

Immunointervention

Immunity and the Pathogenesis of Diseases

Nutrition and Immune Function Interactions

Immunity and Disease Resistance

Working Group Leaders:

- Lanie Bilodeau, Immunologists, Catfish Genetics Research Unit, ARS, Stoneville, Mississippi
- Lou Gasbarre, Research Leader, Bovine Functional Genomics Research Unit, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- Michael H. Kogut, Immunologist, Southern Plains Agricultural Research Center, ARS, College Station, Texas
- Joan Lunney, Immunologist, Animal Parasite Diseases Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland

Research Priority 1

Identification of genes defining host disease resistance

1. Background: There is a need for identification and characterization of genes that confer host resistance to important production and foodborne diseases. Studies must determine whether it is possible to identify food animal species that have the potential to be “healthier.”

2. Issue: Identification and characterization of genes defining host disease resistance will enable producers to reduce disease incidence using genomic approaches. Alternately, identifying disease resistant animals may reveal new approaches for disease control.

3. Research Objectives:

- A Identify/develop commercial/experimental animal resource populations for identification and mapping of immunologically relevant genes
- B. Identify and characterize host genes that control important immunologic responses; determine whether alleles of certain genes would enhance animal health
- C. Define important genetic polymorphisms that affect host resistance/susceptibility

- D. Define gene expression patterns and regulatory phenomena that affect the level of host resistance
- E. Utilize genomic information to reveal new approaches for disease control or biotherapeutic production

4. Proposed Accomplishments:

- A. Mapping projects will identify informative populations for disease studies. These can then be linked with known populations and commercial breeding stock. Information gained can be used to plan modification of populations for specific production/health issues.
- B. Information will be developed on correlation of disease resistance phenotype with desired production traits by phenotyping for disease resistance traits using high throughput (HTS) immune and disease screens coordinated with production parameters. This will prove relevance for utilization of disease resistance in commercial populations/genomic stock.
- C. Expansion of gene expression/regulation approaches, includes microarray [targeted and full genomic], real-time RT-PCR, proteomic studies, RNAi technology, leading to a more thorough understanding of disease resistance mechanisms
- D. Expanded knowledge of targeted genomic areas, Immunoglobulin (Ig), T cell

receptor (TCR), major histocompatibility (MHC), pattern recognition receptors (PRR), Toll-like receptor (TLR) pathways, and their impact on health and disease

responses to infection/invasion of the host. Studies focused on host/pathogen interactions at mucosal surfaces are essential to define early events relevant to disease resistance.

5. Proposed Collaborations: Work with scientists in NP101 and NP106 programs to utilize genomics tools developed; similarly; work with national (NRSP008) and international genome efforts, e.g. sequencing and microarray projects; work with national swine PRRS initiative to target specific disease.

6. Anticipated project duration: Significant progress should be made across taxa for gene identification/characterization studies within 5 years.

7. Impact:

- A. Markers/genes regulating disease resistance will be identified. This information will be used for developing allele tests, e.g., single nucleotide polymorphisms (SNPs) for fast, effective identification of resistant animals; for marker assisted selection (MAS) approaches to increase frequency in target populations; or for development of transgenic or clone lines of disease resistant animals
- B. Definition of novel mechanisms of protective immunity. Apply information gained from genomics to developing new biotherapeutics, vaccine approaches (with APHIS).
- C. Develop approaches for diagnostics and new rationales for immune control of diseases

2. Issue: Identification of pathogen uptake sites/mechanisms will enhance understanding of immune responses. Pathogen entry occurs largely across mucosal membranes. In addition to uptake, the study of induction/regulation of immune responses through mucosal surfaces are needed

3. Research Objectives:

- A. Define cell populations/subsets that affect resistance at mucosal surfaces; determine how neonatal development affects these populations.
- B. Define gene expression patterns and regulatory phenomena that affect the level of host resistance at mucosal surfaces.
- C. Use information from microbial genomics to define substances from pathogens that stimulate/regulate immune responses at mucosal surfaces.
- D. Define the affects of stress/diet/age on immune responses at mucosal surfaces.

4. Proposed Accomplishments:

- A. Identification of regional (respiratory, intestinal, reproduction, skin) areas of infection/points of uptake; Determination of mucosal cell populations involved in disease processes; Development of immune tools to assess immune cell subsets regulating responses; Determination of age-related development associations with infection; Development of gene expression/regulation approaches and reagents to quantitate chemokines and cytokines.
- B. Determine relationship of infection kinetics (in blood/specific mucosal tissues) and mucosal immune function; Identification of relevant genes controlling infection processes, e.g., TLR, anti-microbial peptides; MHC genes involved.
- C. Application of microbial genomics information to probe critical microbial genes influencing host disease response

Research Priority 2

Induction/regulation of immune responses through mucosal surfaces

1. Background: Many infectious diseases enter through respiratory and gastrointestinal tissues and infect through mucosal surfaces. There is a need for a better understanding of what drives immune

and means, or mechanisms, of pathogen invasion/evasion/infection.

5. Proposed Collaborations: Interactions with university scientists through NC projects; with international researchers through UK BBSRC, Spain INIA; with commercial producers/animal health companies

6. Anticipated project duration: Significant progress should be made across taxa for gene identification/characterization studies within 5 years.

7. Impact:

- A. Definition of local protective defense mechanisms; Definition of pathways for interventions, for development of both traditional and therapeutic vaccines, and for preparation of novel mucosally targeted biotherapeutics
- B. Improved vaccine efficacy; verification of efficacy of mucosal delivery vectors and adjuvants.
- C. Develop approaches for diagnostics and rationale for immune control of diseases (with APHIS)

Research Priority 3

Induction/regulation of innate responses; influence/regulation of acquired immunity

1. Background: The role of innate immunity, and its influence on acquired immunity, for disease control are still undefined for most agricultural species. Research in this area will influence development of treatments/vaccines/therapeutics, and improve management practices.

2. Issue: The induction/regulation of innate immunity affects disease kinetics, disease resistance, and ultimately animal production. Determination of induction/regulation mechanisms of innate responses, and their influence on the effectiveness of acquired and protective anti-disease immunity, is needed.

3. Research Objectives:

- A. Define cell populations/subsets that affect innate immunity.
- B. Define gene expression patterns and regulatory phenomena that affect the level of host innate immunity.
- C. Use information from microbial genomics to define substances from pathogens that stimulate/regulate innate immune responses; identify the evolutionarily conserved molecular constituents (PAMPs) of infectious microbes.
- D. Define the affects of stress/diet/age on innate immune responses

4. Proposed Accomplishments:

- A. Knowledge of genomic organization of loci involved in the germ-line encoded pattern recognition receptors (PRR), e.g., TLRs, mannose receptors, scavenger receptors, and complement receptors. Expression levels of PRR on effector cells of innate immunity (macrophages, NK cells, granulocytes, epithelial).
- B. Identification of pathogen-associated molecular patterns (PAMPs) and their interaction with PRR will inform pathogen control measures and help identify new biotherapeutics.
- C. Effect of innate responses on intensity and bias of pathogen-specific effector acquired immune responses, and activation of the co-stimulatory molecules required for T cell responses; determine role of innate and acquired immunity in disease kinetics.
- D. Knowledge of development/regulation of immunity in neonates/early developmental stages
- E. Effect of stress and nutritional status on innate immune function.

5. Proposed Collaborations: Interactions with university scientists through NC projects; with international researchers through UK BBSRC, Spain INIA; with commercial producers/animal health companies.

6. Anticipated project duration: Significant progress should be made across taxa for gene

identification/characterization studies within 5 years.

7. Impact: Identification of animals with broader spectrum disease resistance Development of Biotherapeutics; Definition of Local protective defense mechanisms; Identification of innate immune pathways for interventions/traditional and therapeutic vaccines and for development of new adjuvants and improvement of vaccine efficacy

Research Priority 4

Bioinformatics

1. Background: Improvements in genomics technologies have led to high-throughput sequencing which generates large data sets that require bioinformatics tools and computing power necessary for analysis. Currently, the infrastructure does not exist for handling agency-wide, or in some cases unit-level, collation of data and extensive analyses. These resources need to be developed at the unit, area, agency and national levels.

2. Issue: Large volumes of data are generated that are difficult to analyze. For this reason, bioinformatics and genomics resources need to be developed and integrated into ARS programs and research units.

3. Research Objective:

- A. Develop standardized bioinformatics tools and resources that can be utilized by multiple programs and research units.
- B. Develop tools/databases for species sequence, data mining

4. Proposed Accomplishments:

- A. Improve high-powered computer support and information exchange/accessibility. Tools/Statistical analyses of disease mapping info that adjust to population structure and enable efficient data management. Analysis tools and MAS approaches

- B. Improved *in silico* study tools, e.g., comparative analysis tools; improved annotation tools
- D. Proteomics-based tools

5. Proposed Collaborations: Nationally centered bioinformatics resources but species relevant NSRP-8 and TIGR as potential resources; Need for Long term funding for maintenance and updating of websites

6. Anticipated project duration: Significant progress should be made across taxa for gene identification/characterization studies within 5 years.

7. Impact: Utilize genome sequencing information (animals, microbes); Eliminate redundancy; Improved comparative bioinformatics; Improved information exchange. Tool availability for broader animal health community; *in silico* tools for efficient vaccine development

Research Priority 5

Enhance immune toolkit for major food animal species

1. Background: Basic tools to measure details of immune responses in major food animal species are not available or critical components are missing. For example, there are insufficient panels of monoclonal antibodies (mAb) that identify and quantitate CD antigens on immune cell surfaces, chemokines and cytokines; molecular probes and assays for immune markers; cloned and expressed bioactive chemokines and cytokines. Availability of quality targeted immune gene microarrays will be another important resource.

2. Issue: Because of limited markets for these products, production of high quality reagents is often left to researchers; for some species commercial sources of these reagents are limited or non-existent.

3. Research Objectives:

- A. Establish national and international networks to develop priorities for toolkit needs for each species. Spain INIA; with commercial producers/animal health companies. Need for long term funding for maintenance and updating of immune toolkit websites
- B. Coordinate tool development so that priority reagents become available. Work with commercial sources to market products. Simultaneously work with repositories to have independent sources of hybridoma cell lines and expression vectors for cloned and expressed bioactive chemokines and cytokines so that researchers can produce large quantities for research efforts.

4. Proposed Accomplishments:

- A. Panels of reagents, mAb, cloned and expressed bioactive chemokines and cytokines, immune gene microarrays, will be available to researchers for their disease resistance studies
- B. Critical control points in disease processes, and in immune events controlling these processes, will be revealed
- C. Technology transfer opportunities will be expanded

5. Proposed Collaborations: Interactions with university scientists through NC projects; with international researchers through UK BBSRC,

6. Anticipated project duration: Significant progress should be made across taxa for gene identification/characterization studies within 5 years.

7. Impact: Effective tools to explore phenotyping animal immunity, to determine protective immune responses, to confirm bioactive therapeutics. Tools to quantitate immune responses, to validate models of immunity, to assess differential genetic responses of resistant/susceptible to infectious disease challenges

Note: Specific disease targets will be based on stakeholders' priorities.

Research priorities have been listed but numbers do not indicate relative priorities since all approaches are needed for maximal impact on improving disease resistance.

Immunointervention

Working Group Leaders:

- Marcus Kehrli, Research Leader, Virus and Prion Diseases Research Laboratory, National Animal Disease Center, ARS, Ames, Iowa
- Hyun Lillehoi, Immunologist, Animal Parasite Diseases Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- David White, Veterinary Medical Officer, Arthropod-borne Animal Diseases Research Laboratory, ARS, Laramie, Wyoming

Research Priority 1

Identify mechanisms of innate immunity and their regulation critical in control of animal diseases.

1. Background: Many diseases are not amenable to control by traditional vaccination approaches because they are caused by an array of antigenically distinct opportunistic pathogens. Moreover, these opportunistic diseases are often associated with stress, broad-based immune impairment and/or overwhelming pathogen exposure. Some of these diseases may be better controlled by modulation of the innate immune system at critical growth/production stages to improve the host's capacity to resist opportunistic infections and disease.

2. Issue: There is a lack of knowledge about factors regulating humoral and cellular components of innate immunity.

3. Research Objectives:

- A. Identify critical components of innate immunity in cattle, pigs and poultry and how they are regulated.
- B. Develop effective strategies to evaluate the innate immune response in domestic livestock and poultry, including neonatal and embryonic immunity, innate immune cell function (neutrophils, NK cells, etc.), accessory cell function, and assay development for immunological reagents.

- C. Identify inter-relationships between innate immunity and adaptive immunity in cattle, pigs and poultry, and how innate immunity can impact host adaptive immune responses to vaccination or natural infection.

4. Proposed Accomplishments:

- A. Identification of various receptors and cell signaling pathways in livestock that trigger innate immunity. This should also offer insights on how to regulate the activation of innate immunity.
- B. More effective methods to evaluate methods of immunointervention of the innate immune system.
- C. Identification of key regulatory relationships between the innate and adaptive immune systems. This will provide insight on what critical innate co-stimulatory factors are needed to generate an efficacious immune response.

5. Proposed Collaborations: Appropriate ARS CRIS projects, various university collaborations, and pharmaceutical and biologics companies.

6. Anticipated project duration: 5+ years

7. Impact:

- A. Increased understanding of critical innate immunity components in livestock will provide a knowledge base that can be exploited to make science-based decisions to reduce disease incidence and

- severity through targeted immunointervention techniques.
- B. Increased ability to screen modulators of innate immunity against specific pathogens.
- C. Increased understanding of how innate immunity impacts vaccine responses in livestock will provide a knowledge base for better adjuvant and costimulatory molecule inclusion in vaccines, thus improving vaccine efficacy.

Research Priority 2

Identify vaccines that elicit protective cell-mediated immunity

1. Background: Traditional approaches to disease control have been through vaccination to generate protective antibody responses. Many diseases have been effectively controlled by this approach. However, most animal diseases that persist as problems in the U.S. today have not been amenable to control by these traditional approaches. One explanation of how these pathogens remain uncontrolled is the absence of an appropriate protective immune response. Some pathogens may evade the immune system through many mechanisms and may require generation of antigen-specific cytotoxic T-cells to be appropriately controlled. Identification of protective immune responses to the proper antigens requires immunological tools not currently available for widespread use in livestock or wildlife that are important for us to control.

2. Issue: Traditional methods of assessing vaccine efficacy are to measure antibody responses and protection against disease challenge. Unfortunately antibody responses are not always predictive of a protective immune response against live pathogen challenge. We need better tools to assess cell-mediated immune responses in all livestock species.

3. Research Objectives:

- A. Develop new methods in livestock for detecting and measuring cell-mediated

immunity.

- B. Identify antigens critical for the development of cell-mediated immunity.
- C. Identify methods to regulate the development of cell-mediated immunity.
- D. Develop antigen delivery systems (expression vectors for various host animal species to be used for emerging diseases) that target immune system compartments (mucosal, peripheral, reproductive tract and other privileged sites).
- E. Explore effects of modulation or elimination of T-regulatory cell subsets on induction of protective responses in domestic animals and poultry. Develop novel immunomodulators to suppress regulatory cell.

4. Proposed Accomplishments:

- A. Tools for measuring cell-mediated immunity will enable discovery, development and approval of new effective vaccines with cell-mediated immunity label claims.
- B. Methods to predict and select appropriate pathogen antigens for incorporation into vaccines driving cell-mediated immune responses will enhance our response time in the event of emerging diseases.
- C. Understanding the regulatory factors that contribute to an effective cell-mediated immune response will allow us to create vaccines that do a better job of stimulating immunity than the pathogen does by itself.
- D. An improved capacity to rapidly develop vaccines against new and emerging pathogens.

5. Proposed Collaborations:

6. Anticipated project duration: 5 years +

7. Impact:

- A. Vaccines that aid in the control of diseases not currently effectively controlled.

- B. More rapid vaccine discovery and approval.
- C. Better methods to elicit and control immune responses requiring cell-mediated immunity.
- D. An increased capacity to rapidly respond to emerging disease outbreaks.

Research Priority 3

Identify vaccines that elicit protective humoral immunity

1. Background: Some pathogens elicit strong humoral immune responses that are not highly protective. The survival of these pathogens may depend on their ability to divert the efforts of the immune system towards antigens that do not facilitate their clearance from the host.

2. Issue: There is a lack of basic knowledge in what drives successful humoral immune responses against certain pathogens. More indepth knowledge on regulation of immune responses in different compartments (e.g., respiratory tract, intestinal tract, reproductive tract, mammary gland) of the body is needed to better guide vaccine discovery and development into commercial products.

3. Research Objectives:

- A. Identify factors regulating antibody responses to antigens.
- B. Develop computational methods to increase the efficiency of the design and testing of vaccines.
- C. Develop allied testing methods to distinguish between vaccinated and naturally infected animals.
- D. Develop broad spectrum, targeted delivery systems (target dendritic cells or others) for immunomodulator and/or peptide and/or DNA delivery to effector cells.
- E. Elucidate factors affecting efficacy of current adjuvant systems in domestic animals (i.e. optimized CpG sequences for all livestock species) and develop new

immunoadjuvants from novel sources (snake venoms, modified superantigens, probiotics as adjuvants, cytokines – IL-15, hormones, vitamins and trace minerals, carotenoids, phenolics, etc.).

- F. Delineate the specific conditions required for several different broad-based protein or nucleic acid delivery systems – i.e. needle-less systems, microgels, biodegradable depot systems, *in ovo* injector systems – while concentrating on important factors such as multiple use strategies, limiting the injected volume, and livestock or wildlife behavioural responses (fear and/or aggression in response to noise or pain) to ensure field practicality. Evaluate natural delivery agents (i.e. viruses that cross the placenta naturally to deliver nucleic acids to the developing fetus).

4. Proposed Accomplishments:

- A. A better knowledge base of how antibody genes are selected and arranged for various classes of antigens, we can more effectively design next generation vaccines.
- B. By leveraging the knowledge database from human disease research, we will build a comparative immunology database that can be a guide to antigen selection for use in vaccines.
- C. Marked vaccines to distinguish vaccinates from natural infection.
- D. Adjuvant development (i.e. CpG) and/or delivery, peptide or DNA delivery (i.e. microgels, biodegradable depot systems).
- E. Adjuvants and immune modulators that target more effective immune responses.
- F. Broad-based delivery systems applicable to both livestock and wildlife that account for the practical realities of immunization in the field – especially in the case of wildlife, where very little information exists on the efficacy and reliability of strategies used in domestic livestock and poultry.

5. Proposed Collaborations:

6. Anticipated project duration: 5 years +

7. Impact:

- A. Vaccines that aid in the control of diseases not currently effectively controlled.
- B. More rapid vaccine discovery and approval.
- C. Marked vaccines will enable eradication programs once disease prevalence is below a certain economic threshold.
- D. By partnering with industry early in the antigen selection process we will better ensure viability of developed products.
- E. Effective strategies to immunize and modulate the responses of domestic animals and wildlife benefit agriculture by lesser manipulation of animals and lesser exposure to adjuvants that may impact meat or hide quality.
- F. Effective strategies to immunize wildlife will significantly decrease the risk of disease in domestic animals because of a reduction in the size of the reservoir, both in the U.S. and abroad.

Research Priority 4

Develop immunological reagents for domestic animals, poultry, and wildlife that are necessary to aid in the control of diseases we face today.

1. Background: Although many significant advances have been made in the past 3 decades, cutting-edge research into host immune responses in domestic animals and wildlife has been hampered by the cost of developing the necessary reagents and tools of immunological research. These include but are not limited to species-specific cytokines, effector molecules, and receptors, and monoclonal antibodies against each. These tools can be identified and made available with the necessary resources and investment to produce them. However, mechanisms need to be in place to allow production and sharing of these reagents at reasonable costs.

2. Issue: Reagents for the evaluation of immunological parameters in domestic animals and wildlife are severely lacking. Moreover, syngeneic animals or protocols for culture and preservation of immunological cells are necessary for many studies of basic immunological function.

3. Research Objectives:

- A. Identify an appropriate infrastructure with the capacity to express various recombinant proteins and produce monoclonal antibodies against them. This may represent a government “subsidized” company that in turn recovers some of their manufacturing costs through the sale of these reagents to researchers around the world.
- B. Clone, express in native form, characterize, and develop antibodies to MHC Class I and II, acute phase proteins, complement proteins, cytokines, antibody isotypes, innate immune system receptors (TLR’s and others), effector molecules (i.e. perforin, granzyme, etc.), adhesion molecules, and cell surface markers of lymphocyte subsets for relevant domestic livestock and poultry species. House these reagents in some sort of centralized facility (either government or corporate) while balancing the free exchange of reagents for research purposes with the patent and intellectual property rights that will make them viable commercial products. Use the NIH Tetramer core facility as a model of a government-run facility for vital research reagents.
- C. Develop strategies to produce syngeneic domestic livestock strains (inbreeding) or alternatives to fully syngeneic animal development through *in vitro* manipulation immune cells and/or developing MHC-matched animal lines.
- D. Improve production and reduce cost of protein or nucleic acid substrates to make these strategies viable to animal production systems through high efficiency, scalable fermentation or synthesis processes that will produce native (or at least bioactive) protein.

4. Proposed Accomplishments:

- A. A centrally-maintained research facility that provides high-quality reagents to researchers worldwide for the evaluation of immunological parameters in domestic livestock and poultry.
- B. A reasonably priced supply of new immunological reagents for use in livestock, poultry and wildlife research.

5. Proposed Collaborations:

6. Anticipated project duration: 5 years+ (ongoing supplier)

7. Impact:

- A. Availability of new immunological reagents will increase the number of laboratories capable of investigating host immune responses against disease.
- B. Increasing the number of laboratories investigating immunity against disease with modern immunological reagents and tools will increase the probability of identifying new and more efficacious means of immunointervention.
- C. New avenues of research not currently possible will be opened up for scientists to follow.

Research Priority 5

Disease challenge models are needed for many diseases affecting domestic animals and wildlife.

1. Background: Appropriate disease challenge models will be needed to accurately predict the success of any method of immunointervention in livestock or wildlife. It is possible to not recognize successful methods of immunointervention in instances where the model has an overwhelming pathogen challenge not representative of the natural disease exposure. Animal disease researchers have the unique opportunity of working in the natural host for most diseases that are studied, but reliable evaluation of intervention strategies requires standardized

models and conditions that allow for the delineation of intervention effects. Make protocols and strains of pathogens used in models freely available to any interested researcher. Use models to elucidate immune functions relevant to protection from infection or disease, duration of immunity from different intervention protocols (dose, route, formulation, and timing vs. generation of lifelong immunity), and biomarkers of relevant clinical endpoints for testing and future efficacy evaluation. Finally, use models to define immunogenic relationships within and between pathogens to allow for the development and evaluation of cross-protection between vaccine components or interference between immunomodulatory agents.

2. Issue: Models may fail to accurately predict the outcome of immunointervention strategies in the “real world”.

3. Research Objectives:

- A. Develop challenge models for diseases of domestic animals and wildlife.
- B. Develop generalist *in vivo* tests of immune function that are predictive of disease outcome.

4. Proposed Accomplishments:

- A. Models of disease will help both in the development and continuing evaluation of immunointervention strategies.
- B. Biomarkers of immunointervention success will enhance our capacity to screen for the most efficacious method to elicit a protective immune response.

5. Proposed Collaborations:

6. Anticipated project duration: 5 years

7. Impact:

- A. More rapid and uniform testing of immunointervention strategies across laboratories.
- B. Biomarkers that may be used in large scale studies on efficacious immunointervention strategies will reduce the cost of identifying new disease control methods.

Research Priority 6

Vaccine vigilance

1. Background: There are consequences of vaccination that we need to assess to ensure we are not adversely affecting livestock performance and well-being.

2. Issue: There is a need to investigate and characterize immunopathological mechanisms in domestic livestock and poultry.

3. Research Objectives:

- A. Characterize hyper-responsiveness of the immune system, as an allergic reaction (i.e. peanut allergy), an adverse response to immunointervention, or as a part of other physiopathological processes (i.e. pulmonary hypertension syndrome in chickens, autoimmune uveitis in chickens).
- B. Create a database of field response rates and adverse event rates for evaluation of current and future vaccine efficacy.
- C. Characterize field variation of important pathogens and assess its impact on vaccine failure.
- D. Develop strategies for inclusion of multiple strains of the same pathogen, or multiple pathogens, in a single vaccine dose while avoiding the problems of immunodominance/original antigenic sin, antigen overload, and induction of tolerance.
- E. Optimize use of existing vaccines and immunomodulators for intervention at embryogenesis, birth, weaning, or prior to stress (shipping, comingling, slaughter).
- F. Develop *in vitro* assays for efficacy and safety of immunointervention agents and/or insults to the immune system.

4. Proposed Accomplishments:

- A. Fewer adverse event reports.
- B. Centralized database of vaccine usage and adverse events will lead to better monitoring of efficacy and failure. This database could be developed in conjunction with a broader database of "animal passports/ID".
- C. Vaccines that are more producer-friendly through reduced animal manipulation while retaining the efficacy of individual preparations. Vaccines would also be of great use in wildlife management because of decreased need for contact and manipulation of wildlife.
- D. Ensure efficacy of modulation of relevant immunological endpoints in well-controlled experiments.
- E. Assays would be developed that not only evaluate the immunogenicity of relevant immunity in different systems, but that can also be used to predict those factors for screening of novel compounds for continuing development of immunointervention agents. Assays would concentrate on markers of relevant immunity and/or clinical endpoints and not on simple antigen quantitation.

5. Proposed Collaborations: Industrial partners, Food and Drug Administration

6. Anticipated project duration: 5 years+

7. Impact:

- A. Reduction in adverse reactions to vaccines.
- B. More optimal use of vaccines.

Immunity and the Pathogenesis of Diseases

Working Group Leaders:

- Don Knowles, Research Leader, Animal Disease Research Unit, ARS, Pullman, Washington
- David Swayne, Research Leader, Southeast Poultry Research Laboratory, ARS, Athens, Georgia
- Ray Waters, Veterinary Medical Officer, Bacterial Diseases of Livestock Research Unit, National Animal Disease Center, ARS, Ames, Iowa

Research Priority 1

Characterization of immunity in the newborn

1. Background: Immunointervention strategies for food producing animals are primarily targeted at neonatal/juvenile animals. Additionally, optimal development during the neonatal period is critical for attainment of full production of the animal. Immune development (ontogeny) and host/pathogen interactions of newborn livestock and poultry, however, are poorly characterized.

2. Issue: Numerous developmental, maternal, environmental, and infectious factors influence immune responses elicited by neonates. Characterization of these influences on immune development would enhance strategies to improve immunity.

3. Research Objectives:

- A. Determine the impact of maternal immunity on neonatal protection and immune response
- B. Determine the responses of neonates to vaccines
- C. Determine the impact of animal husbandry and environmental factors on neonatal immunity

4. Proposed Accomplishments:

- A. Identified parameters of immune competence in the newborn
- B. Optimized vaccine strategies for level and duration of immunity

C. Develop new tools for evaluation of immune competence

5. Proposed Collaborations: Collaborations within ARS – across facilities, CRIS projects, and disciplines are in place as well as collaborations with industry and academic institutions.

Additional collaborations should be added as needed. Certain aspects of the project will require international collaborations (e.g., Asia, United Kingdom, Australia, New Zealand, South Africa), of which many are already established.

6. Anticipated project duration: Considering the breadth and impact of this project, it will likely be an on-going project.

7. Impact: Improved strategies to enhance neonatal immunity will result and thereby improving production. More specifically, improved methods to enhance vaccine strategies targeting the neonate and maternal immunity as well as methods to limit factors (e.g., environmental, husbandry, nutritional) that negatively impact neonatal immune development will result.

Research Priority 2

Determination of protective immunity to infectious diseases

1. Background/Issue: An understanding of protective immunity is often required for development of successful strategies to limit disease. Unfortunately, specific protective mechanisms for most infectious diseases of food producing animals are not known.

2. Issue: Our understanding of protective immunity and how to assess it is limited. As a result, this limits advances in development of immunity and immune strategies to control infectious diseases.

3. Research Objectives:

- A. Develop and validate challenge models for vaccine efficacy
- B. Determination of mechanisms of protection

4. Proposed Accomplishments:

- A. Develop validated vaccine challenge models
- B. Identify mechanisms of immune protection

5. Proposed Collaborations: Collaborations within ARS – across facilities, CRIS projects, and disciplines are in place as well as collaborations with industry and academic institutions. Certain aspects of the project will require international collaborations (e.g., Asia, United Kingdom, Australia, New Zealand, South Africa), of which many are already established.

6. Anticipated project duration: In general, these are on-going projects with specific program funding.

7. Impact: Establishment of infection model systems and correlates of protection will enhance our ability to develop effective immunointervention strategies.

Research Priority 3

Regulation of the immune response in control of disease

1. Background/Issue: Exacerbation or induction of disease often results from inappropriate or excessive host responses. Understanding mechanisms of how immunity impacts disease development and resolution will improve our ability to control disease.

2. Issue: Some diseases result from the host immune response. Lack of understanding of this phenomenon for specific diseases has limited progress in prevention and treatment strategies. Important to this issue is the development of and/or sharing of reagents (such as monoclonal antibodies) which allow measurement and characterization of specific components of the immune response.

3. Research Objectives:

- A. Determine the mechanisms of immunosuppression
- B. Determine the mechanisms of pathogen persistence
- C. Determine the role of autoimmunity in disease pathogenesis

4. Proposed Accomplishments:

- A. A determination of the role of pathogen persistence in morbidity and mortality
- B. An understanding of the role of immunosuppression in pathogenesis of disease
- C. New strategies to minimize autoimmunity in infectious disease following pathogen exposure

5. Proposed Collaborations: Collaborations within ARS – across facilities, CRIS projects, and disciplines are in place as well as collaborations with industry and academic institutions. Additional collaborations should be added as needed. Certain aspects of the project will require international collaborations (e.g., Asia, United Kingdom, Australia, New Zealand, South Africa), of which many are already established. Develop a collaborative network for reagent development and exchange.

6. Anticipated project duration: Due to inherent complexity, the duration is difficult to define.

7. Impact: Improved methods to limit immune-associated disease and persistence of pathogens thereby increasing production and decrease spread of disease, respectively

Nutrition and Immune Function Interactions

Working Group Leaders:

- Theodore Elsasser, Research Animal Scientist, Growth Biology Laboratory, Animal and Natural Resources Institute, ARS, Beltsville, Maryland
- Simin Meydani, Chief, Nutritional Immunology Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging, ARS, Tufts University, Boston, Massachusetts
- Joe Urban, Research Leader, Nutrient Requirements and Functions Laboratory, Beltsville Human Nutrition Research Center, ARS, Beltsville, Maryland

Research Priority 1

Defining nutrition related biomarkers that measure risk of disease.

1. Background: Risk of disease during periods of physiological stress, such as birth, lactation, weaning, vaccination, ageing, metabolic disease, etc, can be ameliorated by appropriate nutritional support and intervention strategies.

2. Issue: Nutritional biomarkers to assess optimal health, recommendations for nutritionally based interventions and monitoring of biomarkers to predict clinical outcome are needed.

3. Research Objective:

- A. Evaluate markers of immune function and status that respond to nutritional modification.
- B. Identify dietary components that improve immune function and support optimal health.

4. Proposed Accomplishments:

- A. Develop reagents to identify immune markers for food production animals and man that respond to diet.
- B. Validate animal models to test nutrient and immune interactions in man.
- C. Quantitative risk/benefit evaluation of the dietary strategy to the appropriateness of the response.

- D. Identify specific nutrients with beneficial immune modulating properties that optimize immune function.

5. Proposed Collaborations: Develop ARS Nutrition Network to share information on animal and human models of nutrient interaction with immune function. Need for web-based information sharing.

6. Anticipated project duration: Significant progress should be made across species for identification/characterization of biomarkers within 5 years.

7. Impact: Sharing information on the variety of ARS-directed studies on animal models that measure the effect of nutrition on infectious disease and physiological stressors with human models of chronic inflammatory disease will provide a robust evaluation of multi-functional aspects of immune function and eliminate redundancy. This will improve information exchange and identify novel markers that indicate the status of animal and human health related to diet.

Research Priority 2

Identification of nutrient and gene interactions.

1. Background: Novel dietary biomarkers and their associated genes could identify subpopulations of healthy or animals and humans at risk for disease. Genetic identification in

animals of host resistance to important production and foodborne diseases and in humans of risk factors for chronic and genetic diseases is increasing. Collating these data will assist in designing dietary strategies that promote a healthy phenotype.

2. Issue: Identify and characterize genes that define disease risk factors in humans and determine if gene expression is responsive to dietary interventions. Evaluate the effect of nutrition on promoting optimal health in genetically defined disease resistant livestock.

3. Research Objectives:

- A. Identify and characterize host genes that control important immunologic responses and determine responsiveness to diet.
- B. Define gene expression patterns and regulatory phenomena that are sensitive to diet and affect the level of host resistance
- C. Utilize genomic information from human and animal projects to reveal novel nutrient and immune interactions.
- D. Evaluate nutrients that activate immune gene transcription elements as modifiers of immune function.

4. Proposed Accomplishments:

- A. Mapping projects will identify pathogen and stress resistant livestock populations that can be targeted for nutritional studies related to health outcomes.
- B. Technical advances in gene expression/regulation approaches, includes microarray, real-time RT-PCR, proteomic studies, RNAi technology, etc. can be used to evaluate nutrient and gene interactions related to disease, inflammation and stress.
- C. Dietary interventions with genetically identified populations at risk for immunologically based diseases such as asthma and allergy, inflammatory bowel diseases, type-2 diabetes, obesity, etc. can be evaluated for efficacy.

5. Proposed Collaborations: Work with scientists in NP101, NP106 and NP107 programs to utilize genomics tools or identify animal models that can be used for nutritional studies. Partner with NIH and the human genome project to identify at risk populations for nutritional studies.

6. Anticipated project duration: Significant progress should be made on gene identification, characterization and the role of nutrition in functional expression studies within 5 years.

7. Impact:

- A. Integrated strategies for control of pathogens and stress in livestock as well as inflammatory diseases in humans will include dietary recommendations.
- B. The emergence of genetic profiles will include dietary recommendations for human life style and animal management to optimize health.

Research Priority 3

Nutrient-dependent regulation of innate and acquired immunity.

1. Background: Immunological stress affects nutrient partitioning and nutrient utilization affects immune system efficacy. Appropriate nutritional interventions will affect most components of innate and acquired immunity and influence disease susceptibility, vaccination, immunological memory, and the appropriateness and regulation of a balanced immune response.

2. Issue: Maintenance of optimal health and its restoration following stress from infection or inflammation are dependent on adequate nutrition. Improved understanding of the interaction between innate and acquired immunity and the mechanisms of immune regulation will provide an opportunity to examine specific nutrient requirements for these responses.

3. Research Objectives:

- A. Define the affects of diet on stress and age-related changes in innate and acquired

immunity.

- B. Define the effect of diet on the ontogeny of the immune system and interventions that provide a balanced response to stress and disease.
- C. Evaluate the role of immunity in specific nutrient uptake from the intestine.
- D. Define specific nutrient requirements for effective and appropriate vaccination and immunological memory, and for amelioration of periods of immune deficiency such as during pregnancy, parturition, pediatric development, ageing, stress, etc.
- F. Evaluate the role of nutrition in the amelioration of inflammation and immune pathogenesis.

4. Proposed Accomplishments:

- A. Knowledge of nutrient requirements for the development and regulation of immunity in mothers, neonates and the elderly.
- B. Shared discoveries on the role of specific nutrients in cell and tissue function including receptor mobility, cell signaling pathways, regulation of transcription, mucosal barrier function and physiology, trace mineral-dependent enzymatic activity, reactive oxygen and nitrogen species, lipid inflammatory mediators, carbohydrate glycosyltransferases, oral rehydration solutions, neural transmitters, etc.
- C. More effective vaccines and bioprophylactics.
- D. Nutritional recommendations to improve immune efficacy during periods of stress and disease.

5. Proposed Collaborations: Partnerships with food industry and animal health companies; ARS plant scientists to provide modified and functional food sources; nutritionists involved with nutrient bioavailability.

6. Anticipated project duration: Significant progress in immune regulation and nutrient interactions are assured given the high level of interest.

7. Impact:

- A. Provide recommendations for specific nutrients that contribute to reduced stress and enhanced disease resistance.
- B. Identification of nutrients required for improved vaccine efficacy.
- C. Provide identification of diets that enhance immune development and reduce inflammation.

Research Priority 4

Nutrition and applied immunology

1. Background: Food ingestion and the nutrients provided affect resident microflora in the oral, urogenital and gastrointestinal track, and interact with neural and endocrine pathways to reduce stress. Functional nutrient deficiencies or excess affect the balance of the immune system leading to inflammation or inappropriate responses to infection or stress.

2. Issue: Extend the matrix of information to include indirect interactions between nutrients and the immune system as well as the role of nutrients as modulators of immune function.

3. Research Objectives:

- A. Define probiotic microbial populations that affect immune development and functional resistance at mucosal surfaces and on the efficacy of systemic immunity, and the diets (prebiotics) that support appropriate resident or ingested microbial populations.
- B. Establish a repository of microbial strains that benefit host species and apply genetic or selection tools to improve their application.
- C. Study the role of specific nutrients as modulators of immune function, regulators of inflammation, and as adjuvants.

- D. Evaluate the interaction between diet, environment, behavior and stress on the efficacy of immune function.

4. Proposed Accomplishments:

- A. Identification of functional dietary deficiencies or super nutritional intake of nutrients that result from behavior, environment or stress that affect optimal immune function.
- B. Establishment of animal models that evaluate the long term consequences of diet on immune function and chronic disease.
- C. Definition of the affect of inflammation and immune pathogenesis (obesity, chronic disease, infection, stress) on nutrient uptake and partitioning.
- C. Application of genomics information to characterize microbial genes that improve

host health by the production of useful metabolics, immune stimulators, and competition for sites of pathogen invasion.

5. Proposed Collaborations: Interactions with food industry and animal health companies; food survey groups; microbiologists; food technologists; EPA; FDA.

6. Anticipated project duration: Significant progress should be visible within 5 years with support of intra structure and cooperation.

7. Impact:

- A. Definition of patentable food products that affect human and animal immune function.
- B. Better understanding of the interaction between dietary choices and environment on immune function and optimal health.

Bibliography of ARS Immunology Research 2000-2003

This bibliography contains selected citations of immunology research being performed at ARS Research Centers throughout the United States. The citations were drawn from the CRIS database maintained by the USDA, Cooperative State Research, Education, and Extension Service (<http://cris.csrees.usda.gov/>), AgricOLA database at the National Agricultural Library, CAB Abstracts in the United Kingdom, and MedLine at the National Library of Medicine. It is provided to give an overview of the many facets of ARS immunology research.

It is organized alphabetically by State; within each State it is organized alphabetically by the last name of the staff listed.

Auburn, Alabama

Aquatic Animal Health Research

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=64-20-15-00>

Evans JJ

Evans, J.J., C.A. Shoemaker, and P.H. Klesius. 2001. Distribution of *Streptococcus iniae* in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*) following naive inoculation. *Aquaculture* 194: 233-243.

Klesius, P.H., C.A. Shoemaker, and **J.J. Evans**. 2000. Vaccination: A health management practice for preventing diseases caused by *Streptococcus* in tilapia and other cultured fish. In K. Fitzsimmons and Jomar Carvalho Filho, editors, *Panorama da Aquicultura, Brazil. Tilapia Aquaculture in the 21st Century: Proceedings from the Fifth International Symposium on Tilapia Aquaculture*, p. 558-564.

Klesius, P.H., C.A. Shoemaker, and **J.J. Evans**. *Streptococcus iniae* vaccine. Patent allowance given February 28, 2001. Application No. 09/513,143.

Klesius, P.H., **J.J. Evans**, and C.A. Shoemaker. 2000. Communicable bacterial diseases in fish farming: Prevention by vaccination and immunostimulation. In N. Obaldia III, editor, *Proceedings of the XVII Panamerican Congress of Veterinary Sciences*, p. 13-14.

Shelby, R.A., P.H. Klesius, C.A. Shoemaker, and **J.J. Evans**. 2002. Passive immunization of tilapia, *Oreochromis niloticus* (L.), with anti-*Streptococcus iniae* whole sera. *Journal of Fish Diseases* 25(1): 1-6.

Shelby, R.A., C.A. Shoemaker, **J.J. Evans**, and P.H. Klesius. 2001. Development of an indirect ELISA to detect humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*. *Journal of Applied Aquaculture* 11(3): 35-44.

Shoemaker, C.A., P.H. Klesius, and **J.J. Evans**. 2002. In ovo methods for utilizing the modified live *Edwardsiella ictaluri* vaccine against enteric septicemia in channel catfish. *Aquaculture* 203(3/4): 221-227.

Shoemaker, C.A., P.H. Klesius, and **J.J. Evans**. 2001. Prevalence of *Streptococcus iniae* in tilapia, hybrid striped bass and channel catfish from fish farms in the US. *American Journal of Veterinary Research* 62(2): 174-177.

Shoemaker, C.A., P.H. Klesius, and **J.J. Evans**. 2000. Diseases of tilapia with emphasis on economically important pathogens. In K. Fitzsimmons and Jomar Carvalho Filho, editors, *Panorama da Aquicultura, Brazil. Tilapia Aquaculture in the 21st Century: Proceedings from the Fifth International Symposium on Tilapia Aquaculture*, p. 565-572.

Xu, D.H., P.H. Klesius, C.A. Shoemaker, and **J.J. Evans**. 2000. The early development of *Ichthyophthirius multifiliis* in channel catfish tissue in vitro. *Journal of Aquatic Animal Health* 12: 290-296.

Klesius PH

Barros, M.M., C. Lim, and **P.H. Klesius**. 2002. Effect of soybean meal replacement

by cottonseed meal and iron supplementation on growth, immune response and resistance of Channel Catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture* 207(3/4): 263-279.

Evans, J.J., C.A. Shoemaker, and **P.H. Klesius**. Distribution of *Streptococcus iniae* in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*) following naive inoculation. *Aquaculture* 194: 233-243.

Klesius, P.H., C.A. Shoemaker, and J.J. Evans. *Streptococcus iniae* vaccine. Patent allowance given February 28, 2001. Application No. 09/513,143.

Klesius, P.H., C.A. Shoemaker, J.J. Evans and C. Lim. 2001. Vaccines: Prevention of diseases in aquatic animals. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 317-335.

Lim, C., and **P.H. Klesius**. 2003. Influence of dietary levels of magnesium on growth, tissue mineral content, and resistance of channel catfish *Ictalurus punctatus* challenged with *Edwardsiella ictaluri*. *Journal of the World Aquaculture Society* 34(1): 18-28.

Lim, C., C.A. Shoemaker, and **P.H. Klesius**. 2001. The effect of ascorbic acid on immune response in fish. In K. Dabrowski, editor, *Ascorbic Acid in Aquatic Organisms*, CRC Press: Boca Raton, Florida, p. 149-166.

Lim, C., **P.H. Klesius**, and C.D. Webster. 2001. The role of dietary phosphorus, zinc and selenium in fish health. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 201-212.

Lim, C., **P.H. Klesius**, and C.A. Shoemaker. 2001. Dietary iron and fish health. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 189-199.

Shelby, R.A., **P.H. Klesius**, C.A. Shoemaker, and J.J. Evans. 2002. Passive immunization of tilapia, *Oreochromis niloticus* (L.), with anti-*Streptococcus iniae* whole sera. *Journal of Fish Diseases* 25(1): 1-6.

Shelby, R.A., C.A. Shoemaker, and **P.H. Klesius**. 2003. Development of an immunoassay to measure the humoral immune response of hybrid striped bass *Morone chrysops* x *M. saxatilis*. *Veterinary Immunology and Immunopathology* 91(3/4): 217-25.

Shelby, R.A., C.A. Shoemaker, and **P.H. Klesius**. 2002. Detection of humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*, by a monoclonal antibody-based ELISA. *Journal of Applied Aquaculture* 12(3): 23-31.

Shelby, R.A., C.A. Shoemaker, J.J. Evans, and **P.H. Klesius**. 2001. Development of an indirect ELISA to detect humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*. *Journal of Applied Aquaculture* 11(3): 35-44.

Shoemaker, C.A., **P.H. Klesius**, and C. Lim. 2001. Immunity and disease resistance in fish. . In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 149-162.

Shoemaker, C.A., **P.H. Klesius**, and J.J. Evans. 2002. In ovo methods for utilizing the modified live *Edwardsiella ictaluri* vaccine against enteric septicemia in

channel catfish. *Aquaculture* 203(3/4): 221-227.

Shoemaker, C.A., **P.H. Klesius**, and J.J. Evans (2001). Prevalence of *Streptococcus iniae* in tilapia, hybrid striped bass and channel catfish from fish farms in the US. *American Journal of Veterinary Research* 62(2): 174-177.

Xu, D.H. and **P.H. Klesius**. (2003). Protective effect of cutaneous antibody produced by channel catfish, *Ictalurus punctatus* (Rafinesque), immune to *Ichthyophthirius multifiliis* Fouquet on cohabited non-immune catfish. *Journal of Fish Diseases* 26(5): 287-91.

Xu, D.H., **P.H. Klesius**, and C.A. Shoemaker. (2001). Effect of lectins on the invasion of *Ichthyophthirius* to channel catfish tissues. *Diseases of Aquatic Organisms* 45(2): 115-120.

Yildirim, M., C. Lim, P.J. Wan, and **P.H. Klesius**. 2003. Growth performance and immune response of channel catfish (*Ictalurus punctatus*) fed diets containing graded levels of gossypol-acetic acid. *Aquaculture* 219(1-4): 751-768.

Lim CE

Barros, M.M., **C. Lim**, and P.H. Klesius. 2002. Effect of soybean meal replacement by cottonseed meal and iron supplementation on growth, immune response and resistance of Channel Catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture* 207 (3/4): 263-279.

Klesius, P.H., C.A. Shoemaker, J.J. Evans, and **C. Lim**. 2001. Vaccines: Prevention of diseases in aquatic animals. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 317-335.

Lim, C. and P.H. Klesius. 2003. Influence of dietary levels of magnesium on growth, tissue mineral content, and resistance of channel catfish *Ictalurus punctatus* challenged with *Edwardsiella ictaluri*. *Journal of the World Aquaculture Society* 34(1): 18-28.

Lim, C., P.H. Klesius, and C.A. Shoemaker. 2001. Dietary iron and fish health. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 189-199.

Lim, C., P.H. Klesius, and C.D. Webster. 2001. The role of dietary phosphorus, zinc and selenium in fish health. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 201-212.

Lim, C., C.A. Shoemaker and P.H. Klesius. 2001. The effect of ascorbic acid on immune response in fish. K. Dabrowski, editor, *Ascorbic Acid in Aquatic Organisms*, CRC Press: Boca Raton, Florida, p. 149-166.

Lim, C. and C.D. Webster, editors. 2001. *Nutrition and Fish Health*, Haworth Press, Inc.: New York, 365 pp.

Shoemaker, C.A., P.H. Klesius, and **C. Lim**. 2001. Immunity and disease resistance in fish. In C. Lim and C. Webster, editors, *Nutrition and Fish Health*, Haworth Press, Inc.: New York, p. 149-162.

Yildirim, M., **C. Lim**, P.J. Wan, and P.H. Klesius. 2003. Growth performance and immune response of channel catfish (*Ictalurus punctatus*) fed diets containing graded levels of gossypol-acetic acid. *Aquaculture* 219(1-4): 751-768.

Shelby RA

Shelby, R.A., P.H. Klesius, C.A. Shoemaker, and J.J. Evans. 2002. Passive immunization of tilapia, *Oreochromis niloticus* (L.), with anti-*Streptococcus iniae* whole sera. Journal of Fish Diseases 25(1): 1-6.

Shelby, R.A., C.A. Shoemaker, J.J. Evans, and P.H. Klesius. 2001. Development of an indirect ELISA to detect humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*. Journal of Applied Aquaculture 11 (3) p. 35-44.

Shelby, R.A., C.A. Shoemaker, and P.H. Klesius. 2003. Development of an immunoassay to measure the humoral immune response of hybrid striped bass *Morone chrysops* x *M. saxatilis*. Veterinary Immunology and Immunopathology 91(3/4): 217-25.

Shelby, R.A., C.A. Shoemaker, and P.H. Klesius. 2002. Detection of humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*, by a monoclonal antibody-based ELISA. Journal of Applied Aquaculture 12(3): 23-31.

Shoemaker CA

Evans, J.J., **C.A. Shoemaker**, and P.H. Klesius. 2001. Distribution of *Streptococcus iniae* in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*) following naive inoculation. Aquaculture 194: 233-243.

Klesius, P.H., J.J. Evans, and **C.A. Shoemaker**. 2000. Communicable bacterial diseases in fish farming: Prevention by vaccination and immunostimulation. In N. Obaldia III, editor, Panama. Proceedings of the XVII Panamerican Congress of Veterinary Sciences, p. 13-14.

Klesius, P.H., **C.A. Shoemaker**, and J.J. Evans. *Streptococcus iniae* vaccine. Patent allowance given February 28, 2001. Application No. 09/513,143.

Klesius, P.H., **C.A. Shoemaker**, and J.J. Evans 2000. Vaccination: A health management practice for preventing diseases caused by *Streptococcus* in tilapia and other cultured fish. In K. Fitzsimmons and Jomar Carvalho Filho, editors, Panorama da AQUICULTURA, Brazil. Tilapia Aquaculture in the 21st Century Proceedings from the Fifth International Symposium on Tilapia Aquaculture, p. 558-564.

Klesius, P.H., **C.A. Shoemaker**, J.J. Evans, and C. Lim. 2001. Vaccines: Prevention of diseases in aquatic animals. In C. Lim and C. Webster, editors, Nutrition and Fish Health, Haworth Press, Inc.: New York, p. 317-335.

Lim, C., P.H. Klesius, **C.A. Shoemaker**. 2001. Dietary iron and fish health. In C. Lim and C. Webster, editors, Nutrition and Fish Health, Haworth Press, Inc.: New York, p. 189-199.

Lim, C., **C.A. Shoemaker**, and P.H. Klesius. 2001. The effect of ascorbic acid on immune response in fish. In K. Dabrowski, editor, Ascorbic Acid in Aquatic Organisms, CRC Press: Boca Raton, Florida, p. 149-166.

Shelby, R.A., P.H. Klesius, **C.A. Shoemaker**, and J.J. Evans. 2002. Passive immunization of tilapia, *Oreochromis niloticus* (L.), with anti-*Streptococcus iniae* whole sera. Journal of Fish Diseases 25(1): 1-6.

Shelby, R.A., **C.A. Shoemaker**, J.J. Evans, and P.H. Klesius (2001). Development of an indirect ELISA to detect humoral response to *Streptococcus*

iniae infection of Nile tilapia, *Oreochromis niloticus*. Journal of Applied Aquaculture 11(3): 35-44.

Shelby, R.A., **C.A. Shoemaker**, and P.H. Klesius. 2003. Development of an immunoassay to measure the humoral immune response of hybrid striped bass *Morone chrysops* x *Morone saxatilis*. Veterinary Immunology and Immunopathology 91(3/4): 217-225.

Shelby, R.A., **C.A. Shoemaker**, and P.H. Klesius. 2002. Detection of humoral response to *Streptococcus iniae* infection of Nile tilapia, *Oreochromis niloticus*, by a monoclonal antibody-based ELISA. Journal of Applied Aquaculture 12(3): 23-31.

Shoemaker, C.A., P.H. Klesius, and J.J. Evans. 2002. In ovo methods for utilizing the modified live *Edwardsiella ictaluri* vaccine against enteric septicemia in channel catfish. Aquaculture 203(3/4): 221-227.

Shoemaker, C.A., P.H. Klesius, and J.J. Evans. 2001. Prevalence of *Streptococcus iniae* in tilapia, hybrid striped bass and channel catfish from fish farms in the US. American Journal of Veterinary Research 62(2): 174-177.

Shoemaker, C.A., P.H. Klesius, and J.J. Evans. 2000. Diseases of tilapia with emphasis on economically important pathogens. In K. Fitzsimmons and Jomar Carvalho Filho, editors, Panorama da AQUICULTURA, Brazil. Tilapia Aquaculture in the 21st Century Proceedings from the Fifth International

Symposium on Tilapia Aquaculture, p. 565-572.

Shoemaker, C.A., P.H. Klesius and C. Lim. 2001. Immunity and disease resistance in fish. In C. Lim and C. Webster, editors, Nutrition and Fish Health, Haworth Press, Inc.: New York, p. 149-162.

Xu, D.H., P.H. Klesius, **C.A. Shoemaker**, and J.J. Evans. 2000. The early development of *Ichthyophthirius multifiliis* in channel catfish tissue in vitro. Journal of Aquatic Animal Health 12: 290-296.

Xu, D.H., P.H. Klesius, and **C.A. Shoemaker**. 2001. Effect of lectins on the invasion of *Ichthyophthirius* to channel catfish tissues. Diseases of Aquatic Organisms 45(2): 115-120.

Xu D

Xu, D.H. and P.H. Klesius. 2003. Protective effect of cutaneous antibody produced by channel catfish, *Ictalurus punctatus* (Rafinesque), immune to *Ichthyophthirius multifiliis* Fouquet on cohabited non-immune catfish. Journal of Fish Diseases 26(5): 287-91.

Xu, D.H., P.H. Klesius and C.A. Shoemaker. 2001. Effect of lectins on the invasion of *Ichthyophthirius* to channel catfish tissues. Diseases of Aquatic Organisms 45(2): 115-120.

Xu, D.H., P.H. Klesius, C.A. Shoemaker, and J.J. Evans. 2000. The early development of *Ichthyophthirius multifiliis* in channel catfish tissue in vitro. Journal of Aquatic Animal Health 12: 290-296.

Fayetteville, Arkansas

Poultry Production and Products Safety Research

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=62-26-05-00>

Balog JM

Huff, G.R., W.E. Huff, **J.M. Balog**, and N.C. Rath. 2003. The effects of behavior and environmental enrichment on disease resistance of turkeys. *Brain, Behavior, and Immunity* 17(5): 339-49.

Huff, G.R., W.E. Huff, **J.M. Balog**, N.C. Rath, H. Xie, and R.L. Horst. 2002. Effect of dietary supplementation with vitamin D metabolites in an experimental model of turkey osteomyelitis complex. *Poultry Science* 81(7): 958-965.

Huff, G.R., W.E. Huff, N.C. Rath, and **J.M. Balog**. 2000. Turkey osteomyelitis complex. *Poultry Science* 79(7): 1050-1056.

H. Xie, G.R. Huff, W.E. Huff, **J.M. Balog**, P. Holt, and N.C. Rath. 2002. Identification of ovotransferrin as an acute phase protein in chickens. *Poultry Science* 81(1): 112-20.

Donoghue AM

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and **A.M. Donoghue**. 2002. Prevention of *Escherichia coli* respiratory infection in broiler chickens with bacteriophage (SPR02). *Poultry Science* 81(4): 437-441.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and **A.M. Donoghue**. 2001. Efficacy of bacteriophage to prevent *Escherichia coli* respiratory infection in broiler chickens when administered in the drinking water prior to challenge. *Poultry Science* 80(Suppl.1): 238.

Huff W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and **A.M. Donoghue**. 2001. Bacteriophage prevents *Escherichia coli* respiratory infection in broiler chickens. *Poultry Science* 80(Suppl. 1): 237.

Huff GR

Huff, G.R., W.E. Huff, J.M. Balog, and N.C. Rath. 2003. The effects of behavior and environmental enrichment on disease resistance of turkeys. *Brain, Behavior, and Immunity* 17(5): 339-49.

Huff, G.R., W.E. Huff, J.M. Balog, and N.C. Rath. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Escherichia coli* challenge. 1. Production values and physiological response. *Poultry Science* 80: 1305-1313.

Huff, G.R., W.E. Huff, J.M. Balog, and N.C. Rath. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Escherichia coli* challenge. 2. Resistance to air sacculitis and turkey osteomyelitis complex. *Poultry Science* 80: 1314-1322.

Huff, G.R., W.E. Huff, N.C. Rath, and J.M. Balog. 2001. Effect of feed supplementation with vitamin D metabolites in a dexamethasone-*Escherichia coli* challenge model of turkey osteomyelitis complex. *Poultry Science* 80(Suppl. 1): 238.

Huff, W.E., **G.R. Huff**, N.C. Rath, J.M. Balog, and A.M. Donoghue. 2002. Prevention of *Escherichia coli* infection

in broiler chickens with a bacteriophage aerosol spray. Poultry Science 81(10): 1486-1491.

Huff, W.E., **G.R. Huff**, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2002. Prevention of *Escherichia coli* respiratory infection in broiler chickens with bacteriophage (SPR02). Poultry Science 81(4): 437-441.

Huff, W.E., **G.R. Huff**, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Bacteriophage prevents *Escherichia coli* respiratory infection in broiler chickens. Poultry Science 80(Suppl. 1): 237.

Huff, W.E., **G.R. Huff**, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Efficacy of bacteriophage to prevent *Escherichia coli* respiratory infection in broiler chickens when administered in the drinking water prior to challenge. Poultry Science 80(Suppl.1): 238.

Xie, H., **G.R. Huff**, W.E. Huff, J.M. Balog, and N.C. Rath. 2002. Effects of ovotransferrin on chicken macrophages and heterophil-granulocytes. Developmental and Comparative Immunology 26(9): 805-815.

Xie, H., N.C. Rath, **G.R. Huff**, J.M. Balog, and W.E. Huff. 2001. Inflammation-induced changes in serum modulate chicken macrophage function. Veterinary Immunology and Immunopathology 80 (3/4): 225-235.

Huff WE

Huff, G.R., **Huff, W.E.**, N.C. Rath, and J.M. Balog. 2001. Effect of feed supplementation with vitamin D metabolites in a dexamethasone-*Escherichia coli* challenge model of

turkey osteomyelitis complex. Poultry Science 80(Suppl. 1): 238.

Huff, G.R., **Huff, W.E.**, J.M. Balog, and N.C. Rath. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Escherichia coli* challenge. 1. Production values and physiological response. Poultry Science 80: 1305-1313.

Huff, G.R., **Huff, W.E.**, J.M. Balog, and N.C. Rath. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Escherichia coli* challenge. 2. Resistance to air sacculitis and turkey osteomyelitis complex. Poultry Science 80: 1314-1322.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, and A.M. Donoghue. 2003. Evaluation of aerosol spray and intramuscular injection of bacteriophage to treat an *Escherichia coli* respiratory infection. Poultry Science 82(7): 1108-12.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, and A.M. Donoghue. 2002. Prevention of *Escherichia coli* infection in broiler chickens with a bacteriophage aerosol spray. Poultry Science 81(10): 1486-1491.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2002. Prevention of *Escherichia coli* respiratory infection in broiler chickens with bacteriophage (SPR02). Poultry Science 81(4): 437-441.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Bacteriophage prevents *Escherichia coli* respiratory infection in broiler chickens. Poultry Science 80(Suppl. 1): 237.

Huff, W.E., G.R. Huff, N.C. Rath, J.M. Balog, H. Xie, P.A. Moore, Jr., and

A.M. Donoghue. 2001. Efficacy of bacteriophage to prevent *Eschericheria coli* respiratory infection in broiler chickens when administered in the drinking water prior to challenge. Poultry Science 80(Suppl.1): 238.

Xie, H, G.R. Huff, **W.E. Huff**, J.M. Balog, N.C. Rath. 2002. Effects of ovotransferrin on chicken macrophages and heterophil-granulocytes. Developmental and Comparative Immunology. 26(9): 805-815.

Xie, H., N.C. Rath, G.R. Huff, J.M. Balog, and **W.E. Huff**. 2001. Inflammation-induced changes in serum modulate chicken macrophage function. Veterinary Immunology and Immunopathology 80 (3/4): 225-235.

Rath NC

Huff, G.R., W.E. Huff, **N.C. Rath**, J.M. Balog. 2001. Effect of feed supplementation with vitamin D metabolites in a dexamethasone-*Eschericheria coli* challenge model of turkey osteomyelitis complex. Poultry Science 80(Suppl. 1): 238.

Huff, G.R., W.E. Huff, J.M. Balog, **N.C. Rath**. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Eschericheria coli* challenge. 1. Production values and physiological response. Poultry Science 80: 1305-1313.

Huff, G.R., W.E. Huff, J.M. Balog, and **N.C. Rath**. 2001. Effect of early handling of turkey poults on later responses to a dexamethasone *Eschericheria coli* challenge. 2. Resistance to air sacculitis and turkey osteomyelitis complex. Poultry Science 80: 1314-1322.

Huff, G.R., Zheng, Q., Newberry, L.A., W.E. Huff, J.M. Balog, **N.C. Rath**, K.S. Kim, E.M. Martin, S.C. Goeke, J.K.

Skeeles. 2001. Viral and bacterial agents associated with experimental transmission of infectious proventriculitis of broiler chickens. Avian Diseases 45(4): 828-843.

Huff, W.E., G.R. Huff, **N.C. Rath**, J.M. Balog, and A.M. Donoghue. 2002. Prevention of *Eschericheria coli* infection in broiler chickens with a bacteriophage aerosol spray. Poultry Science 81(10): 1486-1491.

Huff, W.E., G.R. Huff, **N.C. Rath**, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Prevention of *Eschericheria coli* respiratory infection in broiler chickens with bacteriophage (SPR02). Poultry Science 81(4): 437-441.

Huff, W.E., G.R. Huff, **N.C. Rath**, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Bacteriophage prevents *Eschericheria coli* respiratory infection in broiler chickens. Poultry Science 80(Suppl. 1): 237.

Huff, W.E., G.R. Huff, **N.C. Rath**, J.M. Balog, H. Xie, P.A. Moore, Jr., and A.M. Donoghue. 2001. Efficacy of bacteriophage to prevent *Eschericheria coli* respiratory infection in broiler chickens when administered in the drinking water prior to challenge. Poultry Science 80(Suppl.1): 238.

Xie, H, G.R. Huff, W.E. Huff, J.M. Balog, and **N.C. Rath**. 2002. Effects of ovotransferrin on chicken macrophages and heterophil-granulocytes. Developmental and Comparative Immunology 26(9): 805-815.

Xie, H., **N.C. Rath**, G.R. Huff, J.M. Balog, and W.E. Huff. 2001. Inflammation-induced changes in serum modulate chicken macrophage function. Veterinary Immunology and Immunopathology 80(3/4): 225-235.

Little Rock, Arkansas

Arkansas Children's Nutrition Center

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=62-51-10-00>

Badger TM

Badger, T.M., R. Hakkak, and M.J.J. Ronis. 2001. Developmental effects and health aspects of soy protein isolate, casein and whey in male and female rats. *International Journal of Toxicology* 20: 165-175.

Hakkak, R., S. Korourian, T. Fletcher, K. Hale, M. Ferguson, D. Holder, M.J.J. Ronis, J.C. Rowlands, J. Johnston, and **T.M. Badger**. 2001. Effects of whey protein consumption in different stages of life on protection of DMBA-induced mammary tumors in rats. *Proceedings of the American Association of Cancer Research* 42: 311.

Hakkak, R., S. Korourian, D. Irby, J. Sweeny, C. Weatherford, M.J.J. Ronis, J.C. Rowlands, and **T.M. Badger**. 2001. Effects of whey protein consumption on initiation and promotion of DMBA-induced mammary tumors in rats. *Proceedings of American Association for Cancer Research* 42: 892.

Hakkak, R., S. Korourian, M.J.J. Ronis, J. Johnston, and **T.M. Badger**. 2001. Lifetime soy protein isolate consumption protects against azoxymethane- induced colon tumors in male rats. *Cancer Letters* 166: 27-32.

Ronis, M.J.J., C. Mercado, D. Holder, S. Sharp, D. Irby, R. Hakkak, and **T.M.**

Badger. 2001. Inducer-specific effects of dietary protein source on CYP2B1 expression and activity in male rat liver. *The FASEB Journal* 15: A627.

Hakkak R

Badger, T.M., **R. Hakkak**, and M.J.J. Ronis. 2001. Developmental effects and health aspects of soy protein isolate, casein and whey in male and female rats. *International Journal of Toxicology* 20: 165-175.

Hakkak, R., S. Korourian, M.J.J. Ronis, J. Johnston, and T.M. Badger. 2001. Lifetime soy protein isolate consumption protects against azoxymethane- induced colon tumors in male rats. *Cancer Letters* 166: 27-32.

Prior RL

Han, S.N., M. Meydani, D. Wu, B.S. Bender, D.E. Smith, J. Vina, G. Cao, **R.L. Prior**, S.N. Meydani. 2000. Effect of long-term dietary antioxidant supplementation on influenza virus infection. *Journals of Gerontology Biological Sciences and Medical Sciences* 55A: B1-B

Davis, California

Western Human Nutrition Research Center

<http://www.whnrc.usda.gov/>

Hwang DH

Boudreau, M.D., K.H.Sohn, S.H. Rhee, S.W. Lee, J.D. Hunt, and **D.H. Hwang**. 2001. Suppression of tumor cell growth both in nude mice and in culture by n-3 polyunsaturated fatty acids—mediation through cyclooxygenase-independent pathways. *Cancer Research* 61: 1386-1391.

Bray, G.A., J.C. Lovejoy, S.R. Smith, J.P. Delany, M. Lefevre, **D.H. Hwang**, D.H. Ryan, and D.A. York. 2002. The influence of different fats and fatty acids on obesity, insulin resistance and inflammation. *Journal of Nutrition* 132: 2488-2491.

Gao, Z., **D.H. Hwang**, F. Bataille, L. Michael, D. York, M.J. Quon, and J. Ye. 2002. Serine phosphorylation of insulin receptor substrate 1(IRS-1) by inhibitor Kappa B kinase(IKK) complex. *Journal of Biological Chemistry* 277: 48115-48121.

Han, J.A., J.I. Kim, **D.H. Hwang**, L.R. Ballou, A. Mahale, S.A. Aaronson, and S.W. Lee. 2002. p53 induction of COX-2 attenuates p53-or genotoxic stress-mediated apoptosis *EMBO Journal* 21: 5635-5644.

Hwang, D.H. 2001. Signaling pathways for the expression of cyclooxygenase—potential targets of modulation by nutrient. *FASEB Journal* 15: 2556-2564.

Hwang, D.H., V. Fung, and A.J. Dannenberg. 2002. National Cancer Institute Workshop on Chemopreventive Properties of Non-steroidal Anti-inflammatory Drugs: Role of COX-Dependent and -Independent Mechanisms. *Neoplasia* 4: 1-7.

Lee, J.Y. and **D.H. Hwang**. 2002. Docosahexaenoic acid suppresses the activity of peroxisome proliferator-activated receptors in a colon tumor cell line. *Biochemical and Biophysical Research Communications* 298: 667-674.

Lee, J.Y., A. Plakidas, W.H. Lee, P. Chanmugam, A. Heikinnen, G.A. Bray, and **D.H. Hwang**. 2003. Differential modulation of Toll-like receptors by fatty acids—Preferential inhibition by n-3 polyunsaturated fatty acids. *Journal of Lipid Research* 44: 479-486.

Lee, J.Y., K.H. Sohn, S.H. Rhee, and **D.H. Hwang**. 2001. Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *Journal of Biological Chemistry* 276: 16683-16689.

Lee, J.Y., J. Ye, Z. Gao, H.S. Youn, W.H. Lee, L. Zhao, and **D.H. Hwang** (2003). Reciprocal modulation of TLR4 signaling pathways involving MyD88 and PI3K/AKT by saturated and polyunsaturated fatty acids (*J. Biol. Chem.*, in Press).

Paik, J.H., J.Y. Lee, and **D. Hwang**. 2002. Signaling pathways for TNF-induced COX-2 expression: mediation through MAP kinases and NF- κ B, and inhibition by certain nonsteroidal anti-inflammatory drugs. *Advances in Experimental Medicine and Biology* 57: 503-508.

Jacob RA

Stephensen, C.B., **R. Jacob**, G. Aiello, G. Marquis, L. Kruzich, and C.M. Wilson. 2001. For the REACH Project. *Vitamin C*

intake and plasma ascorbic acid in HIV+ and HIV- U.S. adolescents and young adults in the REACH study. *FASEB Journal* 15: A739.

Kelley DS

Benito, P., G.J. Nelson, **D.S. Kelley**, G. Bartolini, P.C. Schmidt, and V.A. Simon. 2001. The effect of conjugated linoleic acid on platelet function, platelet fatty acid composition, and blood coagulation in humans. *Lipids* 36: 221-227.

Benito, P., G.J. Nelson, **D.S. Kelley**, P.C. Schmidt, and V.A. Simon. 2001. The effect of conjugated linoleic acid on plasma lipoproteins and tissue fatty acid composition in humans. *Lipids* 36: 229-236.

Hawkes, W.C., **D.S. Kelley**, and P.C. Taylor. 2001. The effects of dietary selenium on the immune system in healthy men. *Biological Trace Element Research* 81(3): 189-213.

Kelley, D.S. 2001. Modulation of human immune and inflammatory responses by dietary fatty acids. *Nutrition* 17: 669-673.

Kelley, D.S., V.A. Simon, P.C. Taylor, I.L. Rudolph, P. Benito, G.J. Nelson, B.E. Mackey, and K.L. Erickson. 2001. Dietary supplementation with conjugated linoleic acid increased its concentration in human peripheral blood mononuclear cells, but did not alter their function. *Lipids* July 2001. v. 36 (7) p. 669-674.

Kelley, D.S., P.C. Taylor, I.L. Rudolph, P. Benito, G.J. Nelson, B.E. Mackey, K. Erickson. 2000. Dietary conjugated linoleic acid did not alter immune status in young healthy women. *Lipids* 35: 1065-1071.

Kelley, D.S., J.M. Warren, V.A. Simon, G. Bartolini, B.E. Mackey, K.L. Erickson.

2002. Similar effects of c9,t11-CLA and t10,c12-CLA on immune cell functions in mice. *Lipids* 37(7): 725-728.

Pinna, K., **D.S. Kelley**, P.C. Taylor, and J.C. King. 2002. Immune functions are maintained in healthy men with low zinc intake. *Journal of Nutrition* 132(7): 2033-2036.

Rudolph, I.L., **D.S. Kelley**, K.C. Klasing, and K.L. Erickson. 2001. Regulation of cellular differentiation and apoptosis by fatty acids and their metabolites. *Nutrition Research* 21: 381-393.

Stephensen CB

Drammeh, B., G.S. Marquis, **C.B. Stephensen**, and I. Eto. 2001. A randomized fruit-based intervention study to improve vitamin A status or rural Gambian children. *FASEB Journal* 15: A732.

Kemper, C., M. Leung, **C.B. Stephensen**, C.A. Pinkert, M.K. Liszewski, R. Cattaneo, and J.P. Atkinson. 2001. Membrane cofactor protein (MCP; CD46) expression in transgenic mice. *Clinical and Experimental Immunology* 124: 180-89.

Kruzich, L.A., G.S. Marquis, **C.B. Stephensen**, and C.M. Wilson. 2001. Reported micronutrient intakes among HIV-positive and HIV-negative adolescents and young adults in the REACH study. *FASEB Journal* 15: A624. Mitra, A.K., M. Wahed, A. Chowdhury, and **C.B. Stephensen**. 2001. Factors associated with urinary loss of vitamin A in children with watery diarrhea. *FASEB Journal* 15: A256.

Stephensen, C.B. 2003. Vitamin A, beta-carotene, and mother-to-child transmission of HIV. *Nutrition Reviews* 61(8): 280-284.

Stephensen, C.B. 2001. Vitamin A, infection and immune function. *Annual Review of Nutrition* 21: 167-92.

Stephensen, C.B. 2001. Examining the effect of a nutrition intervention on immune function in healthy humans: what do we mean by immune function and who is really healthy anyway. *American Journal of Clinical Nutrition* 74(5): 565-566.

Stephensen, C.B., R. Jacob, G. Aiello, G. Marquis, L. Kruzich, and C.M. Wilson. 2001. Vitamin C intake and plasma ascorbic acid in HIV+ and HIV- U.S. adolescents and young adults in the REACH study. *FASEB Journal* 15: A739.

Stephensen, C.B., R. Rasooly, X. Jiang, M.A. Ceddia, C.T. Weaver, R.A. Chandraratna, and R.P. Bucy. 2002. Vitamin A enhances in vitro Th2 development via retinoid X receptor pathway. *Journal of Immunology* 168(9): 4495-503.

Stephensen, C.B., R. Rasooly, X. Jiang, M.A. Ceddia, C.T. Weaver, R.A.S. Chandraratna, and R.P. Bucy. 2001. Retinoic Acid modulates in vitro Th1/Th2 phenotype development of naive T-cells from DO11.10 T-cell receptor transgenic mice. *FASEB Journal* 15: A1093.

Zunino SJ

Brünke, J.B., K. Fischer, K. Barbin, Y. Schreiter, K. Wachter, F. Mahr, M. Titgemeyer, M. Peipp, **S.J. Zunino**, R. Repp, T. Valerius, and G.H. Fey. 2003. A recombinant bispecific single-chain Fv antibody against HLA class II and FcγRIII (CD16) triggers effective antibody-mediated lysis of lymphoma cells. Manuscript submitted, *British Journal of Hematology*

Dörrie, J., H. Gerauer, Y. Wachter, and

S.J. Zunino. 2001. Resveratrol induces extensive apoptosis by depolarizing mitochondrial membranes and activating caspase-9 in acute lymphoblastic leukemia cells. *Cancer Research* 61: 4731-4739.

Dörrie, J., K. Sapala, and **S.J. Zunino.** 2002. Interferon-γ increases the expression of glycosylated CD95 in B-leukemic cells: An inducible model to study the role of glycosylation in CD95-signaling and trafficking. *Cytokine* 18: 98-107.

Dörrie, J., K. Sapala, and **S.J. Zunino.** 2001. Carnosol-induced apoptosis and downregulation of Bcl-2 in B-lineage leukemia cells. *Cancer Letter* 170: 33-39.

Kellner, C. and **S.J. Zunino.** 2004. Nitric oxide is synthesized in acute leukemia cells after exposure to phenolic antioxidants and initially protects against mitochondrial membrane depolarization. *Cancer Letter* 215(1): 43-52.

Knott, A., Y. Mueller, **S.J. Zunino**, C. Berens, and W. Hillen. 2003. Conditional cell suicide using dox-dependent caspase-2 expression. *Journal of Gene Medicine* 5: 343-354.

Peipp, M., H. Küpers, D. Saul, B. Schlierf, J. Greil, **S.J. Zunino**, M. Gramatzki, and G.H. Fey. 2002. A recombinant CD7-specific single-chain immunotoxin is a potent inducer of apoptosis in acute leukemic T cells. *Cancer Research* 62: 2848-2855.

Peipp, M., D. Saul, K. Barbin, J. Bruenke, **S.J. Zunino**, J.G. van de Winkel, M. Niederweis, and G.H. Fey. 2003. Efficient secretion of fluorescent scFv-fusion proteins directed against CD antigens from insect and mammalian cells. Manuscript submitted, *Journal of Immunological Methods*.

Peipp, M., N. Simon, A. Loichinger, W. Baum, K. Mahr, **S.J. Zunino**, and G.H. Fey. 2001. An improved procedure for the generation of recombinant single-chain Fv

antibody fragments reacting with human CD13 on intact cells. *Journal of Immunological Methods* 251: 161-176.

Athens, Georgia

Southeast Poultry Research Laboratory

<http://sepri.ars.usda.gov>

Gast RK

Gast, R.K. and P.S. Holt. 2001. The relationship between the magnitude of the specific antibody response to experimental *Salmonella enteritidis* infection in laying hens and their production of contaminated eggs. *Avian Diseases* 45(2): 425-431.

Gast, R.K., M.S. Nasir, M.E. Jolley, P.S. Holt, and H.D. Stone. 2002. Detection of experimental *Salmonella enteritidis* and *S. typhimurium* infections in laying hens by fluorescence polarization assay for egg yolk antibodies. *Poultry Science* 81(8): 1128-1131.

Holt, P.S. and **R.K. Gast**. 2002. Comparison of the effects of infection with *Salmonella enteritidis*, in combination with an induced molt, on serum levels of the acute phase protein, alpha1 acid glycoprotein, in hens. *Poultry Science* 81(9): 1295-1300.

Holt, P.S., L.E. Vaughn, **R.K. Gast**, and H.D. Stone. 2002. Development of a lavage procedure to collect crop secretions from live chickens for studying crop immunity. *Avian Pathology* 31(6): 589-592.

Seo, K.H., P.S. Holt, R.E. Brackett, **R.K. Gast**, and H.D. Stone. 2002. Mucosal humoral immunity to experimental *Salmonella enteritidis* infection in the

chicken crop. *Avian Diseases* 46(4): 1015-1020.

Seo, K.H., P.S. Holt, L.E. Vaughn, **R.K. Gast**, and H.D. Stone. 2003. Detection of *Salmonella enteritidis*-specific immunoglobulin A antibodies in crop samples from chickens infected with *Salmonella enteritidis*. *Poultry Science* 82(1): 67-70.

Holt PS

Chaubal, L.H. and **P.S. Holt**. 2000. Characterization of motility and identification of flagella proteins in the avian pathogen *Salmonella pullorum*. *American Journal of Veterinary Research* 60: 1322-1327.

Gast, R.K. and **P.S. Holt**. 2001. The relationship between the magnitude of the specific antibody response to experimental *Salmonella enteritidis* infection in laying hens and their production of contaminated eggs. *Avian Diseases* 45(2): 425-431.

Gast, R.K., M.S. Nasir, M.E. Jolley, **P.S. Holt**, and H.D. Stone. 2002. Detection of experimental *Salmonella enteritidis* and *S. typhimurium* infections in laying hens by fluorescence polarization assay for egg yolk antibodies. *Poultry Science* 81(8): 1128-1131.

Holt, P.S. and R.K. Gast. 2002. Comparison of the effects of infection with *Salmonella enteritidis*, in combination with an induced

molt, on serum levels of the acute phase protein, alpha 1 acid glycoprotein, in hens. Poultry Science 81(9): 1295-1300.

Holt, P.S., L.E. Vaughn, R.K. Gast, and H.D. Stone. 2002. Development of a lavage procedure to collect crop secretions from live chickens for studying crop immunity. Avian Pathology 31(6): 589-592.

Seo, K.H., **P.S. Holt,** R.E. Brackett, R.K. Gast, and H.D. Stone. 2002. Mucosal humoral immunity to experimental *Salmonella enteritidis* infection in the chicken crop. Avian-Diseases 46(4): 1015-1020.

Seo, K.H., **P.S. Holt,** L.E. Vaughn, R.K. Gast, and H.D. Stone. 2003. Detection of *Salmonella enteritidis*-specific immunoglobulin A antibodies in crop samples from chickens infected with *Salmonella enteritidis*. Poultry Science 82(1): 67-70.

Kapczynski DR

Kapczynski, D.R., D.A. Hilt, D. Shapiro, H.S. Sellers, and M.W. Jackwood. 2003. Protection of chickens from infectious bronchitis by in ovo and intramuscular vaccination with a DNA vaccine expressing the S1 glycoprotein. Avian Diseases 47(2): 272-85.

Kapczynski, D.R., H.S. Sellers, G.N. Rowland, and M.W. Jackwood. 2002. Detection of in ovo-inoculated infectious bronchitis virus by immunohistochemistry and in situ hybridization with a riboprobe in epithelial cells of the lung and cloacal bursa. Avian Diseases 46(3): 679-85.

Kapczynski, D.R., T.M. Tumpey. 2003. Development of a virosome vaccine for Newcastle disease virus. Avian Diseases 47(3): 578-87.

Sellers, H.S., P.N. Villegas, J. El-Attrache, **D.R. Kapczynski,** and C.C. Brown. 2001. Detection of infectious bursal disease virus in experimentally infected chickens by in situ hybridization. Avian Diseases 45(1): 26-33.

King DJ

King, D.J. 2000. Efficacy of vaccine in protection against velogenic Newcastle Disease (ND). Proceedings of the United Animal Health Association, p.591-593.

King, D.J., G.D. Kommers, C.C. Brown, and B.S. Seal. 2001. Virulence of pigeon Newcastle disease virus isolates for chickens. Proceedings of the Fiftieth Western Poultry Disease Conference, p.15-16.

King, D.J. 2001. Selection of thermostable Newcastle Disease Virus from reference and vaccine strains. Avian Diseases 45: 512-516.

King, D.J. and B.S. Seal. 2000. Current understanding of molecular correlates of virulence in Newcastle Disease Virus. International Hatchery Practice 14(7)(Suppl.): 10-11.

King, D.J. and D.E. Swayne. 2000. Newcastle disease update: International ND problems. Proceedings of the United Animal Health Association, p.593-596.

Kommers, G.D., **D.J. King,** B.S. Seal, and C.C. Brown. 2003. Virulence of six heterogeneous-origin Newcastle disease virus isolates before and after sequential passages in domestic chickens. Avian-Pathology 32(1): 81-93.

Kommers, G.D., **D.J. King,** B.S. Seal, and C.C. Brown. 2000. Pathogenesis of five different pigeon-origin isolates of Newcastle disease virus for domestic chickens. Veterinary Pathology 37(5): 546 (Abstract No.94).

Locke, D.P., H.S. Sellers, J.M. Crawford, S. Schultz-Cherry, **D.J. King**, R.J. Meinersmann, and B.S. Seal. 2000. Newcastle disease virus phosphoprotein gene analysis and transcriptional editing in avian cells. *Virus Research* 69: 55-68.

Seal, B.S. and **D.J. King**. 2000. Newcastle Disease Virus. Available from: <http://www.els.net> Encyclopedia of Life Sciences.

Sellers, H.S., S. Schultz-Cherry, C.C. Brown, B.S. Seal, and **D.J. King**. 2000. Induction of apoptosis by Newcastle disease virus. Program of the Fifth International Congress of Veterinary Virology, Brescia, Italy. August, 2000.(Abstract).

Suarez, D.L., D.E. Swayne, and **D.J. King**. 2001. The ongoing threat of avian influenza and Newcastle disease to the U.S. poultry industry. Proceedings of the Fiftieth Western Poultry Disease Conference, p.1-7.

Swayne, D.E., D.L. Suarez, and **D.J. King**. 2000. Avian influenza (AI) and velogenic Newcastle disease (VND) update. Proceedings of 35th National Meeting of Poultry Health and Processing, p.37-45.

Perdue ML

Basler, C.F., A.H. Reid, J.K. Dybing, T.A. Janczewski, T.G. Fanning, H. Zheng, M. Salvatore, **M.L. Perdue**, D.E. Swayne, A. Garcia-Sastre, P. Palese, and J.K. Taubenberger. 2001. Sequence of the 1918 pandemic influenza virus non-structural gene (ns) segment and characterization of recombinant viruses bearing the 1918 NS genes. *Proceedings of the National Academy of Science* 98: 2746-2751.

Perdue, M.L., D.L. Suarez, and D.E. Swayne. 2000. Avian influenza in the

90's. *Poultry and Avian Biology Reviews* 11: 1-20.

Schultz-Cherry, S., J.K. Dybing, N.L. Davis, C. Williamson, D.L. Suarez, R. Johnston, and **M.L. Perdue**. 2000. Influenza virus (A/HK/156/97) hemagglutinin expressed by an alphavirus replicon system protects chickens against lethal infection with Hong Kong-origin H5N1 viruses. 255: 55-59.

Swayne, D.E., J.R. Beck, **M.L. Perdue**, and C.W. Beard. 2001. Efficacy of vaccines in chickens against highly pathogenic Hong Kong H5N1 avian influenza. *Avian Diseases* 45: 355-365.

Seal BS

Cameron, K.T., X. Zhang, **B.S. Seal**, M. Rodriguez, and M.K. Njenga. 2001. Antigens to viral capsid and non-capsid proteins are present in brain tissues and antibodies in sera of Theiler's virus-infected mice. *Journal of Virological Methods* 91: 11-19.

Gulati, B.R., K.T. Cameron, **B.S. Seal**, S.M. Goyal, D.A. Halvorson, and M.K. Njenga. 2000. Development of highly sensitive and specific enzyme-linked immunosorbent assay based on recombinant matrix protein for detection of avian pneumovirus antibodies. *Journal of Clinical Microbiology* 38: 4010-4014.

King, D.J., G.D. Kommers, C.C. Brown, and **B.S. Seal**. 2001. Virulence of pigeon Newcastle Disease Virus isolates for chickens. Proceedings of the Fiftieth Western Poultry Disease Conference, p.15-16.

King, D.J. and **B.S. Seal**. 2000. Current understanding of molecular correlates of virulence in Newcastle Disease Virus. *International Hatchery Practice*. 14(7)(Supl.): 10-11.

Kommers, G.D., D.J. King, **B.S. Seal**, and C.C. Brown. 2000. Pathogenesis of five

different pigeon-origin isolates of Newcastle disease virus for domestic chickens. *Veterinary Pathology* 37(5): 546(Abstract No.94).

Locke, D.P., H.S. Sellers, J.M. Crawford, S. Schultz-Cherry, D.J. King, R.J. Meinersmann, and **B.S. Seal**. 2000. Newcastle disease virus phosphoprotein gene analysis and transcriptional editing in avian cells. *Virus Research* 69: 55-68.

Seal, B.S. and H.S. Sellers. 2001. Avian paramyxoviruses evolve independently of their mammalian counterparts and deserve a genus designation among the subfamily Paramyxovirinae in the family Paramyxoviridae. Program of the Ninth Annual Meeting of the Society for Molecular Biology and Evolution, Athens, GA. July, 7-10, 2001. (Abstract).

Seal, B.S. 2001. Molecular evolution of Newcastle disease virus and the application of molecular diagnostics. Program of the Respiratory Diseases of Poultry Symposium, American Association of Avian Pathologists at the American Veterinary Medical Association Annual Meeting in Boston, MA. July 14-18, 2001. (Abstract).

Seal, B.S. 2000. Avian Pneumoviruses and emergence of a new type in the United States. *Animal Health Research Reviews* 1: 67-72.

Seal, B.S. and D.J. King. 2000. Newcastle Disease Virus. Available from: <http://www.els.net> Encyclopedia of Life Sciences.

Sellers, H.S., S. Schultz-Cherry, C.C. Brown, **B.S. Seal**, and D.J. King. 2000. Induction of apoptosis by Newcastle disease virus. Program of the Fifth International Congress of Veterinary Virology, Brescia, Italy. August, 2000. (Abstract).

Wilson, W.C., H.C. Ma, E.H. Venter, A.A. van Djik, **B.S. Seal**, and J.O. Mecham. 2001. Erratum to 'Phylogenetic relationships of bluetongue viruses based on gene S7'. *Virus Research* 73: 201-202.

Suarez DL

Lee, C., C. Song, Y. Lee, I. Mo, M. Garcia, **D.L. Suarez**, and S. Kim. 2000. Molecular and pathogenic characterization of Korean isolates of H9N2 avian influenza virus. *Avian Diseases* 44: 527-535.

Perdue, M.L., **D.L. Suarez**, and D.E. Swayne. 2000. Avian influenza in the 90's. *Poultry and Avian Biology Reviews* 11: 1-20.

Schultz-Cherry, S., J.K. Dybing, N.L. Davis, C. Williamson, **D.L. Suarez**, R. Johnston, and M.L. Perdue. 2000. Influenza virus (A/HK/156/97) hemagglutinin expressed by an alphavirus replicon system protects chickens against lethal infection with Hong Kong-origin H5N1 viruses. *Virology* 255: 55-59.

Suarez, D.L., D.E. Swayne, and D.J. King. 2001. The ongoing threat of avian influenza and Newcastle disease to the U.S. poultry industry. Proceedings of the Fiftieth Western Poultry Disease Conference, p.1-7.

Suarez, D.L. 2000. Influenza of poultry and swine. *Le Medecin Vétérinaire Du Quebec* 30: 187-192.

Suarez, D.L. and S. Schultz-Cherry. 2000. The effect of eukaryotic expression vectors and adjuvants on DNA vaccines in chickens using an avian influenza model. *Avian Diseases* 44: 861-868.

Suarez, D.L., D.E. Swayne, and D.J. King. 2001. The ongoing threat of avian influenza and Newcastle disease to the US poultry industry. Proceedings of the 50th Western Poultry Disease Conference, p.1-7.

Swayne, D.E., **D.L. Suarez**, and D.J. King. 2000. Avian influenza (AI) and velogenic Newcastle disease (VND) update. Proceedings of 35th National Meeting of Poultry Health and Processing, p.37-45.

Swayne, D.E. and **D.L. Suarez**. 2000. Update on international incidences and outbreaks of avian influenza 1999-2000. Proceedings of the US Animal Health Association 104: 600-604.

Swayne DE

King, D.J. and **D.E. Swayne**. 2000. Newcastle disease update: International ND problems. Proceedings of the US Animal Health Association 104: 593-596.

Perdue, M.L., D.L. Suarez, and **D.E. Swayne**. 2000. Avian influenza in the 90's. Poultry and Avian Biology Reviews 11: 1-20.

Perkins, L.E. and **D.E. Swayne**. 2001. Clinicopathologic Aspects of Animal and Zoonotic Diseases of Bioterrorism: Avian Influenza. Clinics of North America 21: 549-590.

Perkins, L.E. and **D.E. Swayne**. 2001. Pathobiology of A/Chicken/Hong Kong/220/97 (H5N1) avian influenza virus in seven Gallinaceous species. Veterinary Pathology 38: 149-164.

Suarez, D.L., **D.E. Swayne**, and D.J. King. 2001. The ongoing threat of avian influenza and Newcastle disease to the US poultry industry. Proceedings of the 50th Western Poultry Disease Conference, p.1-7.

Swayne, D.E. 2001. National and international avian influenza outbreaks and strategies for control. Proceedings

AAAP Symposium on Respiratory Diseases of Poultry, 49-55.

Swayne, D.E. 2000. Avian influenza (AI) and velogenic Newcastle disease (VND) update. Proceedings of the 35th National Meeting on Poultry Health and Processing, p.37-45.

Swayne, D.E., J.R. Beck, M.L. Perdue, and C.W. Beard. 2001. Efficacy of vaccines in chickens against highly pathogenic Hong Kong H5N1 avian influenza. Avian Diseases 45: 355-365.

Swayne, D.E., J.R. Beck, and S. Zaki. 2000. Pathogenicity of West Nile virus for turkeys. Avian Diseases 44: 932-937.

Swayne, D.E. and D.L. Suarez. 2000. Update on international incidences and outbreaks of avian influenza 1999-2000. Proceedings of the US Animal Health Association 104: 600-604.

Swayne, D.E., D.L. Suarez, and D.J. King. 2000. Avian influenza (AI) and velogenic Newcastle disease (VND) update. Proceedings of 35th National Meeting of Poultry Health and Processing, p.37-45.

Tumpey T

Katz, J.M., X. Lu., M. Frace, T. Morken, S.R. Zaki, and **T.M. Tumpey**. 2000. Pathogenesis of and immunity to avian influenza A H5 viruses. Biomedicine Pharmacotherapy 54: 178-187.

Tumpey, T.M., X. Lu, T. Morken, S.R. Zaki, and J.M. Katz. 2000. Depletion of lymphocytes and diminished cytokine production in mice infected with a highly virulent influenza A virus isolated from humans. Journal of Virology 73: 6105-6116.

Tumpey, T.M., M. Renshaw, J. Clements, and J. M. Katz. 2001. Mucosal delivery of inactivated influenza vaccine induces B cell

dependent heterosubtypic protection against a lethal influenza A H5N1 virus. *Journal of Virology* 75: 5141-5150.

West Lafayette, Indiana

Livestock Behavior Research Unit

<http://ag.ansc.purdue.edu/usda/index.html>

Eicher SD

Cheng, H.W., **S.D. Eicher**, Y. Chen, and W.B. Muir. 2001. Effect of genetic selection for group productivity and longevity on immunological and hematological parameters of chickens. *Journal of Poultry Science* 80: 1079-1086.

Eicher, S.D. 2000. Transportation of cattle in the dairy industry: current research and future directions. *Journal of Dairy Science* 84: 19-23.

Eicher, S.D., J. Morrow-Tesch, J.L. Albright, and R.E. Williams. 2001. Tail-docking alters fly numbers, fly avoidance behaviors and cleanliness, but not

physiological measures. *Journal of Dairy Science* 84: 1822- 1828.

McKee, C.A., J.A. Carroll, **S.D. Eicher**, M.E. Zannelli, L.A. Beausang, and R.L. Matteri. 2001. Supplemental vitamin C and beta-glucan alter growth and the LPS-induced immunological responses in young pigs. *Journal of Animal Science* 79(Suppl.2): 34.

Cheng HW

Cheng, H.W., S.D. Eicher, Y. Chen, P.B. Singleton, W.B. Muir. 2001. Effect of genetic selection for group productivity and longevity on immunological and hematological parameters of chickens. *Journal of Poultry Science*. 80: 1079-1086.

Crop Production and Pest Control Research

<http://www.btny.purdue.edu/USDA-ARS/CPPCRU/CPPCRU.html>

Nielsen NC

Helm, R.M., G. Cockrell, C. Connaughton, H.A. Sampson, G.A. Bannon, V. Beilinson, D. Livingstone, **N.C. Nielsen**, and A.W. Burks. 2000. A soybean G2 glycinin allergen. 1. Identification and characterization. *International Archives of Allergy and Immunology* 123(3): 205-212.

Helm, R.M., G. Cockrell, C. Connaughton, H.A. Sampson, G.A. Bannon, V. Beilinson, **N.C. Nielsen**, and A.W. Burks. 2000. A soybean G2 glycinin allergen. 2. Epitope mapping and three-dimensional modeling. *International Archives of Allergy and Immunology* 123(3): 213-219.

Ames, Iowa

National Animal Disease Center

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=36-25-30-00>

Andersen AA

Everett, K.D.E. and **A.A. Andersen**. 2001. Intergenic spacer target sequence for detection and distinguishing Chlamydial strains. 2001. U.S. Patent 6,261,769.

Sharma, M. and **A.A. Andersen**. 2000. Isolation and growth of Chlamydiae in cell cultures. Indian Journal of Animal Sciences 70(11): 1127-1128.

Bricker BJ

Redkar, R., S. Rose, **B. Bricker**, and V. DelVecchio. 2001. Real-time detection of *Brucella abortus*, *Brucella melitensis* and *Brucella suis*. Molecular Cellular Probes 15: 43-52.

Briggs RE

Briggs, R.E. Mucosal Vaccination Against Respiratory Diseases of Ruminants. 2000. Annual Meeting of the United States Animal Health Association. (Invited Oral Presentation).

Frank, G.H., **R.E. Briggs**, G.C. Duff, R.W. Loan, and C. W. Purdy. 2002. Effects of vaccination prior to transit and administration of florfenicol at time of arrival in a feedlot on the health of transported calves and detection of *Mannheimia haemolytica* in nasal secretions. American Journal of Veterinary Research 63(2): 251-256.

Frank, G.H., **R.E. Briggs**, R.W. Loan, C. W. Purdy, E.S. Zehr. 1999. Colonization of the nasopharynx of cattle by *Pasteurella haemolytica*: Effect of tilmicisin.

Haemophilus, *Actinobacillus*, and *Pasteurella* International Conference, p.10.

Fulton, R.W., C. W. Purdy, A.W. Confer, J.T. Saliki, R.W. Loan, **R.E. Briggs**, and L.J. Burge. 2000. Bovine viral diarrhea viral infections in feeder calves with respiratory disease: Interactions with *Pasteurella spp.*, parainfluenza-3 virus, and bovine respiratory syncytial virus. Canadian Journal of Veterinary Research 64: 151-159.

Jeyaseelan, S., M.S. Kannan, **R.E. Briggs**, P. Thumbikat, and S.K. Maheswaran. 2001. *Mannheimia haemolytica* leukotoxin activates a nonreceptor tyrosine kinase signaling cascade in bovine leukocytes, which induces biological effects. Infection and Immunity 69(10): 6131-6139.

Jeyaseelan, S., M.S. Kannan, Hsuan, S.L., Walseth, T.F., Singh, A.K., **R.E. Briggs**, and S.K. Maheswaran. 2000. Role of arachidonic acid metabolites in *Pasteurella haemolytica* leukotoxin-induced cytotoxicity. Conference for Research Workers in Animal Disease, Abstract #148.

Leite, F., F.J. Brown, M.J. Sylte, **R.E. Briggs**, and C.J. Czuprynski. 2000. Recombinant bovine interleukin-1 beta amplifies the effects of partially purified *Pasteurella haemolytica* leukotoxin on bovine neutrophils in a beta-2-integrin-dependent manner. Infection and Immunity 68: 5581-5586.

Purdy, C.W., R.W. Loan, D.C. Straus, **R.E. Briggs**, and G.H. Frank. 2000. Conglutinin and immunoconglutinin titers in stressed calves in a feedlot. American

Journal of Veterinary Research 61(11): 1403-1409.

Storz, J., M. Lin, C. W. Purdy, V.N. Chouljenko, K.G. Kousoulas, F.W. Enright, W.C. Gilmore, **R.E. Briggs**, and R.W. Loan. 2000. Coronavirus and *Pasteurella* infections in bovine shipping fever pneumonia and Evans criteria for causation. Journal of Clinical Microbiology 38: 3291-3298.

Thumbikat, P., S. Jeyaseelan, **R.E. Briggs**, M.S. Kannan, and S.K. Maheswaran. 2000. Functional dissection of *Pasteurella haemolytica* leukotoxin mutants. Conference for Research Workers in Animal Disease. 2000. Abstract #147.

Brockmeier SL

Brockmeier, S.L., M.V. Palmer, and S.R. Bolin. 2000. Effects of intranasal inoculation of porcine reproductive and respiratory syndrome virus, *Bordetella bronchiseptica*, or a combination of both organisms in pigs. American Journal of Veterinary Research 61(8): 892-899.

Brockmeier, S.L., M.V. Palmer, S.R. Bolin, and R.B. Rimler. 2001. Effects of intranasal inoculation with *Bordetella bronchiseptica*, porcine reproductive and respiratory syndrome virus, or a combination of both organisms on subsequent infection with *Pasteurella multocida* in pigs. American Journal of Veterinary Research 62(4): 521-525.

Brockmeier, S.L., K.B. Register, and R.A. Kunkle. 2000. Evaluation of the virulence of a dermonecrotic toxin mutant of *Bordetella bronchiseptica* in swine. Proceedings of the 16th International Congress of the Pig Veterinary Society, p. 480.

Register, K.B., T.F. Ducey, **S.L. Brockmeier**, and D.W. Dyer. 2001.

Reduced virulence of a *Bordetella bronchiseptica* siderophore mutant in neonatal swine. Infection and Immunity 69(4): 2137-2143.

Brogden KA

Ackermann, M.R. and **K.A. Brogden**. 2000. Review. Response of the ruminant respiratory tract to *Mannheimia (Pasteurella) haemolytica*. Microbes and Infection. 2: 1079-1088.

Brogden, K.A., V.C. Kalfa, M.R. Ackermann, D.E. Palmquist, P.B. McCray Jr., and B.F. Tack. 2001. The ovine cathelicidin SMAP29 kills ovine respiratory pathogens in vitro and in an ovine model of pulmonary infection. Antimicrobial Agents and Chemotherapy 45: 331-334.

Caverly, J.M., Z.A. Radi, Andreasen, C.B., R.A. Dixon, **K.A. Brogden**, and M.R. Ackermann. 2001. Comparison of bronchoalveolar lavage fluid obtained from *Mannheimia haemolytica*-inoculated calves with and without prior treatment with the selectin inhibitor TBC1269. American Journal of Veterinary Research 62: 665-672.

Lee, H.Y., M.E. Kehrli, Jr., **K.A. Brogden**, J.M. Gallup, and M.R. Ackermann. 2000. Influence of beta 2-integrin adhesion molecule expression and pulmonary infection with *Pasteurella haemolytica* on cytokine gene expression in cattle. Infection and Immunity 68(7): 4274-4281.

Radi, Z.A., J.M. Caverly, R.A. Dixon, **K.A. Brogden**, and M.R. Ackermann. 2001. Effects of the synthetic selectin inhibitor TBC1269 on tissue damage during acute *Mannheimia haemolytica*-induced pneumonia in neonatal calves. American Journal of Veterinary Research 62: 17-22.

Ramirez-Romero, R., **K.A. Brogden**, J.M. Gallup, R.A. Dixon, and M.R. Ackermann. 2000. Reduction of pulmonary mast cells in areas of acute inflammation in calves with *Mannheimia (Pasteurella) haemolytica* pneumonia. *Journal of Comparative Pathology* 123: 29-35.

Zhao, C., T. Nguyen, L. Liu, R.E. Sacco, **K.A. Brogden**, and R.I. Lehrer. 2001. Gallinacin-3, an inducible epithelial beta-defensin of the chicken. *Infection and Immunity* 69(4): 2684-2691.

Frank GH

Al-Ghamdi, G.M., T.R. Ames, J.C. Baker, R. Walker, C.C.L. Chase, **G.H. Frank**, and S.K. Maheswaran. 2000. Serotyping of *Mannheimia (Pasteurella) haemolytica* isolates from the upper Midwest United States. *Journal of Veterinary Diagnostic Investigations* 12: 576-578.

Frank, G.H., R.E. Briggs, R.W. Loan, C. W. Purdy, E.S. Zehr. 1999. Colonization of the nasopharynx of cattle by *Pasteurella haemolytica*: Effect of tilmicosin. *Haemophilus, Actinobacillus*, and *Pasteurella* International Conference, p. 10.

Frank, G.H. and G.C. Duff. 2000. Effects of tilmicosin phosphate administered prior to transport or at time of arrival and feeding of chlortetracycline after arrival in a feedlot on *Mannheimia haemolytica* organisms in nasal secretions of transported steers. *American Journal of Veterinary Research* 61: 1479-1483.

Purdy, C.W., R.W. Loan, D.C. Straus, R.E. Briggs, and **G.H. Frank**. 2000. Conglutinin and immunoconglutinin titers in stressed calves in a feedlot. *American Journal of Veterinary Research* 61(11): 1403-1409.

Zehr, E.S., **G.H. Frank**, and L.B. Tabatabai. 2001. Characterization of Proteins Extracted from the Outer

Membrane of *Mannheimia haemolytica* by sepharose- bound bovine holotransferrin. 101st General Meeting of the American Society for Microbiology Abstract #100.

Harp JA

Chen, W., **J.A. Harp**, and A.G. Harmsen. 2003. *Cryptosporidium parvum* infection in gene targeted B cell deficient mice. *Journal of Parasitology* 89: 391-393.

Harp, J.A., D. Marancik, and J. Oesper. 2002. In vitro comparison of bovine blood and milk lymphocyte adherence to high endothelial venules in lymph nodes of murine origin. *Conference of Research Workers in Animal Disease* 83: 83P.

Harp, J.A., J.C. Oesper, T.A. Casey. 2003. Oral dosing of mice with *Escherichia coli* expressing recombinant interferon gamma reduces *Cryptosporidium parvum* infection. *Conference of Research Workers in Animal Disease* 84: 93P. (Abstract).

Harp, J.A., J.R. Stabel, B.A. Pesch, and J.P. Goff. 2004. Expression of adhesion molecules on milk and blood leukocytes from periparturient dairy cattle with Johne's disease. *Veterinary Immunology and Immunopathology* 98: 69-76.

Harp, J.A., J.R. Stabel, B.A. Pesch, and J.P. Goff. 2001. Expression of adhesion molecules on milk and blood leukocytes from periparturient cows. *Conference of Research Workers in Animal Disease* 82: 62P.

Harp, J.A., J.R. Stabel, T.E. Rahner, B.A. Pesch, and J.P. Goff. 2003. Comparison of adhesion molecule expression on lymphocytes from blood and milk of normal cows and cows with Johne's disease. *Conference of Research Workers in Animal Disease* 84: 71P.

Harp, J.A., T.E. Waters, and J.P. Goff. 2004. Lymphocyte subsets and adhesion molecule expression in milk and blood of periparturient dairy cattle. *Veterinary Immunology and Immunopathology*. In press.

Heidari, M., **J.A. Harp**, and M.E. Kehrli. 2001. Expression, purification, and in vitro biological activities of recombinant bovine granulocyte colony stimulating factor. *Veterinary Immunology and Immunopathology* 81: 45-57.

Kehrli, M.E. and **J.A. Harp**. 2001. Immunity in the mammary gland. *Veterinary Clinics of North America Food Animal Practice* 17: 495-516.

Kimura, K., J.P. Goff, M.E. Kehrli, **J.A. Harp**, and B.J. Nonnecke. 2002. Effects of mastectomy on the composition of peripheral blood mononuclear cell populations in periparturient dairy cows. *Journal of Dairy Science* 85: 1437-1444.

Nonnecke, B.J., M.A. Fowler, B.A. Pesch, B.L. Miller, R.L. Horst, T.E. Johnson, H.B. Perry, and **J.A. Harp**. 2000. Function and composition of circulating leukocyte populations from milk-replacer-fed, neonatal calves: Effects of dietary vitamins A and E. *ADSA Annual Mtg.*, Baltimore, Maryland, July 24-28, 2000. (Abstract).

Oesper, J.C., S.M. Preszler, and **J.A. Harp**. 2003. Production of recombinant bovine gamma interferon by *Escherichia coli* for use in an oral vaccine to control *Cryptosporidium parvum* infection. *Conference of Research Workers in Animal Disease* 84: 91P.

Pesch, B.A. and **J.A. Harp**. 2000. Comparative expression of homing receptors on blood and milk cells of dairy cattle. *Great Lakes International Imaging and Flow Cytometry Association Annual Meeting*, Detroit, October 13-15, 2000.

Rahner, T.E., **J.A. Harp**, and J.P. Goff. 2003. Expression of homing receptors on blood and milk granulocytes during the parturient period of dairy cattle. *Great Lakes International Imaging and Flow Cytometry Association Annual Meeting*, Madison, WI, October 3-5, 2003.

Smith, J.M., B.J. Nonnecke, M.E. Van Amburgh, B.A. Pesch, and **J.A. Harp**. 2000. Effects of dietary conjugated linoleic acid (CLA) on the composition and function of peripheral blood mononuclear leukocyte populations in heifer calves. *ADSA/ASAS Annual Mtg.*, Indianapolis, Indiana, July 1-5, 2001.

Sonea, I.M., M.K. Schwarz, and **J.A. Harp**. 2001. Regulation of interferon- γ production by substance P following infection with *Cryptosporidium parvum*. *Conference of Research Workers in Animal Disease* 82: 54P.

Stabel, J.R., J.P. Goff, K. Kimura, **J.A. Harp**, and R.H. Whitlock. 2002. Effects of supplemental energy on periparturient immunosuppression in dairy cows with Johne's disease. *7th Intl. Colloquium on Paratuberculosis*, Bilboa, Spain, June 11-14, 2002. (Abstract).

Waters, W.R., M.V. Palmer, M.J. Wannemuehler, R.E. Sacco, **J.A. Harp**. 2000. B cells are required for the induction of intestinal inflammatory lesions in TCR α -deficient mice persistently infected with *Cryptosporidium parvum*. *Journal of Parasitology* 86: 1073-1077.

Horst RL

Nonnecke, B.J., M.R. Waldron, T. Nishida, T.R. Overton, and **R.L. Horst**. 2001. Circulating leukocyte populations, serum cytokines, and plasma vitamins A and E in mid-lactation dairy cows infused with varied doses of lipopolysaccharide.

Kehrli ME Jr

Kehrli, M.E. Jr. and J.A. Harp. 2001.
Immunity in the mammary gland.
Veterinary Clinics of North America, The
Food Animal Practice 17(3): 495-516.

Kimura, K., J.P. Goff, **M.E. Kehrli Jr.**,
J.A. Harp, and B.J. Nonnecke. 2002.
Effects of mastectomy on composition of
peripheral blood mononuclear cell
populations in periparturient dairy cows.
Journal of Dairy Science 85(6): 1437-
1444.

Nonnecke, B.J., K. Kimura, J.P. Goff, and
M.E. Kehrli Jr.. 2003. Effects of the
mammary gland on functional capacities of
blood mononuclear leukocyte populations
from periparturient cows. Journal of Dairy
Science 86(7): 2359-68.

Kimura K

Kimura, K., J.P. Goff, M.E. Kehrli Jr.,
J.A. Harp, and B.J. Nonnecke. 2002.
Effects of mastectomy on composition of
peripheral blood mononuclear cell
populations in periparturient dairy cows.
Journal of Dairy Science June 85(6): 1437-
1444.

Kimura, K., J.P. Goff, M.E. Kehrli Jr, and
T.A. Reinhardt. 2002. Decreased
neutrophil function as a cause of retained
placenta in dairy cattle. Journal of Dairy
Science 85(3): 544-50.

Nonnecke, B.J., **K. Kimura**, J.P. Goff, and
M.E. Kehrli Jr. Effects of the mammary
gland on functional capacities of blood
mononuclear leukocyte populations from
periparturient cows. Journal of Dairy
Science 2003 Jul, 86(7): 2359-68.

Lager, K.M. and W.L. Mengeling. 2000.
PRRS: Nature of the RNA virus and how it
causes disease. Proceedings of
International Pig Veterinary Society 16th
Congress, p. 538-543.

Mengeling, W.L., S.R. Bolin, **K.M. Lager**,
A.C. Vorwald, and R.D. Wesley. 2000.
Immunology and epidemiology of porcine
reproductive and respiratory syndrome,
pseudorabies, and postweaning
multisystemic wasting syndrome.
Proceedings of 2nd National Meeting of
the Italiana Veterinari per Animali da
Reddito, p. 86-92.

Mengeling, W.L., D.F. Clouser, A.C.
Vorwald, and **K.M. Lager**. 2000. Genetic
recombination among strains of porcine
reproductive and respiratory syndrome
virus. Conference of Research Workers in
Animal Disease 81st Annual Meeting,
Abstract No. 134.

Mengeling, W.L., R.C. Cutlip, A.C.
Vorwald, and **K.M. Lager**. 2000. The role
of porcine reproductive and respiratory
syndrome virus and pseudorabies virus in
swine respiratory disease. Proceedings of
2nd National Meeting of the Italiana
Veterinari per Animali da Reddito, p. 93-
97.

Mengeling, W.L. and **K.M. Lager**. 2000.
Porcine reproductive and respiratory
syndrome (PRRS) in the U.S.A.
Proceedings of XVII PanAmerican
Congress of Veterinary Science, p.41-43.

Mengeling, W.L., **K.M. Lager**, K.
Burkhart, A.C. Vorwald, M.B. Roof, and
D.E. Gorcyca. Porcine reproductive and
respiratory syndrome virus vaccine based
on isolate JA-142. U.S. Patent Application
S/N 09/298,110.

Mengeling, W.L., **K.M. Lager**, A.C. Vorwald, R.D. Wesley, and C. Prieto. 2000. Porcine reproductive and respiratory syndrome: vaccines and immunity. *Magazyn Weterynaryjny*, p. 16-18.

Mengeling, W.L., A.C. Vorwald, N.A. Cornick, **K.M. Lager**, and H.W. Moon. 2000. The use of porcine alveolar macrophages for detecting Shiga Toxin. *Proceedings of American Association of Veterinary Laboratory Diagnosticians 43rd Annual Meeting*, p. 82.

Yaeger, M., J. Benson, J. Christopher-Hennings, and **K. Lager**. 2000. Evaluation of the sensitivity of fetal serology, IHC, and PCR for the diagnosis of PRRSV in fetuses. *Proceedings of American Association of Veterinary Laboratory Diagnosticians 43rd Annual Meeting*, p. 12.

Lehmkuhl HD

Boyce, W.M., L.W. Woods, K. Keel, N.J. MacLachlan, C.O. Porter, and **H.D. Lehmkuhl**. 2000. An epizootic of adenovirus induced hemorrhagic disease in captive black-tailed deer (*Odocoileus hemionus*). *Journal of Zoo and Wildlife Medicine* 31(3): 370-373.

Cutlip, R.C., J.M. Miller, A.N. Hamir, J. Peters, M.M. Robinson, A.L. Jenny, **H.D. Lehmkuhl**, W.D. Taylor, F.D. Bisplinghoff. 2001. Resistance of cattle to scrapie by the oral route. *Canadian Journal of Veterinary Research* 65(2): 131-132.

Debey, B.M., and **H.D. Lehmkuhl**. 2000. Fatal infections in lambs associated with an adenovirus. 43rd Annual Meeting, American Association of Veterinary Laboratory Diagnosticians, p. 16.

Lehmkuhl, H.D., B.M. Debey, and R.C. Cutlip. 2001. A new serotype adenovirus isolated from a goat in the United States.

Journal of Veterinary Diagnostic Investigation 13(3): 195-200.

Lehmkuhl, H.D. and L.W. Woods 2001. Characterization of a new adenovirus species isolated from black-tailed deer in California. *Archives of Virology* 146(6): 1187-1196.

Lehmkuhl, H.D., B.M. Debey, and R.C. Cutlip. 2001. A new serotype adenovirus isolated from a goat in the United States. *Journal of Veterinary Diagnostic Investigation* 13(3): 195-200.

Saliki, J.T., S.L. Caseltine, R.W. Fulton, and **H.D. Lehmkuhl**. 2000. Isolation of bovine adenovirus type 7 from calves with respiratory disease. 43rd Annual Meeting, American Association of Veterinary Laboratory Diagnosticians, p. 88.

Woods, L.W., P.K. Swift, **H.D.**

Lehmkuhl, P.H. Chiu, R.S. Hanley, R.W. Nordhausen, M.H. Stillian, and M.L. Drew. 2001. Experimental adenovirus hemorrhagic disease in white-tailed deer fawns. *Journal of Wildlife Diseases* 37(1): 153-158.

Mengeling WL

Lager, K.M. and **W.L. Mengeling**. 2000. PRRS: Nature of the RNA virus and how it causes disease. *Proceedings of International Pig Veterinary Society 16th Congress*, p. 538-543.

Mengeling, W.L., S.R. Bolin, K.M. Lager, A.C. Vorwald, and R.D. Wesley. 2000. Immunology and epidemiology of porcine reproductive and respiratory syndrome, pseudorabies, and postweaning multisystemic wasting syndrome. *Proceedings of 2nd National Meeting of the Italiana Veterinari per Animali da Reddito*, p. 86-92.

Mengeling, W.L., D.F. Clouser, A.C. Vorwald, and K.M. Lager. 2001. Genetic recombination among strains of porcine reproductive and respiratory syndrome virus. Conference of Research Workers in Animal Disease 81st Annual Meeting, Abstract No. 134.

Mengeling, W.L., R.C. Cutlip, A.C. Vorwald, and K.M. Lager. 2000. The role of porcine reproductive and respiratory syndrome virus and pseudorabies virus in swine respiratory disease. Proceedings of 2nd National Meeting of the Italiana Veterinari per Animali da Reddito, p. 93-97.

Mengeling, W.L. and K.M. Lager. 2000. Porcine reproductive and respiratory syndrome (PRRS) in the U.S.A. Proceedings of XVII PanAmerican Congress of Veterinary Science, p. 41-43.

Mengeling, W.L., K.M. Lager, A.C. Vorwald, R.D. Wesley, and C. Prieto. 2000. Porcine reproductive and respiratory syndrome: vaccines and immunity. *Magazyn Weterynaryjny*, p. 16-18.

Mengeling, W.L., K.M. Lager, K. Burkhardt, A.C. Vorwald, M.B. Roof, and D.E. Gorcyca. Porcine reproductive and respiratory syndrome virus vaccine based on isolate JA-142. U.S. Patent Application S/N 09/298,110.

Mengeling, W.L., A.C. Vorwald, N.A. Cornick, K.M. Lager, and H.W. Moon. 2000. The use of porcine alveolar macrophages for detecting Shiga toxin. Proceedings of American Association of Veterinary Laboratory Diagnosticians 43rd Annual Meeting, p. 82.

Miller JM

Cutlip, R.C., **J.M. Miller**, A.N. Hamir, J. Peters, M.M. Robinson, A.L. Jenny, H.D. Lehmkuhl, W.D. Taylor, and F.D.

Bisplinghoff. 2001. Resistance of cattle to scrapie by the oral route. *Canadian Journal of Veterinary Research* 65(2): 131-132.

Hamir, A. N., R.C. Cutlip, **J.M. Miller**, E.S. Williams, M.J. Stack, M.W. Miller, K.I. O'Rourke, and M.J. Chaplin. 2001. Preliminary findings on the experimental transmission of chronic wasting disease agent of mule deer to cattle. *Journal of Veterinary Diagnostic Investigation* 13: 91-96.

Neill JD

Endsley, J.J., J.F. Ridpath, **J.D. Neill** Cell mediated immunity to bovine viral diarrhea virus in calves protected by maternal antibody. Conference of Research Workers in Animal Disease. 2000. Abstract No. 91.

Leibler-Tenorio, E.M., J.F. Ridpath, I. Greiser-Wilke, J.F. Pohlenz, and **J.D. Neill** Comparison of lesions and distribution of viral antigen in cattle with acute BVDV-2 infection and mucosal disease. European Veterinary Pathology Meeting. 2000. Abstract No. 31.

Neill, J.D. and J.F. Ridpath. 2000. Recombination with a cellular mRNA encoding a DNAJ-like protein results in biotype conversion in genotype 2 bovine viral diarrhea viruses. United States-Japan Natural Resources Animal and Avian Health Panel, Abstract No. 11.

Neill, J.D. and J.F. Ridpath. 2000. Further biochemical characterization of three bovine capases involved in apoptosis of virus-infected cells. Conference of Research Workers in Animal Disease, Abstract No. 130P.

Neill, J.D. and J.F. Ridpath. 2001. Changes in host cell gene expression following infection with bovine viral diarrhea virus genotype 2. American

Society for Virology, Abstract No. W41-1, p.135.

Ridpath, J.F., S. Hietla, S. Sorden, and **J.D. Neill**. 2000. Comparison of diagnostic techniques for detection of acute BVDV infections. Conference of Research Workers in Animal Disease, Abstract No. 125P.

Ridpath, J.F. and **J.D. Neill**. 2000. Characterization of BVDV from North America. United States-Japan Natural Resources Animal and Avian Health Panel, Abstract No. 10.

Ridpath, J.F., **J.D. Neill**, M. Frey, and J.G. Landgraf. 2000. Phylogenetic, antigenic and clinical characterization of type 2 BVDV from North America. *Veterinary Microbiology* 77(1-2): 145-155.

Ridpath, J.F., **J.D. Neill**, J.J. Endsley, J.A. Roth. 2000. Passive immunity and the development of a protective response against BVDV. Conference of Research Workers in Animal Disease, Abstract No. 77P.

Nonnecke BJ

Nonnecke, B.J., R.L. Horst, D.C. Hammell, and S.T. Franklin. 2000. Effects of supplemental vitamin A on retinoic acid concentrations in the plasma of preruminant calves. *International Journal of Vitamin and Nutrition Research* 70(6): 278-286.

Nonnecke, B.J., K. Kimura, J.P. Goff, and M.E. Kehrli Jr. 2003. Effects of the mammary gland on functional capacities of blood mononuclear leukocyte populations from periparturient cows. *Journal of Dairy Science* 86(7): 2359-68.

Nonnecke, B.J., M.P. Robert, J.D. Godkin, D.C. Hammell, and S.T. Franklin. 2001. Influence of supplemental, dietary

vitamin A on retinol-binding protein concentrations in the plasma of preruminant calves. *Journal of Dairy Science* 84: 641-648.

Nonnecke, B.J., M.R. Waldron, T. Nishida, T.R. Overton, and R.L. Horst. 2001. Circulating leukocyte populations, serum cytokines, and plasma vitamins A and E in mid-lactation dairy cows infused with varied doses of lipopolysaccharide. International Animal Agriculture and Food Science Conference, Abstract p. 385.

Smith, J.A., **B.J. Nonnecke**, M.E. van Amburgh, B.A. Pesch, and J.A. Harp. 2001. Effects of dietary conjugated linoleic acid (CLA) on the composition and function of peripheral blood mononuclear leukocyte populations in heifer calves. International Animal Agriculture and Food Science Conference, Abstract No. 1806, p. 436.

Waldron, M.R., T. Nishida, **B.J. Nonnecke**, and T.R. Overton. 2001. Metabolic responses of lactating dairy cows to intravenous infusion of increasing amounts of lipolysaccharide. International Animal Agriculture and Food Science Conference, Abstract p. 385.

Waldron, M.R., T. Nishida, **B.J. Nonnecke**, and T.R. 2001. Overton Hepatic oxidative metabolism in lactating cows is modulated by increasing doses of intravenous lipopolysaccharide. International Animal Agriculture and Food Science Conference, Abstract p. 385.

Dean-Nystrom EA

Dean-Nystrom, E.A., L. Gansheroff, E. Twiddy, H.W. Moon, and A.D. O'Brien. 2000. Passive protection of suckling piglets from *Escherichia coli* O157:H7 infection by vaccination of pregnant sows with intiminO157. International Symposium and Workshop on Shiga Toxin

(Verotoxin)-Producing *Escherichia coli* Infections, p. 175.

Dean-Nystrom, E. A., A.R. Melton-Celsa, J.F. Pohlenz, H.W. Moon, and A.D. O'Brien. 2003. Comparative Pathogenicity of *Escherichia coli* O157 and Intimin-Negative Non-O157 Shiga Toxin-Producing *E. coli* Strains in Neonatal Pigs. *Infection and Immunity* 71(11): 6526-6533.

Olsen SC

Adone, R., F. Ciuchini, and **S.C. Olsen.** 2001. RB51 antigen validation of complement fixation test to detect in a blind assay calves of Iowa (USA) vaccinated with *Brucella abortus* RB51. *Journal of Clinical Microbiology* 8: 385-387.

Adone, R., F. Ciuchini, and **S.C. Olsen.** 2000. Antibody responses of heifers vaccinated with *Brucella abortus* RB51 monitored by complement fixation test. *Brucellosis 2000, including the 53rd Brucellosis Research Conference: Scientific program and abstracts.* September 7-9. Nimes, France 2000:38.

Capsel, R.L., **S.C. Olsen**, N.F. Cheville, and C.O. Thoen. 2000. Survival of *Brucella abortus* strain RB51 lyophilized and as a liquid vaccine under different storage conditions. *Biologicals* 28: 209-216.

Januszewski, M.C., **S.C. Olsen**, R.G. McLean, L. Clark, and J.C. Rhyan. 2001. Experimental infection of nontarget species of rodents and birds with *Brucella abortus* strain RB51 vaccine. *Journal of Wildlife Diseases* 37: 532-537.

Olsen, S.C. 2000. Biosafety Issues in Animal Care and Use. American Biological Science Safety Association, Washington, DC, October 22.

Olsen, S.C. 2000. Comparison of immune responses to ballistic or hand delivery of *Brucella abortus* strain RB51 vaccine to bison. *Proceedings of the Conference on Research Workers in Animal Disease*, p.61.

Olsen, S.C. 2000. Update on brucellosis research at NADC at the Greater Yellowstone Interagency Brucellosis Committee Meeting in Jackson, WY on November 1, 2000.

Olsen, S.C. 2000. Immune responses and protection of brucellosis vaccines. USDA/APHIS/Veterinary Services, Designated Brucellosis Epidemiologist Training, Ames, IA, on August 11, 2000.

Olsen, S.C. 2000. Safety and efficacy of the *Brucella abortus* strain RB51 vaccine. 35th Meeting of the United States-Japan Cooperative Program in Natural Resources at the National Veterinary Services Laboratory, September 26, 2000.

Olsen, S.C. and I.K. Lee. 2000. Effect of recombinant interleukin-12 on immune responses to brucellosis vaccines. *Research Workers in Animal Diseases Conference Proceedings*, p. 71.

Palmer MV

Brockmeier, S.L., **M.V. Palmer**, and S.R. Bolin. 2000. Effects of intranasal inoculation of porcine reproductive and respiratory syndrome virus, *Bordetella bronchiseptica*, or a combination of both organisms in pigs. *American Journal of Veterinary Research*. 2000 61(8): 892-899.

Brockmeier, S.L., **M.V. Palmer**, S.R. Bolin, and R.B. Rimler. 2001. Effects of intranasal inoculation with *Bordetella bronchiseptica*, porcine reproductive and respiratory syndrome virus, or a combination of both organisms on subsequent infection with *Pasteurella*

multocida in pigs. American Journal of Veterinary Research 62(4): 521-525.

Reinhardt TA

Kimura, K., J.P. Goff, and T.A.

Reinhardt. 2001. Parturition blunts calcium signals in immune cells. Federation of American Societies for Experimental Biology Conference, Abstract p. 136.

Kimura, K., J.P. Goff, **T.A. Reinhardt**, and S. Sato. 2001. Association between retained placenta and blood interleukin-8 concentration and lack of association of retained placenta with energy and calcium metabolic profiles. International Animal Agriculture and Food Science Conference, Abstract p. 151.

Register KB

Brockmeier, S.L., **K.B. Register**, and R.A. Kunkle. 2000. Evaluation of the virulence of a dermonecrotic toxin mutant of *Bordetella bronchiseptica* in swine. Proc. 16th International Congress of the Pig Veterinary Society, p. 480.

Raffell, T.R., **K.B. Register**, and L.M. Temple. 2000. Incidence of *Bordetella avium* in wild birds as measured by serology and tracheal cultures. 100th Meeting of the American Society for Microbiology, Abstract #C5, p. 140.

Register, K.B. 2001. Genetic polymorphism and its implications in *Bordetella bronchiseptica*. Respiratory Diseases V. Pig Progress 16: 20-21.

Register, K.B. 2001. Novel genetic and phenotypic heterogeneity in *Bordetella bronchiseptica* pertactin. Infection and Immunity 69(3): 1917- 1921.

Register, K.B. 2000. RFLP and sequence analysis reveals heterogeneity among the

pertactin genes of *Bordetella bronchiseptica* vaccine and field strains. Proceedings of the 16th Congress of the International Pig Veterinary Society, p. 481.

Register, K.B. 2000. Recent advances in understanding the role of *Bordetella bronchiseptica* and its virulence factors in swine respiratory disease. Proc. 16th Congress of the International Pig Veterinary Society, p. 483-488.

Register, K.B., T.F. Ducey, S.L. Brockmeier, and D.W. Dyer. 2001. Reduced virulence of a *Bordetella bronchiseptica* siderophore mutant in neonatal swine. Infection and Immunity 69(4): 2137-2143.

Register, K.B., R.E. Sacco, and G. Foster. 2000. Ribotyping and restriction endonuclease analysis reveal a novel clone of *Bordetella bronchiseptica* in seals. Journal of Veterinary Diagnostic Investigation 12(6): 535-540.

Sacco, R.E., **K.B. Register**, and G.E. Nordholm. 2000. Restriction endonuclease analysis discriminates *Bordetella bronchiseptica* isolates. Journal of Clinical Microbiology 38(12): 4387-4393.

Richt JA

Kiermayer, S., I. Kraus, **J.A. Richt**, W. Garten, and M. Eickmann. 2002. Identification of the amino terminal subunit of the glycoprotein of Borna disease virus. FEBS Letters 531(2): 255-258.

Richt, J.A., K.M. Lager, B.H. Janke, R.D. Woods, R.G. Webster, and R.J. Webby. 2003. Pathogenic and antigenic properties of phylogenetically distinct reassortant H3N2 swine influenza viruses cocirculating in the United States. Journal

of Clinical Microbiology 41(7): 3198-3205.

Mueller, C.A., **J.A. Richt**, R. Meyermann, M. Deininger, and H. Schluesener. 2003. Accumulation of the proinflammatory cytokine endothelial-monocyte-activating polypeptide II in ramified microglial cells in brains of Borna virus infected Lewis rats. *Neuroscience Letters* 339(3): 215-218.

Wolff, T., G. Unterstab, G. Heins, **J.A. Richt**, and M. Kann. 2002. Characterization of an unusual importin alpha binding motif in the Borna disease virus p10 protein that directs nuclear import. *Journal of Biological Chemistry* 277(14): 12151-12157.

Ridpath JF

Endsley, J.J., **J.F. Ridpath**, and J.D. Neill. 2000. Cell mediated immunity to bovine viral diarrhea virus in calves protected by maternal antibody. Conference of Research Workers in Animal Disease, Abstract No. 91.

Flores, E.F., L.H. Gil, S.A. Botton, R. Weiblen, **J.F. Ridpath**, L.C. Kreutz, C. Pilati, D. Driemeyer, V. Moojen, and A.C. Wendelstein. 2000. Clinical, pathological and antigenic aspects of bovine viral diarrhea virus (BVDV) type 2 isolates identified in Brazil. *Veterinary Microbiology* 77(1-2): 175-183.

Leibler-Tenorio, E.M., **J.F. Ridpath**, I. Greiser-Wilke, J.F. Pohlenz, and J.D. Neill. 2000. Comparison of lesions and distribution of viral antigen in cattle with acute BVDV-2 infection and mucosal disease. European Veterinary Pathology Meeting, Abstract No. 31.

Neill, J.D. and **J.F. Ridpath**. 2001. Changes in host cell gene expression following infection with bovine viral

diarrhea virus genotype 2. American Society for Virology Abstract No. W41-1, p. 135.

Neill, J.D. and **J.F. Ridpath**. 2000. Further biochemical characterization of three bovine capases involved in apoptosis of virus-infected cells. Conference of Research Workers in Animal Disease Abstract No. 130P.

Neill, J.D. and **J.F. Ridpath**. 2000. Recombination with a cellular mRNA encoding a DNAJ-like protein results in biotype conversion in genotype 2 bovine viral diarrhea viruses. United States-Japan Natural Resources Animal and Avian Health Panel Abstract No. 11.

Ridpath, J.F. 2001. Detection and molecular typing of ruminant pestiviruses. Proceedings of Conference on Pestivirus Contamination of Bovine Sera, p. 19-24.

Ridpath, J.F., S. Hietla, S. Sorden, and J.D. Neill. 2000. Comparison of diagnostic techniques for detection of acute BVDV infections. Conference of Research Workers in Animal Disease Abstract No. 125P.

Ridpath, J.F. and J.D. Neill. 2000. Characterization of BVDV from North America. United States-Japan Natural Resources Animal and Avian Health Panel Abstract No. 10.

Ridpath, J.F., J.D. Neill, J. Endsley, and J.A. Roth. 2000. Passive immunity and the development of a protective response against BVDV. Conference of Research Workers in Animal Disease Abstract No. 77P.

Ridpath, J.F., J.D. Neill, M. Frey, and J.G. Landgraf. 2000. Phylogenetic, antigenic and clinical characterization of type 2 BVDV from North America. *Veterinary Microbiology* 77(1-2): 145-155.

Stoffregen, B., S.R. Bolin, **J.F. Ridpath**, and J. Pohlenz. 2000. Morphologic lesions in type 2 BVDV infections experimentally induced by strain BVDV2-1373 recovered from a field case. *Veterinary Microbiology* 77(1-2): 157-162.

Van Campen, H., **J. Ridpath**, E. Williams, J. Cavender, J. Edwards, S. Smith, and H. Sawyer. 2001. Isolation of bovine viral diarrhea virus from a free-ranging mule deer in Wyoming. *Journal of Wildlife Diseases* 37(2): 306-311.

Vilcek, S., **J.F. Ridpath**, H. van Campen, and J. Cavender. 2001. Identification of a novel pestivirus genotype isolated from pronghorn antelope. *American Society for Virology Abstract No. P25-9*, p. 196.

Sacco RE

Butler, J.E., N. Wertz, J. Sun, and **R.E. Sacco**. 2004. Comparison of the expressed porcine V β and J β repertoire of thymocytes and peripheral T cells. (submitted).

Brogden, K.A., M. Heidari, **R.E. Sacco**, D. Palmquist, J.M. Guthmiller, G.K. Johnson, H.P. Jia, P.B. McCray Jr., and B.F. Tack. 2003. Defensin induced adaptive immunity and its potential in preventing periodontal disease. *Oral Microbiology and Immunology* 18: 95-99.

Register, K.B., **R.E. Sacco**, G. Foster. 2000. Ribotyping and restriction endonuclease analysis reveal a novel clone of *Bordetella bronchiseptica* in seals. *Journal of Veterinary Diagnostic Investigation* 12(6): 535-540.

Sacco, R.E., M. Hagen, M. Sandor, J.V. Weinstock, and R.G. Lynch. 2002. Established Th1 granulomatous responses induced by active *Mycobacterium avium* infection switch to Th2 following

challenge with *Schistosoma mansoni*. *Clinical Immunology* 104:274-281.

Sacco, R.E., D. Hutto, W.R. Waters, L. Xiasong, M.E. Kehrli Jr., F.A. Zuckermann, and M.J. Wannemuehler. 2000. Reduction in inflammation following blockade of CD18 or CD29 adhesive pathways during the acute phase of a spirochetal-induced colitis in mice. *Microbial Pathogenesis* 29(5): 289-299.

Sacco, R.E., K.B. Register, and G.E. Nordholm. 2000. Restriction endonuclease analysis discriminates *Bordetella bronchiseptica* isolates. *Journal of Clinical Microbiology* 38(12): 4387-4393.

Sacco, R.E., W.R. Waters, K.M. Rudolph, and M.L. Drew 2004. Comparative nitric oxide production by LPS-stimulated monocyte-derived macrophages from *Ovis canadensis* and *Ovis aries*. (submitted).

Sonea, I.M., A.E. Jergens, **R.E. Sacco**, Y. Niyo, E.A. Merten, L.A. Kaufman, and P.F. Moore. 2000. Flow cytometric analysis of canine colonic and small intestinal lymphocytes obtained by endoscopic biopsy in the healthy dog. *Veterinary Immunology and Immunopathology* 77: 103-119.

Waters, W.R., R. Hontecillas, **R.E. Sacco**, F.A. Zuckermann, K.R. Harkins, J. Bassaganya-Riera, and M.J. Wannemuehle. 2000. Antigen-specific proliferation of porcine CD8 $\alpha\alpha$ cells to an extracellular bacterial pathogen. *Immunology* 101: 333-341.

Waters, W.R., M.V. Palmer, S.C. Olsen, **R.E. Sacco**, and D.L. Whipple. 2003. Immune responses of elk to *Mycobacterium bovis* bacille Calmette Guerin vaccination. *Vaccine* 21:1518-1526.

Waters, W.R., M.V. Palmer, **R.E. Sacco**, and D.L. Whipple. 2002. Nitric oxide production as an indication of *Mycobacterium bovis* infection of white-tailed deer (*Odocoileus virginianus*). Journal of Wildlife Disease 38: 338-343.

Waters, W.R., M.V. Palmer, M.J. Wannemuehler, **R.E. Sacco**, and J.A. Harp. 2000. B cells are required for the induction of intestinal inflammatory lesions of TCR- α -deficient mice persistently infected with *Cryptosporidium parvum*. Journal of Parasitology 86: 1073-1077.

Waters, W.R., **R.E. Sacco**, Fach, S.J., M.V. Palmer, S.C. Olsen, and T.J. Kreeger. 2002. Analysis of mitogen-stimulated lymphocyte subset proliferation and nitric oxide production by peripheral blood mononuclear cells of captive elk (*Cervus elaphus*). Journal of Wildlife Diseases 38: 344-351.

Zhao, C., T. Nguyen, L. Liu, **R.E. Sacco**, K.A. Brogden, and R.I. Lehrer. 2001. Gallinacin-3, an inducible epithelial beta-defensin of the chicken. Infection and Immunity 69(4): 2684-2691.

Stabel JR

Dargatz, D.A., B.A. Byrum, L.K. Barber, R.W. Sweeney, R.H. Whitlock, W.P. Shulaw, R.H. Jacobson, and **J.R. Stabel**. 2001. Evaluation of a commercial ELISA for diagnosis of paratuberculosis in cattle. Journal of American Veterinary Medical Association 218: 1163-1166.

Pillai, S.R., B.M. Jayarao, J.D. Gummo, E.C. Gue, D. Tiwari, **J.R. Stabel**, and R.H. Whitlock. 2001. Identification and subtyping of *Mycobacterium avium* subsp. *paratuberculosis* and *Mycobacterium avium* subsp. *avium* by randomly amplified polymorphic DNA. Veterinary Microbiology 79: 275-284.

Stabel, J.R. 2001. On-farm batch pasteurization destroys *Mycobacterium paratuberculosis* in waste milk. Journal of Dairy Science 82: 524-527.

Stabel, J.R. and R.H. Whitlock. 2001. An evaluation of a modified interferon-gamma assay for the detection of paratuberculosis in dairy herds. Veterinary Immunology and Immunopathology 79: 69-81.

Stabel TJ

Fitzsimmons, C.J., L. Marklund, **T.J. Stabel**, L. Galina, and C.K. Tuggle. 2001. Evaluation of cross-species hybridization with porcine mRNA to human microarrays. International Conference Plant and Genome Research, p. 212.

Greiner, L.L., T.S. Stahly, and **T.J. Stabel**. 2001. The effect of dietary soy genistein on pig growth and viral replication during a viral challenge. Journal of Animal Science 79: 1272-1279.

Greiner, L.L., T.S. Stahly, and **T.J. Stabel**. 2000. Quantitative relationship of systemic virus concentration on growth and immune response in pigs. Journal of Animal Science 78: 2690-2695.

Meyerholz, D.K. and **T.J. Stabel**. 2000. Preferential cellular adherence of *Salmonella typhimurium* DT104 in a swine gut-loop model. Conference of Research Workers in Animal Disease, Abstract p. 1.

Stabel, T.J., S.R. Bolin, and B.A. Pesch. 2000. A simple and rapid flow cytometric method for detection of porcine cell surface markers. Journal of Immunological Methods 245: 147-152.

Stanton TB

Alt, D.P., D.J. Trott, and **T.B. Stanton**. 2000. Molecular cloning, expression, and DNA sequence analysis of a gene encoding

an outer membrane protein of *Brachyspira* (*Serpulina*) *pilosicoli*. 16th International Pig Veterinary Society Abstract p. 9.

Li, C., L. Corum, D. Morgan, E.L. Rosey, **T.B. Stanton**, and N.W. Charon. 2000. The spirochete FlaA periplasmic flagellar sheath protein impacts flagellar helicity. *Journal of Bacteriology* 182: 6698-6706.

Trott, D.J., R.L. Zuerner, M.J. Wannemuehler, and **T.B. Stanton**. 2000. Monoclonal antibodies reacting with *Brachyspira* (*Serpulina*) *pilosicoli* membrane proteins. 16th International Pig Veterinary Society, Abstract p. 41.

Tabatabai LB

Bartleson, C.J. and **L.B. Tabatabai**. 2001. Identification of iron-regulated outer membrane proteins of *Haemophilus parasuis*. *FASEB Journal* 15(4): A533.

Belzer, C.A., **L.B. Tabatabai**, and G.H. Frank. 2000. Purification and characterization of the *Pasteurella haemolytica* 35 Kilodalton periplasmic iron-regulated protein. *Preparative Biochemistry and Biotechnology* 30: 343-355.

Tabatabai, L.B., J.M. Gephart, and K.W. Post. 2001. Two Lys-C endoprotease peptide fragments from a major immunoreactive protein from *Haemophilus parasuis* are homologous to *H. influenzae* elongation Factor Tu (EFTU). *FASEB Journal* 15(4): A534.

Zehr, E.S., G.H. Frank, and **L.B. Tabatabai**. 2001. Characterization of Proteins Extracted from the Outer Membrane of *Mannheimia haemolytica* by sepharose- bound bovine holotransferrin. 101st General Meeting of the American Society for Microbiology, Abstract #100.

Dorn, A.D., **W.R. Waters**, V.M. Byers, B.A. Pesch, and M.J. Wannemuehler. 2002. Characterization of mitogen-stimulated porcine lymphocytes using multicolor flow cytometry. *Veterinary Immunology and Immunopathology* 87: 1-10.

Endsley, J.J., J.L. Furrer, M.A. Endsley, M.A. McIntosh, A.C. Maue, **W.R. Waters**, D.R. Lee, and D.M. Estes. 2004. Characterization of a bovine homologue of granulysin and NK-lysin with broad spectrum antimicrobial activity. *Journal of Immunology*, in press.

Nonnecke, B.J., **W.R. Waters**, M.R. Foote, R.M. Fowler, R.L. Horst, and B.L. Miller. 2003. Interferon-gamma and TNF-alpha secretion by mononuclear leukocytes from peripheral blood of young and adult cattle vaccinated with *Mycobacterium bovis* BCG: modulation by 1,25-dihydroxyvitamin D3. *International Journal of Vitamin and Nutrition Research*, 73: 235-244.

Palmer, M.V., **W.R. Waters**, D.D. Pedersen, and W.C. Stoffregen. 2004. Vaccination of reindeer with a killed West Nile virus vaccine. *Journal of Wildlife Diseases*, in press.

Palmer, M.V., **W.R. Waters**, D.L. Whipple, R.E. Slaughter, and S.L. Jones. 2004. Analysis of interferon-gamma production by *Mycobacterium bovis* infected white-tailed deer (*Odocoileus virginianus*) using an in-vitro based assay. *Journal of Veterinary Diagnostic Investigation* 16: 16-21.

Palmer, M.V., D.L. Whipple, and **W.R. Waters**. 2001. Tuberculin skin testing in white-tailed deer (*Odocoileus virginianus*). *Journal of Veterinary Diagnostic Investigation* 13: 530-533.

Waters WR

- Sacco, R.E., D.L. Hutto, **W.R. Waters**, X. Li, M.E. Kehrli, F.A. Zuckermann, and M.J. Wannemuehler. 2000. Reduction in inflammation following blockade of CD18 or CD29 adhesive pathways during the acute phase of a spirochetal-induced colitis in mice. *Microbial Pathogenesis* 29: 289-299.
- Thacker, E.L., B.J. Thacker, M.P. Kuhn, A. Hawkins, and **W.R. Waters**. 2000. Evaluation of local and systemic immune responses induced by intramuscular injection of a *Mycoplasma hyopneumoniae* bacterin to pigs. *American Journal of Veterinary Research* 61: 1384-1389.
- Trott D.J., M.R. Moeller, R.L. Zuerner, J. Goff, **W.R. Waters**, D.P. Alt, R.L. Walker, and M.J. Wannemuehler. 2003. Isolation, characterization, and immune reactivity of *Treponema phagedenis*-like spirochetes from papillomatous digital dermatitis lesions in dairy cattle. *Journal of Clinical Microbiology* 41: 2522-2529.
- Waters, W.R.**, K.R. Harkins, and M.J. Wannemuehler. 2002. Five-color flow cytometric analysis of swine lymphocytes for detection of proliferation, apoptosis, viability, and phenotype. *Cytometry* 48: 146-152.
- Waters, W.R.**, R. Hontecillas, R.E. Sacco, F.A. Zuckermann, K.R. Harkins, J. Bassaganya-Riera, and M.J. Wannemuehler. 2000. Antigen-specific proliferation of porcine CD8 $\alpha\alpha$ cells to an extracellular bacterial pathogen. *Immunology* 101: 333-341.
- Waters W.R.**, J.M. Miller, M.V. Palmer, J.R. Stabel, D.E. Jones, K.A. Koistinen, E.M. Steadham, M.J. Hamilton, W.C. Davis, and J.P. Bannantine. 2003. Early induction of a humoral and cellular immune response during experimental *Mycobacterium avium* subsp. *paratuberculosis* infection of calves. *Infection and Immunity* 71: 5130-5138.
- Waters W.R.**, B.J. Nonnecke, M.R. Foote, A.C. Maue, T.E. Rahner, M.V. Palmer, D.L. Whipple, R.L. Horst, D.M. Estes. 2003. *Mycobacterium bovis* bacille Calmette Guerin vaccination of cattle: activation of bovine CD4 $^{+}$ and gamma-delta TCR $^{+}$ cells and modulation by 1,25-dihydroxyvitamin D3. *Tuberculosis* 83: 287-297.
- Waters, W.R.**, B.J. Nonnecke, M.V. Palmer, S. Robbe-Austermann, J.P. Bannantine, J.R. Stabel, D.L. Whipple, J.B. Payeur, D.M. Estes, J.E. Pitzer, F.C. Minion. 2004. Differentiation of *Mycobacterium bovis* infection of cattle from *M. avium* subsp. *avium* and *M. avium* subsp. *paratuberculosis* sensitization using a recombinant ESAT-6:CFP-10 fusion protein. *Clinical and Diagnostic Laboratory Immunology*, in press.
- Waters, W.R.**, B.J. Nonnecke, T.E. Rahner, M.V. Palmer, D.L. Whipple, and R.L. Horst. 2001. Modulation of *Mycobacterium bovis* specific responses of bovine peripheral blood mononuclear cells by 1,25-dihydroxyvitamin D3. *Clinical and Diagnostic Laboratory Immunology* 8: 1204-1212.
- Waters W.R.**, M.V. Palmer, B.J. Nonnecke, D.L. Whipple, and R.L. Horst. 2004. *Mycobacterium bovis* infection of vitamin D-deficient NOS2 $^{-/-}$ mice. *Microbial Pathogenesis* 36: 11-17.
- Waters W.R.**, M.V. Palmer, S.C. Olsen, R.E. Sacco, and D.L. Whipple. 2003. Immune responses of elk to *Mycobacterium bovis* bacille Calmette Guerin vaccination. *Vaccine* 21: 1518-1526.
- Waters, W.R.**, M.V. Palmer, B.A. Pesch, S.C. Olsen, M.J. Wannemuehler, and D.L.

Whipple. 2000. MHC class II-restricted, CD4+ T cell proliferative responses of peripheral blood mononuclear cells from *Mycobacterium bovis*-infected white-tailed deer. *Veterinary Immunology and Immunopathology* 76: 215-229.

Waters, W.R., M.V. Palmer, B.A. Pesch, S.C. Olsen, M.J. Wannemuehler, and D.L. Whipple. 2000. Lymphocyte subset proliferative responses of *Mycobacterium bovis*-infected cattle to purified protein derivative. *Veterinary Immunology and Immunopathology* 77: 257-273.

Waters, W.R., M.V. Palmer, R.E. Sacco, and D.L. Whipple. 2002. Nitric oxide production as an indication of *Mycobacterium bovis* infection of white-tailed deer (*Odocoileus virginianus*). *Journal of Wildlife Diseases* 38: 317-322.

Waters, W.R., M.V. Palmer, M.J. Wannemuehler, R.E. Sacco, and J.A. Harp. 2000. B cells are required for the induction of intestinal inflammatory lesions of Tcr+/-deficient mice persistently infected with *Cryptosporidium parvum*. *Journal of Parasitology* 86: 1073-1077.

Waters, W.R., M.V. Palmer, and D.L. Whipple. 2002. *Mycobacterium bovis*-infected white-tailed deer (*Odocoileus virginianus*): detection of immunoglobulin specific to crude mycobacterial antigens by ELISA. *Journal of Veterinary Diagnostic Investigation* 14: 470-475.

Waters W.R., M.V. Palmer, D.L. Whipple, M.C. Carlson, and B.J. Nonnecke. 2003. Diagnostic implications of antigen-induced IFN-gamma, nitric oxide, and TNF-alpha production by blood mononuclear cells from *Mycobacterium bovis*-infected cattle. *Clinical and Diagnostic Laboratory Immunology* 10: 960-966.

Waters W.R., M.V. Palmer, D.L. Whipple, R.E. Slaughter, and S.L. Jones. 2004. Immune responses of white-tailed deer (*Odocoileus virginianus*) to *Mycobacterium bovis* BCG vaccination. *Journal of Wildlife Diseases* 40: 66-78.

Waters, W.R., R.E. Sacco, Fach, S.J., M.V. Palmer, S.C. Olsen, and T.J. Kreeger. 2002. Analysis of mitogen-stimulated lymphocyte subset proliferation and nitric oxide production by peripheral blood mononuclear cells of captive elk (*Cervus elaphus*). *Journal of Wildlife Diseases* 38: 323-330.

Waters W.R., T.E. Rahner, M.V. Palmer, D. Cheng, B.J. Nonnecke, and D.L. Whipple. 2003. Expression of L-selectin (CD62L), CD44, and CD25 on activated bovine T cells. *Infection and Immunity* 71: 317-326.

Wesley RD

Mengeling, W.L., S.R. Bolin, K.M. Lager, A.C. Vorwald, and **R.D. Wesley**. 2000. Immunology and epidemiology of porcine reproductive and respiratory syndrome, pseudorabies, and postweaning multisystemic wasting syndrome. *Proceedings of 2nd National Meeting of the Italiana Veterinari per Animali da Reddito*, p. 86-92.

Mengeling, W.L., K.M. Lager, A.C. Vorwald, **R.D. Wesley**, and C. Prieto. 2000. Porcine reproductive and respiratory syndrome: vaccines and immunity. *Magazyn Weterynaryjny*, p. 16-18.

Wesley, R.D. 2000. Minimum viral dose for protective lactogenic immunity against transmissible gastroenteritis. *Proceedings of International Pig Veterinary Society 16th Congress*, p. 572-573.

Whipple DL

Palmer, M.V., W.R. Waters, and **D.L. Whipple**. 2002. Susceptibility of raccoons (*Procyon lotor*) to infection with *Mycobacterium bovis*. *Journal of Wildlife Diseases* 38(2): 266-274.

Palmer, M.V., **D.L. Whipple**, S.C. Olsen, and R.H. Jacobson. 2000. Cell mediated and humoral immune responses of white-tailed deer experimentally infected with *Mycobacterium bovis*. *Research in Veterinary Science* 68(1): 95-98.

Waters, W.R., M.V. Palmer, R.E. Sacco, and **D.L. Whipple**. 2002. Nitric oxide production as an indication of *Mycobacterium bovis* infection in white-tailed deer (*Odocoileus virginianus*). *Journal of Wildlife Diseases* 38(2): 338-343.

Waters W.R., T.E. Rahner, M.V. Palmer, D. Cheng, B.J. Nonnecke, and **D.L. Whipple**. 2003. Expression of L-Selectin (CD62L), CD44, and CD25 on activated bovine T cells. *Infection and Immunity* 71(1): 317-326.

Whipple, D.L., M.V. Palmer, R.E. Slaughter, and S.L. Jones. 2001. Comparison of purified protein derivatives and effect of skin testing on results of a commercial gamma interferon assay for diagnosis of tuberculosis in cattle. *Journal of Veterinary Diagnostic Investigation* 13(2): 117-122.

Yersin AG

Edens, F.W., **A.G. Yersin**, D.G. Simmons. 1999. Tryptophan methylester modulation of poult responses to *Bordetella avium*. *Poultry Science* 78(3): 327-335.

Yersin, A.G., F.W. Edens, and D.G. Simmons. 1998. The effects of *Bordetella avium* infection on elastin and collagen content of turkey trachea and aorta. *Poultry Science* 77(11): 1654-1660.

Zuerner RL

Brenot, A., D. Trott, I. Saint Girons, and **R. Zuerner**. 2001. Penicillin-binding proteins in *Leptospira interrogans*. *Antimicrobial Agents and Chemotherapy* 45: 870-877.

Hsu, T., D.L. Hutto, F.C. Minion, **R.L. Zuerner**, and M.J. Wannemuehler. 2001. Cloning of a beta-hemolysin gene of *Brachyspira (Serpulina) hyodysenteriae* and its expression in *Escherichia coli*. *Infection and Immunity* 69: 706-711.

Trott, D.J., **R.L. Zuerner**, M. Wannemuehler, and T.B. Stanton. 2000. Monoclonal antibodies reacting with *Brachyspira (Serpulina) pilosicoli* membrane proteins. 16th International Pig Veterinary Society, Abstract p. 41.

Zuerner, R., D. Haake, B. Adler, and R. Segers. 2000. Technological advances in the molecular biology of *Leptospira*. *Journal of Molecular Microbiology and Biotechnology* 2: 455-462.

New Orleans, Louisiana

Southern Regional Research Center

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=64-35-56-00>

Champagne ET

Chung, S.Y., C.L. Butts, and **E.T. Champagne**. 2001. Effect of curing on peanut allergenicity. American Peanut Research and Education Society, Abstract p. 21.

Chung, S.Y. and **E.T. Champagne**. 2001. Association of end-product adducts with increased IgE binding of roasted peanuts. Journal Agricultural Food Chemistry 49: 3592-3597.

Chung, S.Y. and **E.T. Champagne**. 2001. Effect of peanut maturity on IgE binding. Journal Allergy Clinical Immunology 107: 189.

Chung, S.Y. and **E.T. Champagne**. 2000. End products are potential cause for the increase in IgE-binding of roasted peanuts. American Research Education Society, Abstract p. 24.

Chung, S.Y., S.J. Maleki, **E.T. Champagne**, K.L. Buhr, and D.W. Gobet. 2000. Peanut allergenicity could be enhanced by biochemical reactions occurring during peanut roasting. Journal Allergy Clinical Immunology 105(1, part 2): 395.

Maleki, S.J. and **E.T. Champagne**. 2001. Allergenic and biophysical properties of purified peanut proteins before and after roasting. Journal Allergy Clinical Immunology 103(1, part 2): 624.

Maleki, S.J. and **E.T. Champagne**. 2001. Screening the core germplasm

collection for peanuts with reduced allergenic potential. American Peanut Research Education Society, Abstract p. 47.

Maleki, S.J. and **E.T. Champagne**. 2000. Comparing the peanut allergens before and after roasting. American Peanut Research Education Society, Abstract p. 55.

Maleki S.J., S.Y. Chung, **E.T. Champagne**, and J.P. Raufman. 2000. The effects of processing on the allergenic properties of peanut proteins. Journal Allergy Clinical Immunology 106: 763-768.

Chung S

Chung, S.Y., C.L. Butts, and E.T. Champagne. 2001. Effect of curing on peanut allergenicity. American Peanut Research and Education Society, Abstract p. 21.

Chung, S.Y. and E.T. Champagne. 2001. Association of end-product adducts with increased IgE binding of roasted peanuts. Journal Agricultural Food Chemistry 49: 3592-3597.

Chung, S.Y. and E.T. Champagne. 2001. Effect of peanut maturity on IgE binding. Journal Allergy Clinical Immunology 107: 189.

Chung, S.Y. and E.T. Champagne. 2000. End products are potential cause for the increase in IgE-binding of roasted peanuts. American Research Education Society, Abstract p. 24.

Chung, S.Y., S.J. Maleki, E.T. Champagne, K.L. Buhr, and D.W. Gobet. 2000. Peanut allergenicity could be enhanced by

biochemical reactions occurring during peanut roasting. 2000. *Journal Allergy Clinical Immunology* 105(1, part 2): 395.

Maleki S.J., **S.Y Chung**, E.T. Champagne, and J.P. Raufman. 2000. The effects of processing on the allergenic properties of peanut proteins. *Journal Allergy Clinical Immunology* 106: 763-768.

Maleki SJ

Airee, R., A. Coke, S.K. Airee, and **S.J. Maleki**. 2000. Correlation between aflatoxins, allergens and ingredients of peanuts? American Chemical Society National Meeting, Abstract p.848.

Bannon, G.A., R.A. Kopper, **S.J. Maleki**, D.S. Shin, H.A. Sampson, and A.W. Burks. 2000. Structure plays a critical role in the allergenicity of food proteins. *Journal Allergy Clinical Immunology* 105(1, part 2): 554.

Chung, S.Y., **S.J. Maleki**, E.T. Champagne, K.L. Buhr, and D.W. Gobet. 2000. Peanut allergenicity could be enhanced by biochemical reactions occurring during peanut roasting. *Journal Allergy Clinical Immunology* 105(1, part 2): 395.

Maleki, S.J. and E.T. Champagne. 2001. Allergenic and biophysical properties of purified peanut proteins before and after roasting. *Journal Allergy Clinical Immunology* 103(1, part 2): 624.

Maleki, S.J. and E.T. Champagne. 2001. Screening the core germplasm collection for peanuts with reduced allergenic potential. American Peanut Research Education Society, Abstract p. 47.

Maleki, S.J. and E.T. Champagne. 2000. Comparing the peanut allergens before and after roasting. American Peanut Research Education Society, Abstract p. 55.

Maleki S.J., S.Y. Chung, E.T. Champagne, and J.P. Raufman. 2000. The effects of processing on the allergenic properties of peanut proteins. *Journal Allergy Clinical Immunology* 106: 763-768.

Maleki S.J., R.A. Kopper, D.S. Shin, S.J. Stanley, H.A. Sampson, A.W. Burks, and G.A. Bannon. 2000. Structure of the Major Peanut Allergen Ara h 1 May protect IgE-Binding Epitopes from degradation. *Journal Immunology* 164: 5844-5849.

Sen, M., R.A. Kopper, **S.J. Maleki**, G.A. Bannon, H.A. Sampson, and A.W. Burks. 2001. Structure plays a critical role in the allergenicity of food proteins. *Journal Allergy Clinical Immunology* 103(1, part 2): 612.

Beltsville, Maryland

Henry A. Wallace Beltsville Agricultural Research Center

<http://www.ba.ars.usda.gov/>

Allen PC

Allen, P.C. 2001. Oxidative stress during avian coccidiosis. 8th Coccidiosis Conference, p. 156.

Allen, P.C. and R.H. Fetterer. 2001. Effects of dietary Vitamin E supplementation on *Eimeria maxima* infections in chickens. 8th Coccidiosis Conference, p. 156.

Zhu, J., H.S. Lillehoj, **P. Allen**, H.H. Cheng, M. Emara, M. Sadjadi, and D. Pollock. 2001. Mapping quantitative trait loci affecting disease resistance to avian coccidiosis. Proceedings of the 73rd Northeastern Conference on Avian Diseases, June 27-29. Virginia-Maryland College of Veterinary Medicine, University of Maryland, College Park, p34.

Zhu, J., H.S. Lillehoj, C. van Tassel, M. Emara, **P. Allen**, H. Cheng, D. Pollock, M. Sadjadi, and T. Sonstegard. 2001. Mapping QTL loci affecting growth and disease resistance to avian coccidiosis. Proceedings of the International Animal Agricultural and Food Sciences Conference, July 24-28, Indianapolis, Indiana p. 43.

Augustine PC

Augustine, P.C. 2001. *Eimeria adenoeides* sporozoites: enhanced invasion of cultured cells previously inoculated with *Eimeria acervulina* sporozoites. Parasitology Research 87(7): 548-52.

Augustine, P.C. 2001. Cell: sporozoite interactions and invasion by apicomplexan parasites of the genus *Eimeria*. International Journal for Parasitology 31(1): 1-8.

Augustine, P.C. 2001. Invasion in different cell types by sporozoites of *Eimeria* species and effects of monoclonal antibody 1209-C2 on invasion of cells by sporozoites of several apicomplexan parasites. Journal of Eukaryotic Microbiology 48: 177-181.

Augustine, P., G. Olsen, H. Danforth, G. Gee, and M. Novilla. 2001. Use of monoclonal antibodies developed against chicken coccidia (*Eimeria*) to study invasion and development of *Eimeria reichenowi* in Florida sandhill cranes (*Grus canadensis*). Journal of Zoo and Wildlife Medicine 32(1): 65-70.

Bannerman DD

Bannerman, D.D., K.T. Eiting, R.K. Winn, and J.M. Harlan. 2004. FLICE-like inhibitory protein (FLIP) protects against apoptosis and suppresses NF- κ B activation induced by bacterial lipopolysaccharide. American Journal of Pathology 165: 1423-1431.

Bannerman, D.D. and S.E. Goldblum. 2003. Mechanisms of bacterial lipopolysaccharide-induced endothelial apoptosis. American Journal of Physiology - Lung Cellular and Molecular Physiology 284(6): L899-L914.

Bannerman, D.D., M.J. Paape, J.P. Goff, K. Kimura, J.D. Lippolis, and J.C. Hope. 2004. Innate immune response to intramammary infection with *Serratia marcescens* and *Streptococcus uberis*. Veterinary Research 35(6): 681-700.

Bannerman, D.D., M.J. Paape, W.R. Hare, and J.C. Hope. 2004. Characterization of the bovine innate immune response to intramammary infection with *Klebsiella pneumoniae*. Journal of Dairy Science 87(8): 2420-2432.

Bannerman, D.D., M.J. Paape, W.R. Hare, and E.J. Sohn. 2003. Increased levels of LPS-binding protein in bovine blood and milk following bacterial lipopolysaccharide challenge. Journal of Dairy Science 86: 3128-3137.

Bannerman, D.D., M.J. Paape, J.W. Lee, X. Zhao, J.C. Hope, P. Rainard. 2004. *Escherichia coli* and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection. Clinical and Diagnostic Laboratory Immunology 11: 463-472.

Erwert, R.D., K.T. Eiting, J.C. Tupper, R.K. Winn, J.M. Harlan, and **D.D. Bannerman**. 2003. Shiga toxin induces decreased expression of the anti-apoptotic protein Mcl-1 concomitant with the onset of endothelial apoptosis. Microbial Pathogenesis 35(2): 87-93.

Li, X., J.C. Tupper, **D.D. Bannerman**, R.K. Winn, C.J. Rhodes, J.M. Harlan. 2003. Phosphoinositide 3 kinase mediates Toll-like receptor 4-induced activation of NF- κ B in endothelial cells. Infection and Immunity 71: 4414-4420.

Paape, M.J., **D.D. Bannerman**, X. Zhao, and J.W. Lee. 2003. The bovine neutrophil: structure and function in blood and milk. Veterinary Research 34: 597-627.

Sakarya, S., S. Rifat, J. Zhou, **D.D. Bannerman**, N.M. Stamatos, A.S. Cross, and S.E. Goldblum. 2004. Mobilization of neutrophil sialidase

activity desialylates the pulmonary vascular endothelial surface and increases resting neutrophil adhesion to and migration across the endothelium. Glycobiology 14: 481-494.

Sohn, E.J., M.J. Paape, R.P. Peters, R.H. Fetterer, N.C. Talbot, and **D.D. Bannerman**. 2004. The production and characterization of anti-bovine CD14 monoclonal antibodies. Veterinary Research 35 (5): 597-608.

Dawson H

Almeria, S., T. De-Marez, **H. Dawson**, R. Araujo, J.P. Dubey, and L.C. Gasbarre. 2003. Cytokine gene expression in dams and fetuses after experimental *Neospora caninum* infection of heifers at 110 days of gestation. Parasite Immunology 25(7): 383-392.

Dubey JP

Dubey, J.P., M.M. Garner, M.D. Stetter, A.E. Marsh, and B.C. Barr. 2001. Acute *Sarcocystis falcatula*-like infection in a carmine bee-eater (*Merops nubicus*) and immunohistochemical cross reactivity between *S. falcatula* and *Sarcocystis neurona*. Journal of Parasitology 87: 824-831.

Kringel, H., **J.P. Dubey**, E. Beshah, R. Hecker, and J.F. Urban. 2001. Effect of CpG oligodeoxynucleotide (ODN)-induced immune activation of porcine immunity to *Toxoplasma gondii*. American Association of Veterinary Parasitology 46: 15.

Lindsay, D.S. and **J.P. Dubey**. 2001. Direct agglutination test for the detection of antibodies to *Sarcocystis neurona* in experimentally infected animals. Veterinary Parasitology 95: 179-186.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, D. Zarlenga, L. Jauregui, M. Cosio, L.W. Douglass, **J.P. Dubey**, and J.K. Lunney. 2001. Cytokine and lymphocyte

profile after oral infection with *Toxoplasma gondii* oocysts. Scandinavian Journal of Immunology 54: 93.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, D. Zarlenga, L. Jauregui, M. Cosio, L.W. Douglass, **J.P. Dubey**, and J.K. Lunney. 2001. Cytokine and lymphocyte profile in miniature swine after oral infection with *Toxoplasma gondii* oocysts. International Journal of Parasitology 31: 187-195.

Elsasser TH

Elsasser, T.H., K.C. Klasing, N. Filipov, F. Thompson, G.P. Moberg (ed.), and J.A. Mench. 2000. The metabolic consequences of stress: targets for stress and priorities of nutrient use. In G. Moberg, editor, The biology of animal stress: basic principles and implications for animal welfare, CABI Publishing: Wallingford, UK, p. 77-110.

Kahl, S., **T.H. Elsasser**, J.L. Sartin, and R. Fayer. 2002. Effect of progressive cachectic parasitism and growth hormone treatment on hepatic 5'-deiodinase activity in calves. Domestic Animal Endocrinology 22(4): 211-221.

Perkins, K.H., M.J. VandeHaar, J.L. Burton, J.S. Liesman, R.J. Erskine, and **T.H. Elsasser**. 2002. Clinical responses to intramammary endotoxin infusion in dairy cows subjected to feed restriction. Journal of Dairy Science 85(7): 1724-1731.

Pio, R., A. Martinez, **T.H. Elsasser**, and F. Cuttitta. 2000. Presence of immunoreactive adrenomedullin in human and bovine milk. Peptides 21(12): 1859-1863.

Richards, M.P., T.J. Caperna, **T.H. Elsasser**, C.M. Ashwell, and J.P.

McMurtry. 2000. Design and application of a polyclonal peptide antiserum for the universal detection of leptin protein. Journal of Biochemical and Biophysical Methods 45: 147-156.

Evans JD

Evans, J.D. 2000. Microsatellite loci in the honey bee parasitic mite *Varroa jacobsoni*. Molecular Ecology 9: 1436-1438.

Fayer R

Higgins, J., M.C. Jenkins, **R. Fayer**, L. Xiao, A.A. Lal, B. Bennet, and J. Richards. 2000. New methods for molecular diagnostics for emerging waterborne pathogens. Proceedings of the International Conference on Emerging and Infectious Diseases, p. 38.

Jenkins, M.C., J. Trout, J. Higgins, and **R. Fayer**. 2000. The mRNA levels for amyloglucosidase as measured by RT-PCR correlate with infectivity and fecundity of *Cryptosporidium parvum* oocysts for neonatal BALB/c mice. Proceedings of the International Conference on Emerging and Infectious Diseases, p. 9.

Jenkins, M.C., J. Trout, C. Murphy, J. Harp, J. Higgins, W. Wergin, and **R. Fayer**. 1999. Cloning and expression of a DNA sequence encoding a 41-kilodalton *Cryptosporidium parvum* oocyst wall protein. Clinical and Diagnostic Laboratory Immunology 6(6): 912-920.

Fetterer RH

Allen, P.C. and **R.H. Fetterer**. 2001. Effects of dietary Vitamin E supplementation on *Eimeria maxima* infections in chickens. 8th Coccidiosis Conference, p. 156.

Allen, P.C. and **R.H. Fetterer**. 2000. Effect of *Eimeria acervulina* infections on plasma l-arginine. Poultry Science 79: 1414-1417.

Fetterer, R.H. and P.C. Allen. 2000. *Eimeria acervulina* elevates plasma and muscle 3-methylhistidine levels in chickens. *Journal of Parasitology* 86: 783-781.

Gasbarre LC

Gasbarre, L.C., E.A. Leighton, and T. Sonstegard. 2001. Role of the bovine immune system and genome in resistance to gastrointestinal nematodes. *Veterinary Parasitology* 98: 51-64.

Gasbarre, L.C., W.L. Stout, L. Lohr, and J. Guerrero. 2001. Gastrointestinal nematodes on a grazing dairy operation: Effect of parasite control on production and parasite transmission. *Proceedings of the American Association of Veterinary Parasitology*, p. 51.

Hediger-Weithaler, B.M., V.T. Heussler, R. Eicher, A. Zakher, D.S. Zarlenga, **L.C. Gasbarre**, and A.S. Waldvogel. 2000. Interferon- and interleukin 4 mRNA expression by PBMCs from BVDV seropositive pregnant and non-pregnant cattle. *Veterinary Immunology and Immunopathology* 77: 201-212.

Sonstegard, T.S., **L.C. Gasbarre**, C.P. van Tassell, and T. Padilha. 2001. Genomic tools to improve parasite resistance. *Proceedings of the American Association of Veterinary Parasitologists* 46: 46.

Zarlenga, D.S., M.B. Chute, **L.C. Gasbarre**, and P. Boyd. 2001. A multiplex PCR assay for differentiating economically important gastrointestinal nematodes of cattle. *Veterinary Parasitology* 97: 199-209.

Jenkins MC

Baker, S.A., **M.C. Jenkins**, and C.L. Chappell. 2001. Validating a

recombinant protein antigen in a serodiagnostic test for antibodies to *Cryptosporidium parvum*. *Proceedings of the 51st Annual Meeting of the American Society of Parasitologists*, Albuquerque, New Mexico, p. 50.

Dyer, R.M., **M.C. Jenkins**, O.C.H. Kwok, L.W. Douglas, and J.P. Dubey. 2000. Serologic survey of *Neospora caninum* infection in a closed dairy cattle herd in Maryland: risk of serologic reactivity by production groups. *Veterinary Parasitology* 90: 171-181.

Fayer, R., E.J. Lewis, J.M. Trout, T.K. Graczyk, **M.C. Jenkins**, J. Higgins, L. Xiao, and A.A. Lal. 1999. *Cryptosporidium parvum* in oysters from commercial harvesting sites in the Chesapeake Bay. *Emerging Infectious Diseases* 5: 706-710.

Higgins, J., **M.C. Jenkins**, R. Fayer, L. Xiao, A.A. Lal, B. Bennet, and J. Richards. 2000. New methods for molecular diagnostics for emerging waterborne pathogens. *Proceedings of the International Conference on Emerging and Infectious Diseases*, p. 38.

Jenkins, M.C., S. Botero, P. Allen, and H. Danforth. 2000. Application of non-attenuated and gamma-irradiated eimerian oocysts to control avian coccidiosis. *Proceedings of the Annual Meeting of the American Association of Veterinary Parasitologists*, p. 85.

Jenkins, M.C., J.A. Caver, C. Bjorkman, T.C. Anderson, S. Romand, B. Vinyard, A. Ugglia, P. Thulliez, and J.P. Dubey. 2000. Serological investigation of an outbreak of *Neospora caninum*-associated abortion in a dairy herd in the southeastern United States. *Veterinary Parasitology* 94(1-2): 17- 26.

Jenkins, M.C., J. Trout, M.S. Abrahamsen, C.A. Lancto, J. Higgins, and R. Fayer. 2000. Estimating viability of *Cryptosporidium parvum* oocysts using reverse transcriptase-

polymerase chain reaction (RT-PCR) directed at mRNA encoding amyloglucosidase. *Journal of Microbial Methods* 43: 97-106.

Jenkins, M.C., J. Trout, and R. Fayer. 2001. Treatment of *Cryptosporidium parvum* sporozoites with antibodies to a recombinant CP47 surface antigen inhibits in vitro parasite development. Proceedings of the Annual Meeting of the American Association of Veterinary Parasitologists, Boston, Massachusetts, p. 49.

Jenkins, M.C., J. Trout, J. Higgins, and R. Fayer. 2000. The mRNA levels for amyloglucosidase as measured by RT-PCR correlate with infectivity and fecundity of *Cryptosporidium parvum* oocysts for neonatal BALB/c mice. Proceedings of the International Conference on Emerging and Infectious Diseases Late Breakers, p. 9.

Jenkins, M.C., J. Trout, C. Murphy, J. Harp, J. Higgins, W. Wergin, and R. Fayer. 1999. Cloning and expression of a DNA sequence encoding a 41-kilodalton *Cryptosporidium parvum* oocyst wall protein. *Clinical and Diagnostic Laboratory Immunology* 6(6): 912-920.

Lillehoj, H.S., K.D. Choi, **M.C.**

Jenkins, V.N. Vakharia, K.D. Song, J.Y. Han, and E.P. Lillehoj. 2000. A recombinant *Eimeria* protein inducing interferon- production. Comparison of different gene expression systems and immunization strategies for vaccination against coccidiosis. *Avian Diseases* 44: 379-389.

Lillehoj, H.S., **M.C. Jenkins,** and V. Vakharia. 2000. *Eimeria* protein inducing interferon- production. Comparison of different gene expression systems and immunization strategies for

vaccination against coccidiosis. World Poultry Congress, p. 17-13.

Lillehoj HS

Babu, U., R.A. Dalloul, M. Okamura, **H.S. Lillehoj,** H. Xie, R. Raybourne, D. Gaines, and R.A. Heckert. 2004. *Salmonella enteritidis* clearance and immune responses in chickens following *Salmonella* vaccination and challenge. *Veterinary Immunology and Immunopathology* 101: 251-257.

Babu, U., M. Scott, M.J. Myers, M. Okamura, D. Gaines, R. Raybourne, H.F. Yancy, **H.S. Lillehoj,** and R. Heckert. 2002. Effects of live attenuated and killed *Salmonella* vaccine on T-lymphocyte mediated immunity in laying hens. *Veterinary Immunology and Immunopathology* 6729: 1-6.

Choi, K.D. and **H.S. Lillehoj.** 2000. Effect of chicken IL-2 on T-cell growth and function. Increased T cell response following DNA immunization. *Veterinary Immunology and Immunopathology* 73: 309-312.

Constantinoiu, C.C., **H.S. Lillehoj,** M. Matsubayashi, Y. Hosoda, H. Tani, H. Matsuda, L.K. Sasai, and E. Baba. 2003. Analysis of cross-reactivity of five new chicken monoclonal antibodies which recognize the apical complex of *Eimeria* using confocal laser immunofluorescence assay *Veterinary Parasitology* 118: 29-35.

Constantinoiu, C.C., **H.S. Lillehoj,** M. Matsubayashi, H. Tani, H. Matsuda, L.K. Sasai, and E. Baba. 2004. Characterization of stage-specific and cross-reactive antigens from *Eimeria acervulina* by chicken monoclonal antibodies. *Journal of Veterinary Medical Science* 66: 403-408.

Dalloul, R., **H.S. Lillehoj,** X.A. Ding, D.M.C. Klinman, and R.A. Heckert. 2004. In ovo administration of CpG oligodeoxynucleotides and the recombinant

microneme protein MIC2 protects against *Eimeria* infections. In press.

Dalloul, R.A., **H.S. Lillehoj**, and J.A. Doerr. 2001. Probiotics in broilers: Modulation of intestinal immune function. Proceedings of the 48th Maryland Nutrition Conference for Feed Manufacturers, p. b30-41.

Dalloul, R.A., **H.S. Lillehoj**, M. Okamura, H. Xie, W. Min, X. Ding, and R.A. Heckert. 2004. In vivo effects of CpG oligodeoxynucleotide on *Eimeria* infection in chickens. *Avian Diseases* 48(4): 783-790.

Dalloul, R.A., **H.S. Lillehoj**, T.A. Shellem, and J.A. Doerr. 2003. Enhanced mucosal immunity against *Eimeria acervulina* in broilers fed a *Lactobacillus*-based probiotic. *Poultry Science* 82: 62-66.

Dalloul, R.A., **H.S. Lillehoj**, T.A. Shellem, and J.R. Doerr. 2003. Intestinal immunomodulation by Vitamin A deficiency and *Lactobacillus*-based probiotic in *Eimeria acervulina*-infected broiler chickens. *Avian Diseases* 47: 1313-1320.

Dalloul, R.A., **H.S. Lillehoj**, T.A. Shellem, and J.A. Doerr. 2002. Effect of Vitamin A deficiency on host intestinal immune response to *Eimeria acervulina* in broiler chickens. *Poultry Science* 81: 1509-1515.

Ding, X., **H.S. Lillehoj**, M.A. Quiroz, E. Berensee, and E.P. Lillehoj. 2004. 3 Protective immunity against *Eimeria acervulina* following in ovo immunization with a recombinant subunit vaccine and cytokine genes. *Infection and Immunity* 72(12): 6939-6944.

Emara, M.G., H. Kim, J. Zhu, R.R. Lapierre, N. Lakshmanan, and **H.S. Lillehoj**. 2002. Genetic diversity at the major histocompatibility complex (B) and microsatellite loci in three commercial broiler pure lines. *Poultry Science* 81(11): 1609-1617.

Emara, M.G., R.R. Lapierre, G.M. Greene, M. Knieriem, J.K. Rosenberger, D.L. Pollock, M. Sadjadi, C.D. Kim, and **H.S. Lillehoj**. 2002. Phenotypic variation among three broiler pure lines for Marek's Disease, coccidiosis and antibody response to sheep red blood cells. *Poultry Science* 81:642-648.

Gehad, A.E., **H.S. Lillehoj**, G.L. Hendrick III, and M. Mashaly. 2002. I. The role of cytokines and hormones in the initiation of humoral immunity using T-independent and T-dependent antigens. *Developmental & Comparative Immunology* 26(8): 751-759.

Gehad, A.E., **H.S. Lillehoj**, G.L. Hendrick, and M. Mashaly. 2002. II. The effects of T-independent and T-dependent antigens on the distribution of lymphocyte populations. *Developmental & Comparative Immunology* 26(8): 761-771.

Gehad A.E., **H.S. Lillehoj**, and M. Mashaly. 2001. The role of lymphocytes and hormones during the development of the delayed hypersensitivity response in immature male chickens. In K.A. Schat, editor, *Current Progress on Avian Immunology Research*, American Association of Avian Pathologists: Kennett Square, Pennsylvania, p. 308-314.

Guangxing, L., E. Lillehoj, and **H.S. Lillehoj**. 2001. Interleukin-2 production in SC and TK chickens infected with *Eimeria tenella*. *Avian Diseases* 46: 2-9.

Guangxing L., **H.S. Lillehoj**, and W. Min. 2001. Production and characterization of monoclonal antibodies detecting the chicken interleukin-15 receptor alpha chain.

Veterinary Immunology and Immunopathology 82: 215-227.

Hwang, K.C., K.D. Song, T.H. Kim, H.K. Lee, S.H. Sohn, **H.S. Lillehoj**, and J.Y. Han. 2001. Genetic linkage mapping of RAPD markers segregating in Korean Ogo chickens-White Leghorn backcross population. Asian-Australian Journal of Animal Science 14: 302-306.

Jenkins, M., P. Allen, R. Fetterer, **H.S. Lillehoj**, K. Miska. 2003. Are there immunovariants of coccidia that can affect vaccine efficacy? Watt Poultry USA.

Kim, J.K., **H.S. Lillehoj**, W.G. Min, J.Y. Han, K. Sasai, E.J. Sohn, and E.P. Lillehoj. 2001. Recombinant single chain fragment variable region antibody constructed from chicken monoclonal antibody detects apical complex protein of *Eimeria acervulina*. In K.A. Schat, editor, Current Progress on Avian Immunology Research, American Association of Avian Pathologists: Kennett Square, Pennsylvania, p. 64-70.

Kim, J.K., W. Min, **H.S. Lillehoj**, S.W. Kim, E.J. Sohn, K.D. Song, and J.Y. Han. 2001. Generation and characterization of recombinant scFv antibodies detecting *Eimeria acervulina* surface antigens. Hybridoma 20: 175-181.

Kliger, C.A., A.E. Gehad, R.M. Hulet, W.B. Roush, **H.S. Lillehoj**, and M.M. Mashaly. 2000. Effect of melatonin and photoperiod on broiler immune system. Poultry Science 79: 18-25.

Lillehoj, E.P., C.H. Yun, and **H.S. Lillehoj**. 2000. Vaccine for avian enteropathogens *Eimeria*, *Cryptosporidium* and *Salmonella*. Animal Health Research Reviews 1: 47-65.

Lillehoj, H.S. 2002. Beyond genomics revolution: Emergence of agricultural biotechnology. Proceeding of Future Perspectives in Animal Biotechnology, Seoul National University, College of Animal Science and Technology, Suwon, Korea, p. 5-21.

Lillehoj, H.S. 2002. Coccidiosis: This poultry disease's impact is anything but paltry. Healthy Animals <http://www.ars.usda.gov/is/np/ha/ha10.htm>

Lillehoj, H.S. 2002. Gut immune responses and intervention strategies against *Eimeria* and *Salmonella*. Proceedings of International Seminars on Avian Pathology and Production, p. 91-131.

Lillehoj, H.S. 2001. Application of DNA marker technology to the development of coccidiosis control strategy. Poultry Biotechnology and its Applications, Proceedings of Korean Society of Poultry Science Symposium, p. 1-17.

Lillehoj, H.S. 2000. Mucosal immune response to coccidiosis. Proceedings of the 21st World Poultry Congress, S1.10. (CD).

Lillehoj, H.S. 2000. Review on vaccine development against enteric parasites *Eimeria* and *Cryptosporidium*. Japanese Poultry Science 37: 117-141.

Lillehoj, H.S. 2000. Subcutaneous injection with IL-2 gene enhances cell-mediated immunity. Proceedings of the 21st World Poultry Congress, P17.14. (CD).

Lillehoj, H.S., K.D. Choi, M.C. Jenkins, V.N. Vakharia, K.D. Song, J.Y. Han, and E.P. Lillehoj. 2000. A recombinant *Eimeria* protein inducing interferon- γ production. Comparison of different gene expression systems and immunization strategies for vaccination against coccidiosis. Avian Diseases 44: 379-389.

- Lillehoj, H.S.,** R. Dalloul, and W. Min. 2003. Enhancement of intestinal immunity to coccidiosis using cytokines, oligodinucleotides and probiotics. 2003. World Poultry: Special Edition, p. 18-23.
- Lillehoj, H.S.** and L. Guangxing. 2004. Nitric oxide production by macrophages stimulated with sporozoites, lipopolysaccharide, or IFN- γ and its dynamic changes in SC and TK strains of chickens infected with *Eimeria tenella*. Avian Diseases 48: 244-253.
- Lillehoj, H.S.,** M.C. Jenkins, and V.N. Vakharia. 2000. *Eimeria* protein inducing interferon- γ production. Comparison of different gene expression systems and immunization strategies for vaccination against coccidiosis. Proceedings of the 21st World Poultry Congress, P17.13. (CD).
- Lillehoj, H.S.** and E.P. Lillehoj. 2000. Avian coccidiosis. A review of acquired intestinal immunity and vaccination strategies. Avian Diseases 44: 408-425.
- Lillehoj, H.S.** and W.G. Min. 2002. *Gallus gallus* mRNA for interleukin-16. Genbank. Accession Number. AJ508678.
- Lillehoj, H.S.** and W.G. Min 2002. *Gallus gallus* mRNA for putative common cytokine receptor gamma chain β . Genbank. Accession Number. AJ419896.
- Lillehoj, H.S.,** W.G. Min 2002. *Gallus gallus* mRNA for putative common cytokine receptor gamma chain α . Genbank. Accession Number. AJ419898.
- Lillehoj, H.S.,** W.G. Min, K.D. Choi, U. Babu, J. Burnside, T. Miyamoto, and E.P. Lillehoj. 2001. Functional characterization of chicken cytokines homologous to mammalian IL-15 and IL-2. In K.A. Schat, editor, Current Progress on Avian Immunology Research, American Association of Avian Pathologists: Kennett Square, Pennsylvania, p. 8-13.
- Lillehoj, H.S.,** W. Min, K.D. Choi, U.S. Babu, J. Burnside, T. Miyamoto, B.M. Rosenthal, and E.P. Lillehoj. 2001. Molecular, cellular, and functional characterization of chicken cytokines homologous to mammalian IL-15 and IL-2. Veterinary Immunology and Immunopathology 82: 229-244.
- Lillehoj, H.S.,** W.G. Min, and R.A. Dalloul. 2004. Recent progress on the cytokine regulation of intestinal immune responses to *Eimeria*. Poultry Science 83:611-623.
- Lillehoj, H.S.** and M. Okamura. 2003. Host immunity and vaccine Development to coccidia and *Salmonella* infections in chickens. Journal of Poultry Science 40(3): 151-193.
http://www.jstage.jst.go.jp/article/jpsa/40/3/151/_pdf
- Lillehoj, H.S.,** C.H. Yun, and M. Nichols. 2000. Monoclonal antibodies detecting chicken IFN-gamma and development of in vitro assay. Patent Serial Number 09/534,002, Document Number 0165.99.
- Lillehoj, H.S.,** J. Zhu, and W.G. Min. 2001. Application of DNA marker technology and functional genomics to the development of coccidiosis control strategy. Korean Journal of Poultry Science 28: 107-123.
- Min, W., J.K. Kim, **H.S. Lillehoj,** E.J. Sohn, J.Y. Han, K.D. Song, and E.P. Lillehoj. 2001. Synthetic construct for anti-*Eimeria acervulina* apical protein antibody, clone 6D12. Genbank. Accession Number. AJ3000836.
- Min, W., J.K. Kim, **H.S. Lillehoj,** E.J. Sohn, J.Y. Han, K.D. Song, and E. P. Lillehoj. 2001. Characterization of recombinant scFv

antibody reactive with an apical antigen of *Eimeria acervulina*. Biotechnology Letters 23: 949-955.

Min, W. and **H.S. Lillehoj**. 2004. Identification and characterization of chicken interleukin-16 cDNA. Developmental and Comparative Immunology 28: 153-162.

Min, W. and **H.S. Lillehoj**. 2002. Sequence and functional properties of chicken interleukin-17. Journal of Interferon and Cytokine Research 22: 1125-1130.

Min, W. and **H.S. Lillehoj**. 2000. Synthetic construct for anti-*Eimeria acervulina* surface ScFv antibody, clone 5D11. Genbank, Accession Number. AJ298108.

Min, W. and **H.S. Lillehoj**. 2000. Synthetic construct for anti-*Eimeria acervulina* surface ScFv antibody, clone 2-1. Genbank, Accession Number. AJ298107.

Min, W.G., **H.S. Lillehoj**, J. Burnside, H.B. Kim, and E. Lillehoj. 2001. Northern blot analysis of concanavalin A-induced expression of chicken gene homologous to the mammalian IL-2 receptor chain. In K.A. Schat, editor, Current Progress on Avian Immunology Research, American Association of Avian Pathologists: Kennett Square, Pennsylvania, p. 346-351.

Min, W., **H.S. Lillehoj**, J. Burnside, K.C. Weining, P. Staeheli, and M. Nichols. 2002. Adjuvant effects of IL-1beta, IL-8, IL-15, IFN-alpha, IFN-gamma, TGF-beta and Lymphotoxin on DNA vaccination against *Eimeria acervulina*. Vaccine 20: 267-274.

Min, W.G., **H.S. Lillehoj**, J. Burnside, K.C. Weining, P. Staeheli, and M.B.

Nichols. 2001. Effect of IL-1, IL-8, IL-15, IL-2, IFN- α , IFN- γ , lymphotoxin and TGF- β 4 on DNA vaccination against *Eimeria acervulina*. In K.A. Schat, editor, Current Progress on Avian Immunology Research, American Association of Avian Pathologists: Kennett Square, Pennsylvania, p. 352-356.

Min, W., **H.S. Lillehoj**, and R. Fetterer. 2002. Identification of an alternatively spliced transcript isoform of the common cytokine receptor γ chain. Biochemical and Biophysical Research Communication 299: 321-327.

Min, W., **H.S. Lillehoj**, L. Guangxing, E.J. Sohn, and T. Miyamoto. 2002. Development and characterization of monoclonal antibodies to chicken interleukin-15. Veterinary Immunology and Immunopathology 88: 49-56.

Min, W., **H.S. Lillehoj**, S. Kim, J.J. Zhu, H. Beard, N. Alkharouf, and B.F. Matthews. 2003. Profiling local gene expression changes associated with *Eimeria maxima* and *Eimeria acervulina* using cDNA microarray. Applied Microbiology and Biotechnology 62: 392-399.

Miska, K., R. Fetterer, W. Min, **H.S. Lillehoj**. Identification of a heat shock protein 90 gene in two *Eimeria* species: Expression and evolutionary analysis. Molecular and Biochemical Parasitology. In press.

Miyamoto, T., **H.S. Lillehoj**, E.J. Sohn, and W. Min. 2001. Production and characterization of monoclonal antibodies detecting chicken interleukin-2 and the development of an antigen capture enzyme-linked immunosorbent assay. Veterinary Immunology and Immunopathology 80: 245-257.

Miyamoto, T., W. Min, and **H.S. Lillehoj**. 2001. Kinetics of interleukin-2 production in chickens infected with *Eimeria tenella*.

Comparative Immunology,
Microbiology and Infectious Diseases
25: 149-158.

Miyamoto, T., W. Min, and **H.S. Lillehoj**. 2001. Lymphocyte proliferation response during *Eimeria tenella* infection assessed by a new, reliable, non-radioactive colorimetric assay. *Avian Diseases* 46: 10-16.

Okamura, M., **H.S. Lillehoj**, R.B. Raybourne, U.S. Babu, R.A. Heckert. 2004. Cell-mediated immune responses to a killed *Salmonella enteritidis* vaccine. *Comparative Immunology, Microbiology and Infectious Diseases* 27(4): 255-272.

Okamura, M., **H.S. Lillehoj**, R.B. Raybourne, U. Babu, and R. Heckert. 2003. Antigen-specific lymphocyte proliferation, Interleukin-6 (IL-6) and IL-2 production in chickens immunized with outer membrane protein of *Salmonella enteritidis*. *Avian Diseases* 47: 1331-1338.

Olah, I., K.H. Gumati, N. Nagy, A. Magyar, B. Kaspers, and **H.S. Lillehoj**. 2002. Diverse expression of the K-1 antigen by cortico-medullary and reticular epithelial cells of the bursa of Fabricius in chicken and guinea fowl. *Developmental and Comparative Immunology* 26: 481-488.

Pinard-van der Laan, M.H., **H.S. Lillehoj**, and J. Zhu. 2003. Disease resistance and transmission. Genetic resistance and transmission of avian parasites. In W.M. Muir and S.E. Aggrey, editors, *Poultry Genetics, Breeding and Biotechnology*, CABI Publishing.

Sasai, K., M. Aita, **H.S. Lillehoj**, T. Miyamoto, T. Fukata, and E. Baba. 2000. Dynamics of lymphocyte

subpopulation changes in the cecal tonsils of chickens infected with *Salmonella enteritidis*. *Veterinary Microbiology* 74: 345-351.

Song, K.D., J.Y. Han, W. Min, **H.S. Lillehoj**, S. Kim, and J. Kim. 2001. Molecular cloning and characterization of cDNA encoding immunoglobulin heavy and light chain variable regions from four chicken monoclonal antibodies specific to surface antigens of intestinal parasites, *Eimeria acervulina*. *Korean Journal of Microbiology* 39(1): 49-55.

Song, K.D., **H.S. Lillehoj**, K.D. Choi, M.S. Parcels, J.T. Huynh, C.H. Yun, and J.Y. Han. 2000. A DNA vaccine encoding a conserved *Eimeria* protein induces protective immunity against live *Eimeria acervulina* challenge. *Vaccine* 19: 243-252.

Withanage, G.S., K. Sasai, T. Fukata, T. Miyamoto, **H.S. Lillehoj**, and E. Baba. 2003. Increased lymphocyte subpopulations and macrophages in the ovaries and oviducts of laying hens infected with *Salmonella enterica* serovar *Enteritidis*. *Avian Pathology* 32: 583-590.

Xie, H., R.B. Raybourne, U. Babu, **H.S. Lillehoj**, and R. Heckert. 2003. Effects of CPG on IL-6 and nitric oxide production, surface antigen expression, phagocytosis, proliferation and apoptosis of chicken macrophage cell line HD11 cells. *Developmental and Comparative Immunology* 27(9): 823-834.

Yamaguchi, M., and **H.S. Lillehoj**. 2002. Immunity to protozoan parasites in poultry. In Thankam Mathew, editor, *Modern Concepts of Immunology in Veterinary Medicine-Poultry Immunology*, (Advances in Medical and Veterinary Virology, Immunology and Epidemiology), Thajema Press, p. 183-242.

Yun, C.H. and **H.S. Lillehoj**. 2000. Intestinal immune responses to coccidiosis. *Developmental & Comparative Immunology* 24(2-3): 303-324.

Yun, C.H., **H.S. Lillehoj**, and K.D. Choi. 2000. Chicken IFN- γ monoclonal antibodies and their application in enzyme-linked immunosorbent assay. *Veterinary Immunology and Immunopathology* 73: 297-308.

Yun, C.H., **H.S. Lillehoj**, and K.D. Choi. 2000. *Eimeria tenella* infection induces local IFN- γ production and intestinal lymphocyte subpopulation changes. *Infection and Immunity* 68(3): 1282-1288.

Yun, C.H., **H.S. Lillehoj**, J. Zhu, and W.G. Min. 2000. Kinetic differences in intestinal and systemic interferon- γ and antigen-specific antibodies in chickens infected with *Eimeria maxima*. *Avian Diseases* 44: 305-312.

Zhang, S., **H.S. Lillehoj**, M.C. Jenkins, N. Lally, and W. Min. 2001. *Gallus gallus* mRNA for putative FK506-binding protein (FKBP25 gene). Genbank, Accession Number. AJ312905.

Zhu J., **H.S. Lillehoj**, P.C. Allen, W. Min, C. Van Tassell, T.S. Sonstegard, H.H. Cheng, D.L. Pollock, M. Sadjadi, and M. Emara. 2003. Mapping quantitative trait loci associated with resistance to coccidiosis and growth. *Poultry Science* 82: 9-16.

Zhu, J., **H.S. Lillehoj**, P. Allen, C.H. Yun, D. Pollock, and M. Emara. 2000. Analysis of disease resistance associated parameters in broiler chickens challenged with *Eimeria maxima*. *Poultry Science* 79: 619-625.

Zhu, J., **H.S. Lillehoj**, H.H. Cheng, D. Pollock, M. Sadjadi, and M. Emara. 2001. Screening for highly heterozygous chickens in outbred commercial broiler lines to increase detection power for mapping quantitative trait loci. *Poultry Science* 80: 6-21.

Zhou, H., **H.S. Lillehoj**, and S. Lamont. 2003. Association of chicken interferon- γ genotype and protein level with antibody response kinetics. *Avian Diseases* 46: 869-876.

Lunney JK

Alvarez, B., C. Sanchez, R. Bullido, A. Marina, **J.K. Lunney**, F. Alonso, A. Ezquerro, and J. Dominguez. 2000. Porcine SWC3 is a member of the SIRP family and associates with protein-tyrosine phosphatase SHP-1. *Tissue Antigens* 55: 342-351.

Bannerman, D., M.J. Paape, **J.K. Lunney**. 2004. Immunity: Innate. In W.G. Pond and A.W. Bell, editors, *Encyclopedia of Animal Science*, Marcel Dekker, Inc.: New York. In Press.

Davis, W.C., K. Haverson, A. Saalmüller, H. Yang, **J.K. Lunney**, M.J. Hamilton, and M.D. Pescovitz. 2001. Analysis of monoclonal antibodies reacting with molecules expressed on $\gamma\delta$ T cells. *Veterinary Immunology and Immunopathology* 80: 53-62.

Dawson, H.D., E. Beshah, S. Nishi, G. Solano-Aguilar, M. Morimoto, A. Zhao, K.B. Madden, T.K. Ledbetter, J.P. Dubey, T. Shea-Donohue, **J.K. Lunney**, and J.F. Urban Jr. Localized multi-gene expression patterns support an evolving Th1/Th2-like paradigm in response to infections with *Toxoplasma gondii* and *Ascaris suum* in pigs. *Infection and Immunity*. In Revision.

Dawson, H., S. Nishi, E. Beshah, G. Solano-Aguilar, D. Zarlenga, J.F. Urban, and **J.K.**

- Lunney**. 2002. Functional genomic assessment of porcine disease resistance: real time assays of porcine immune gene expression. Proceedings 7th World Congress on Genetics Applied to Livestock Production, p. 13-41.
- Dawson, H.D., A.R. Royae, S. Nishi, D. Kuhar, W.M. Schnitzlein, F. Zuckermann, J.F. Urban, and **J.K. Lunney**. 2004. Identification of key immune mediators regulating T helper 1 responses in swine. *Veterinary Immunology and Immunopathology* 100: 105-111.
- Gerrits, R.J., **J.K. Lunney**, L.A. Johnson, V.G. Pursel, R.R. Kraeling, G.A. Rohrer, and J.R. Dobrinsky. 2005. Perspectives for artificial insemination and genomics to improve global swine populations. *Theriogenology* 63(2): 283-299.
- Haverson, K., A. Saalmüller, B. Alvarez, F. Alonso, M. Bailey, A.T.J. Bianchi, W.J.A. Boersma, Z. Chen, W.C. Davis, J. Dominguez, H. Engelhardt, A. Ezquerro, L.S. Grosmaire, M.J. Hamilton, E. Hollemweguer, C. Huang, K.V. Khanna, G. Kuebart, G. Lackovic, J.A. Ledbetter, R. Lee, D. Llanes, **J.K. Lunney**, K.C. McCullough, T. Molitor, J. Nielsen, T.A. Niewold, M.D. Pescovitz, J. de la Lastra Perez, Z. Rehakova, H. Salmon, W.M. Schnitzlein, J. Seebach, A. Simon, J. Sinkora, M. Sinkora, C.R. Stocks, A. Summerfield, L. Sver, E. Thacker, I. Valpotic, H. Yang, F.A. Zuckermann, and R. Zwart. 2001. Overview of the Third International Workshop on Swine Leukocyte Differentiation Antigens. *Veterinary Immunology and Immunopathology* 80: 5-23.
- Jauregui, L.H., J. Higgins, D. Zarlenga, J.P. Dubey, and **J.K. Lunney**. 2001. Development of a real-time PCR assay for detection of *Toxoplasma gondii* in pig and mouse tissues. *Journal of Clinical Microbiology* 39: 2065-2071.
- Johnson, P., N.T. Keen, **J.K. Lunney**, and M.J. Sadowsky. 2001. Conference editorial: Agricultural microbes genome 2. *Comparative and Functional Genomics* 2: 10-13. [<http://www3.interscience.wiley.com/cgi-bin/issuetoc?ID=77002017>].
- Lunney, J.K.** 2003. Are there immune gene alleles that determine whether a pig will be healthy? In M. Boggess, editor, *Genetics of Pig Health Symposium*, National Pork Board Press: Des Moines, Iowa, p.63-72.
- Lunney, J.K.** 2003. In search of disease-resistant pigs. *National Hog Farmer*, Apr 15, 2003, p. 30-34.
- Lunney, J.K.**, C. Fossum, G.V. Alm, F. Steinbach, and E. Wattrang. 2002. *Veterinary immunology: opportunities and challenges*. *Trends in Immunology* 23: p. 4-6.
- Lunney, J.K.**, M.J. Paape, and D. Bannerman. 2004. Immunity: Acquired. In W.G. Pond and A.W. Bell, editors, *Encyclopedia of Animal Science*, Marcel Dekker, Inc.: New York. In Press.
- Martens, G W., **J.K. Lunney**, J.E. Baker, and D.M. Smith. 2003. Rapid assignment of swine leukocyte antigen (SLA) haplotypes in pedigreed herds using a polymerase chain reaction based assay. *Immunogenetics* 55: 395-401.
- Navarro, P., R. Christenson, G. Ekhardt, B. Bosworth, **J.K. Lunney**, M. Rothschild, J. Lemke, and J.E. Butler. 2000. Genetic differences in the frequency of the hinge variants of porcine IgA is breed dependent. *Veterinary Immunology and Immunopathology* 73: 287-295.
- Royae, A.R., R. Husmann, H.D. Dawson, G. Calzada-Nova, W.M. Schnitzlein, F.

Zuckermann, and **J.K. Lunney**. 2004. Deciphering the involvement of innate immune factors in the development of the host responses to PRRSV vaccination. *Veterinary Immunology and Immunopathology* 102(3): 199-216.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, D. Zarlenga, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and **J.K. Lunney**. 2001. Cytokine and lymphocyte profile after oral infection with *Toxoplasma gondii* oocysts. *Scandinavian Journal of Immunology* 54: 93.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, D. Zarlenga, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and **J.K. Lunney**. 2001. Cytokine and lymphocyte profile in miniature swine after oral infection with *Toxoplasma gondii* oocysts. *International Journal of Parasitology* 31: 187-195.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, D. Zarlenga, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and **J.K. Lunney**. 2000. Cytokine and lymphocyte profile after *Toxoplasma gondii* infection in miniature swine. *Proceedings of the 16th Congress of the International Pig Veterinary Society, Australia*, p.183.

Solano-Aguilar, G.I., L. Galina, L.W. Douglass, and **J.K. Lunney**. 2002. Developmental immunity in neonatal pigs monitored by the function of peripheral blood mononuclear cells. *Proceedings of the 17th Congress of the International Pig Veterinary Society, Iowa*, p. 259.

Solano-Aguilar, G.I., J.F. Urban, E. Beshah, L. Schopf, H. Kringel, A. Domingo, J.P. Dubey, and **J.K. Lunney**. 2000. Mucosal immune response of young pigs after *Toxoplasma gondii*

infection. *Proceedings of the 16th Congress of the International Pig Veterinary Society, Australia*, p. 171.

Solano-Aguilar, G.I., K. Vengroski, E. Beshah, and **J.K. Lunney**. 2000. Isolation and purification of lymphocyte subsets from gut associated lymphoid tissue in neonatal swine. *Journal of Immunological Methods* 241: 185-199.

Solano-Aguilar, G.I., K. Vengroski, E. Beshah, L.W. Douglass, and **J.K. Lunney**. 2001. Characterization of lymphocyte subsets from mucosal tissues in neonatal swine. *Developmental and Comparative Immunology* 25: 245-263.

Solano-Aguilar, G.I., D. Zarlenga, E. Beshah, K. Vengroski, L. Gasbarre, D. Junker, M. Cochran, C. Weston, D. Valencia, C. Chiang, H. Dawson, J. Urban, and J.K. Lunney. 2002. Limited effect of recombinant porcine Interleukin-12 on porcine lymphocytes due to a low expression of IL-12 beta2 receptor. *Veterinary Immunology and Immunopathology* 89: 133-148. [Correction 95:183, 2003].

Thacker, E., A. Summerfield, K. McCullough, A. Ezquerro, J. Dominguez, F. Alonso, **J.K. Lunney**, J. Sinkora, and K. Haverson. 2001. Summary of workshop findings for porcine myelomonocytic markers. *Veterinary Immunology and Immunopathology* 80: 93-109.

Paape MJ

Burvenich, C., J. Detilleux, **M.J. Paape**, and A.M. Massart-Leen. 2000. Physiological and genetic factors that influence the cows resistance to mastitis especially during early lactation. *Proceedings of IDF Symposium on Immunology of Ruminant Mammary Gland*, p. 9-20.

Diez-Fraile, A., G.M. Tomita, E. Meyer, **M.J. Paape**, and C. Burvenich. 2000. CD62L

expression on circulating neutrophils in relation to *Escherichia coli* mastitis. Proceedings of International Dairy Federation Symposium on Immunology of Ruminant Mammary Gland, p. 177-178.

Ledbetter, T.K., **M.J. Paape**, and L.W. Douglass. 2001. Cytotoxic effects of peroxynitrite, polymorphonuclear neutrophils, free-radical scavengers, inhibitors of myeloperoxidase, and inhibitors of nitric oxide synthase on bovine mammary secretory epithelial cells. American Journal of Veterinary Research 62: 286-293.

Paape, M.J. and A. Contreras. 2000. Limitaciones legales problematica de los recuentos celulares in leche de vaca y cabra en los Estados Unidos. Ovis 67: 13-23.

Paape, M.J., M.I. Duenas, R.P. Wettemann, and L.W. Douglass. 2000. Effects of intramammary infection and parity on calf weaning weight and milk quality in beef cows. Journal of Animal Science 78: 2508-2514.

Paape, M.J., K. Shafer-Weaver, A.V. Capuco, K. Van Oostveldt, and C. Burvenich. 2000. Immune surveillance of mammary tissue by phagocytic cells. Advances in Experimental Medicine and Biology 480: 259-277.

Peters, R.R., S. Domaragiri, **M.J. Paape**, and L.W. Douglass. 2000. Evaluation of 1.6% phenol as a premilking and postmilking teat dip in preventing new bovine intramammary infections. Journal of Dairy Science 83: 1750-1757.

Rainard, P., C. Riollot, B. Poutrel, and **M.J. Paape**. 2000. Phagocytosis and killing of *Staphylococcus aureus* by bovine neutrophils after priming by tumor necrosis factor-alpha and the des-arginine derivative of C5a. American

Journal of Veterinary Research 61: 951-959.

Tomita, G.M., Y. Wang, **M.J. Paape**, B. Poutrel, and P. Rainard. 2000. Influence of bispecific antibodies on the in vitro bactericidal activity of bovine neutrophils against *Staphylococcus aureus*. Journal of Dairy Science 83: 2269-2275.

Van Oostveldt, K., G.M. Tomita, **M.J. Paape**, and C. Burvenich. 2000. Apoptosis of circulating bovine neutrophils during experimentally induced *Escherichia coli* and endotoxin mastitis. Proceedings of International Dairy Federation Symposium on Immunology of Ruminant Mammary Gland, p. 319-325.

Shafer-Weaver KA

Paape, M.J., **K. Shafer-Weaver**, A.V. Capuco, K. Van Oostveldt, and C. Burvenich. 2000. Immune surveillance of mammary tissue by phagocytic cells. Advances in Experimental Medicine and Biology 480: 259-277.

Solano-Aguilar GI

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, **D. Zarlenga**, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and J.K. Lunney. 2001. Cytokine and lymphocyte profile in miniature swine after oral infection with *Toxoplasma gondii* oocysts. International Journal of Parasitology 31: 187-195.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, **D. Zarlenga**, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and J.K. Lunney. 2001. Cytokine and lymphocyte profile after oral infection with *Toxoplasma gondii* oocysts. Scandinavian Journal of Immunology 54: 93.

Solano-Aguilar, G.I., K. Vengroski, E. Beshah, L.W. Douglass, and J.K. Lunney. 2001. Characterization of lymphocyte

subsets from mucosal tissues in neonatal swine. *Developmental and Comparative Immunology* 25(3): 245-63.

Solano-Aguilar, G.I., D. Zarlenga, E. Beshah, K. Vengroski, L. Gasbarre, D. Junker, M. Cochran, C. Weston, D. Valencia, C. Chiang, H. Dawson, J.F. Urban, and J.K. Lunney. 2002. Limited effect of recombinant porcine interleukin-12 on porcine lymphocytes due to a low level of IL-12 beta2 receptor. *Veterinary Immunology and Immunopathology* 89(3-4): 133-48.

Trout JM

Jenkins, M.C., **J. Trout**, M.S. Abrahamsen, C.A. Lancto, J. Higgins, and R. Fayer. 2000. Estimating viability of *Cryptosporidium parvum* oocysts using reverse transcriptase-polymerase chain reaction (RT-PCR) directed at mRNA encoding amyloglucosidase. *Journal of Microbial Methods* 43: 97-106.

Jenkins, M.C., **J. Trout**, and R. Fayer. 2001. Treatment of *Cryptosporidium parvum* sporozoites with antibodies to a recombinant CP47 surface antigen inhibits in vitro parasite development. *Proceedings of the Annual Meeting of the American Association of Veterinary Parasitologists*. Boston, Massachusetts, p. 49.

Jenkins, M.C., **J. Trout**, J. Higgins, and R. Fayer. 2000. The mRNA levels for amyloglucosidase as measured by RT-PCR correlate with infectivity and fecundity of *Cryptosporidium parvum* oocysts for neonatal BALB/c mice. *Proceedings of the International Conference on Emerging and Infectious Diseases*. Late Breakers, p. 9.

Jenkins, M.C., **J. Trout**, C. Murphy, J. Harp, J. Higgins, W. Wergin, and R. Fayer. 1999. Cloning and expression of a

DNA sequence encoding a 41-kilodalton *Cryptosporidium parvum* oocyst wall protein. *Clinical and Diagnostic Laboratory Immunology* 6(6): 912-920.

Tuo W

Goff, W.L., W.C. Johnson, S.M. Parish, G.M. Barrington, **W. Tuo**, R.A. Valdez. 2001. The age-related immunity in cattle to *Babesia bovis* infection involves the rapid induction of IL-12, IFN-gamma and iNOS mRNA expression in the spleen. *Parasite Immunology* 23: 453-461.

Tuo, W., D. Zhuang, D.P. Knowles, W.P. Cheevers, M.S. Sy, and K.I. O'Rourke. 2001. Prp-c and Prp-Sc at the fetal-maternal interface. *Journal of Biological Chemistry* 276: 18229-18234.

Urban JF Jr

Elliott, D.E., **J.F. Urban Jr.**, C.K. Argo, and J.V. Weinstock. 2000. Does the failure to acquire helminthic parasites predispose to Crohn's disease? *The Federation of American Societies for Experimental Biology Journal* 14: 1848-1855.

Finkelman, F.D. and **J.F. Urban**. 2001. The other side of the coin: The protective role of the type 2 (Th2) cytokines. *Journal of Allergy and Clinical Immunology* 107: 772-780.

Kringel, H., J.P. Dubey, E. Beshah, R. Heckert, and **J.F. Urban**. 2001. Effect of CpG oligodeoxynucleotide (ODN)-induced immune activation of porcine immunity to *Toxoplasma gondii*. *American Association of Veterinary Parasitology* 46: 15.

Urban, J.F., L. Schopf, K.B. Madden, N. Nobern-Trauth, F.D. Finkelman. 2001. Nematode biology determines the requirements for expression of IL-4 receptor alpha chain on lymphoid and/or non-lymphoid cells that regulate intestinal

expulsion. The Federation of American Societies for Experimental Biology Journal 15: A690.

Zarlenga DS

Hediger-Weithaler, B.M., V.T. Heussler, R. Eicher, A. Zakher, **D.S. Zarlenga**, L.C. Gasbarre, and A.S. Waldvogel. 2000. Interferon- and interleukin 4 mRNA expression by PBMCs from BVDV seropositive pregnant and non-pregnant cattle. *Veterinary Immunology and Immunopathology* 77: 201-212.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, **D. Zarlenga**, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, and J.K. Lunney. 2001. Cytokine and

lymphocyte profile in miniature swine after oral infection with *Toxoplasma gondii* oocysts. *International Journal of Parasitology* 31: 187-195.

Solano-Aguilar, G.I., E. Beshah, K. Vengroski, **D. Zarlenga**, L. Jauregui, M. Cosio, L.W. Douglass, J.P. Dubey, J.K. Lunney. 2001. Cytokine and lymphocyte profile after oral infection with *Toxoplasma gondii* oocysts. *Scandinavian Journal of Immunology* 54: 93.

Wang, Y., **D.S. Zarlenga**, M.J. Paape, and G.E. Dahl. 2002. Recombinant bovine soluble CD14 sensitizes the mammary gland to lipopolysaccharide. *Veterinary Immunology and Immunopathology* 86(1/2): 115-124.

Boston, Massachusetts

The Jean Mayer USDA Human Nutrition Research Center on Aging

<http://hnrc.tufts.edu/>

Han SN

Chung, H., D. Wu, R. Gay, **S.N. Han**, B. Goldin, R. Bronson, J. Mason, D.E. Smith, and S.N. Meydani. 2003. Effect of age on susceptibility to Azoxymethane-induced colonic aberrant crypt foci formation in C57BL/6JNIA mice. *Journals of Gerontology Biological Sciences and Medical Sciences* 58(5): B400-B405.

Chung, H., D. Wu, **S.N. Han**, R. Gay, B. Goldin, R.E. Bronson, J.B. Mason, D.E. Smith, and S.N. Meydani. 2003. Vitamin E supplementation does not alter Azoxymethane-induced colonic aberrant crypt foci formation in young or old mice. *Journal of Nutrition* 133(2): 528-532.

Han, S.N., L.S. Leka, A.H. Lichtenstein, L.M. Ausman, and S.N. Meydani. 2003. Effect of therapeutic lifestyle change (TLC) diet on immune functions of moderately hypercholesterolemic humans. *Journal of Lipid Research* 44(12): 2304-2310.

Han, S.N., L.S. Leka, A.H. Lichtenstein, L.M. Ausman, E.J. Schaefer, and S.N. Meydani. 2002. Effect of hydrogenated and saturated, relative to polyunsaturated, fat on immune and inflammatory responses of adults with moderate hypercholesterolemia. *Journal of Lipid Research* 43(3): 445-452.

Han, S.N., L.S. Leka, A.H. Lichtenstein, L.M. Ausman, E.J. Schaefer, S.N. Meydani. 2001. Hydrogenated fat containing diets high in trans fatty acids do not adversely affect immune function in humans. *The Federation of American Societies for Experimental Biology Journal* 15(5): A1094.

Han, S.N., M. Meydani, D. Wu, B.S. Bender, D.E. Smith, J. Vina, G. Cao, R.L. Prior, and S.N. Meydani. 2000. Effect of long-term dietary antioxidant supplementation on influenza virus infection. *Journals of Gerontology Biological Sciences and Medical Sciences* 55A: B1-B8.

Han, S.N., D. Wu, W.K. Ha, Beharka, A.A., Smith, D., B.S. Bender, and S.N. Meydani. 2000. Vitamin E supplementation increases Th1 cytokine production in old mice infected with influenza virus infection. *Immunology* 100: 487-493.

Meydani S

Adolfsson, O., B.T. Huber, and **S.N. Meydani**. 2001. Vitamin E increases IL-2 production by T cells from old mice through a post-transcriptional mechanism. *The Federation of American Societies for Experimental Biology Journal* 15(5): A293.

Beharka, A.A., M. Meydani, D. Wu, L.S. Leka, A. Meydani, and **S.N. Meydani**. 2001. IL-6 production does not increase with age. *Journals of Gerontology Biological Sciences and Medical Sciences* 56A(2): B81-B88.

Beharka, A.A., S. Paiva, L.S. Leka, J. Ribaya-Mercado, R.M. Russell, and **S.N. Meydani**. 2001. Effect of age on the gastrointestinal-associated mucosal immune response of humans. *Journals of Gerontology Biological Sciences and Medical Sciences* 56A(5): B218-B223.

Bugianesi, R., M Serafini, F. Simone, D. Wu, **Meydani, S.,** A. Ferro-Luzzi, E. Azzini, and G. Maiani. 2000. High-performance liquid chromatography with coulometric electrode array detector for the determination of quercetin levels in

cells of the immune system. *Analytical Biochemistry* 284: 296-300.

Chung, H., D. Wu, R. Gay, B. Goldin, R. Bronson, J. Mason, D. Smith, and **S.N. Meydani**. 2001. Aged mice are more susceptible to Azoxymethane (AOM)-induced colon cancer. *The Federation of American Societies for Experimental Biology Journal* 15: A994.

Han, S.N., L.S. Leka, A.H. Lichtenstein, L.M. Ausman, E.J. Schaefer, and **S.N. Meydani**. 2001. Hydrogenated fat containing diets high in trans fatty acids do not adversely affect immune function in humans. *The Federation of American Societies for Experimental Biology Journal* 15(5): A1094.

Han, S.N., M. Meydani, D. Wu, B.S. Bender, D.E. Smith, J. Vina, G. Cao, R.L. Prior, and **S.N. Meydani**. 2000. Effect of long-term dietary antioxidant supplementation on influenza virus infection. *Journals of Gerontology Biological Sciences and Medical Sciences* 55A: B1-B8.

Han, S.N., D. Wu, W.K. Ha, A.A. Beharka, D. Smith, B.S. Bender, **S.N. Meydani**. 2000. Vitamin E supplementation increases Th1 cytokine production in old mice infected with influenza virus infection. *Immunology* 100: 487-493.

Hayek, M.G. and **S.N. Meydani**. 2000. Effect of vitamin E on infection and immunity. *The Science of Geriatrics* 2: 583-594.

King, J.C., R.E. Black., M.P. Doyle, K.L. Fritsche, B.H. Halbrook, O.A. Levander, **S.N. Meydani**, W.A. Walker, and C.E. Woteki. 2000. Foodborne illnesses and nutritional status: A statement from an American Society for Nutritional Sciences working group. *Journal of Nutrition* 130: 2613-2617.

Meydani, S.N. 2000. Advances in preventive medicine: nutrition and immune function in the elderly. In A.L. Goldstein, editor, *Frontiers in Biomedicine*. Kluwer Academic/Plenum Publishers: New York, p. 19-29.

Meydani, S.N. and A.A. Beharka. 2000. Vitamin E and immune response in the aged. *Bibliotheca Nutrition et Dieta* 55: 148-158.

Meydani, S.N., W. Fawzi, and S.N. Han. 2000. The effect of vitamin deficiencies (E and A) and supplementation on infection and immune response. *Nutrition, Immunity, and Infectious Diseases in Infants and Children* 45: 213-241.

Meydani, S.N., T.S. McElreavy, and M. Meydani. 2000. Antioxidants and immune function. In M.A. Fiatarone, editor, *Exercise, Nutrition, and the Older Woman*, CRC Press: Boca Raton, Florida, p. 183-205.

Meydani, S.N. and M. Meydani. 2000. Effect of dietary supplementation with black currant seed oil on the immune response of healthy elderly subjects and dietary conjugated linoleic acid influences the immune response of young and old C57BL/6NCrIBR mice. *Annual Bibliography of Significant Advances in Dietary Supplement Research*, p. 25-26.

Wu DY

D. Wu, M.G. Hayek, and S.N. Meydani. 2001. Vitamin E and macrophage cyclooxygenase regulation in the aged. *Journal of Nutrition* 131: 382S-388S.

Wu, D., M. Meydani, S.N. Han, L.S. Leka, and S.N. Meydani. 2000. Effect of dietary supplementation with fish oil in combination with different levels of vitamin E on immune response in healthy elderly human subjects. *The Federation of American Societies for Experimental Biology Journal* 14: A238.

East Lansing, Michigan

The Avian Disease and Oncology Laboratory

<http://www.msu.edu/~arsadol/index.htm>

Bacon LD

Bacon, L.D., H.D. Hunt, and H.H. Cheng. 2001. Genetic resistance to Marek's Disease. In K. Hirai, editor, *Marek's Disease*, Springer Verlag: New York, p. 121-142.

Bacon, L.D., H.D. Hunt, and H.H. Cheng. 2000. A review of the development of chicken lines to resolve genes determining resistance to diseases. *Poultry Science* 79: 1082-1093.

Witter, R.L., **L.D. Bacon,** H.D. Hunt, R.E. Silva, and A.M. Fadly. 2001. Avian

leukosis virus subgroup J infection profiles in broiler breeder chickens: associations with virus transmission to progeny. *Avian Disease* 44: 913-931.

Cheng HH

Bacon, L.D., H.D. Hunt, and **H.H. Cheng.** 2001. Genetic resistance to Marek's Disease. In K. Hirai, editor, *Marek's Disease*, Springer Verlag: New York, p. 121-142.

Bacon, L.D., H.D. Hunt, and **H.H. Cheng.** 2000. A review of the development of chicken lines to resolve genes determining

resistance to diseases. *Poultry Science* 79: 1082-1093.

Liu, H.C., H.J. Kung, J.E. Fulton, R.M. Morgan, and **H.H. Cheng**. 2001. Growth hormone interacts with the Marek's disease virus SORF2 protein and is associated with disease resistance in chicken. *Proceedings of the National Academy of Sciences USA*, 98: 9203-9208.

Maki-Tanila, A., **H.H. Cheng**, M.E. Delany, J. Burnside, and S. Mizuno. 2000. First report on chicken genes and chromosomes. *Cytogenetics and Cell Genetics* 90: 169-218.

Schmid, M., I. Nanda, M. Guttenbach, C. Steinlein, H. Hoehn, M. Scharlt, T. Haaf, S. Weigend, R. Fries, J.M. Buerstedde, K. Wimmers, D.W. Burt, J. Smith, S. A'Hara, A. Law, D.K. Griffin, N. Bumstead, J. Kaufman, P.A. Phomsom, T. Burke, M.A.M. Groenen, and **H.H. Cheng**. 2000. First report on chicken genes and chromosomes. *Cytogenetics and Cell Genetics* 90: 169-218.

Suchyta, S.P., **H.H. Cheng**, J. Burnside, and J.B. Dodgson. 2001. Comparative mapping of chicken anchor loci orthologous to genes on human chromosomes 1, 4 and 9. *Animal Genetics* 32: 12-18.

Yonash, N., **H.H. Cheng**, J. Hillel, E.D. Heller, and A. Cahaner. 2001. DNA microsatellites linked to quantitative trait loci affecting antibody response and survival rate in meat-type chickens. *Poultry Science* 80: 22-28.

Zhu, J., H.S. Lillehoj, P. Allen, **H.H. Cheng**, M.G. Emara, M. Sadjadi, and D. Pollock. 2001. Mapping quantitative trait loci affecting disease resistance to avian coccidiosis. *Proceedings of the 73rd*

Northeastern Conference on Avian Diseases, p. 34.

Zhu, J., H.S. Lillehoj, **H.H. Cheng**, D. Pollock, M. Sadjadi, and M. Emara. 2001. Screening for highly heterozygous chickens in outbred commercial broiler lines to increase detection power for mapping quantitative trait loci. *Poultry Science* 80(1): 6-12

Zhu, J., H. Lillehoj, M. Emara, **H. Cheng**, D. Pollock, M. Sadjadi, P. Allen, and C. Yun. 2000. Mapping QTL associated with disease resistance to coccidiosis in commercial broiler chickens. 89th *Poultry Science Meeting*, p. 9.

Zhu, J., H.S. Lillehoj, C. van Tassel, M. Emara, P. Allen, **H. Cheng**, D. Pollock, M. Sadjadi, and T. Sonstegard. 2001. Mapping QTL loci affecting growth and disease resistance to avian coccidiosis. *Proceedings of the International Animal Agricultural and Food Sciences Conference*, p. 43.

Fadly AM

Fadly, A.M., R.E. Silva, and L.F. Lee. 2000. Antigenic characterization of selected field isolates of subgroup J avian leukosis virus. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, *Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses*, Rauschholzhausen, Germany, p. 13-22.

Fadly, A.M. 2000. Isolation and identification of avian leukosis viruses: a review. *Avian Pathology* 29: 529-535.

Hunt, H.D., V. Leathers, C. Lamichhane, G. Rosales, E. Jensen, J. McKay, and **A. Fadly**. 2000. Analysis of serum antibody to ALV-J in commercial broilers: correlation between ELISA, flow cytometry and Virus Neutralization. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, *Proceedings of the International Symposium*

on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 127-140.

Hunt, H., B. Lupiani, and **A.M. Fadly**. 2000. Recombination between ALV-J and endogenous subgroup E viruses. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 50-60.

Hussain, A.I., V. Shanmugam, W.M. Switzer, S.X. Tsang, **A.M. Fadly**, D. Thea, R. Helfand, W. Bellini, T.M. Folks, and W. Heneine. 2001. Lack of evidence of endogenous avian leukosis virus and endogenous avian retrovirus transmission to measles, mumps and rubella vaccine recipients. *Emerging Infectious Diseases* 7: 66-72.

Lee, L.F., **A.M. Fadly**, and H.D. Hunt. 2000. Avian leukosis virus subgroup J envelope gene product for diagnosis and immunogenic composition. United States Patent #6,146,641.

Williams, S.M., W.M. Reed, and **A.M. Fadly**. 2000. Influence of age of exposure on the response of line 0 and line 63 chickens to infection with subgroup J avian leukosis virus. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 67-76.

Witter, R.L., L.D. Bacon, H.D. Hunt, R.E. Silva, and **A.M. Fadly**. 2001. Avian leukosis virus subgroup J infection profiles in broiler breeder chickens: associations with virus transmission to progeny. *Avian Disease* 44: 913-931.

Hunt HD

Bacon, L.D., **H.D. Hunt**, and H.H. Cheng. 2001. Genetic resistance to Marek's Disease. In K. Hirai, editor, *Marek's Disease*, Springer Verlag: New York, p. 121-142.

Bacon, L.D., **H.D. Hunt**, and H.H. Cheng. 2000. A review of the development of chicken lines to resolve genes determining resistance to diseases. *Poultry Science* 79: 1082-1093.

Hunt, H.D., V. Leathers, C. Lamichhane, G. Rosales, E. Jensen, J. McKay, and A. Fadly. 2000. Analysis of serum antibody to ALV-J in commercial broilers: correlation between ELISA, flow cytometry and Virus Neutralization. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 127-140.

Hunt, H.D., B. Lupiani, and A.M. Fadly. 2000. Recombination between ALV-J and endogenous subgroup E viruses. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 50-60.

Hunt, H.D., B. Lupiani, M.M. Miller, I. Gimeno, L.F. Lee, and M.S. Parcells. 2001. Marek's disease virus down regulates surface expression of MHC (B complex) class I (BF) glycoproteins during active but not latent infection of chicken cells. *Virology* 282: 198-205.

Lee, L.F., A.M. Fadly, and **H.D. Hunt**. 2000. Avian leukosis virus subgroup J envelope gene product for diagnosis and immunogenic composition. United States Patent #6,146,641.

Parcells, M.S., S.F. Lin, R.L. Dienglewicz, V. Majerciak, D.R. Robinson, H.C. Chen, Z.

Wu, G.R. Dubyak, P. Brunovskis, **H.D. Hunt**, L.F. Lee, and H.J. Kung. 2001. Marek's disease virus (MDV) encodes an interleukin-8 homolog (vIL-8): Characterization of the vIL-8 protein and a vIL-8 deletion mutant MDV. *Journal of Virology* 75(11): 5159-5173.

Preisler, M.T., R.J. Tempelman, R.J. Erskine, **H.D. Hunt**, and J.L. Burton. 2000. Glucocorticoid receptor expression profiles in mononuclear leukocytes of periparturient Holstein cows. *Journal of Dairy Science* 83: 38-47.

Preisler, M.T., P.S.D. Weber, R.J. Tempelman, R.J. Erskine, **H.D. Hunt**, and J.L. Burton. 2000. Glucocorticoid receptor down-regulation in neutrophils of periparturient cows. *American Journal of Veterinary Research* 61: 14-19.

Witter, R.L., L.D. Bacon, **H.D. Hunt**, R.E. Silva, and A.M. Fadly. 2001. Avian leukosis virus subgroup J infection profiles in broiler breeder chickens: associations with virus transmission to progeny. *Avian Disease* 44: 913-931.

Lee LF

Fadly, A.M., R.F. Silva, and **L.F. Lee**. 2000. Antigenic characterization of selected field isolates of subgroup J avian leukosis virus. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, *Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses*, Rauschholzhausen, Germany, p. 13-22.

Hunt, H.D., B. Lupiani, M.M. Miller, I.M. Gimeno, **L.F. Lee**, and M.S. Parcells. 2001. Marek's disease virus down regulates surface expression of MHV (B complex) class I (BF) glycoproteins during active but not latent

infection of chicken cells. *Virology* 282(1): 198- 205.

Kung, H.J., L. Xia, P. Brunovskis, D. Li, J.L. Liu, and **L.F. Lee**. 2001. Meq: An MDV-specific bZIP transactivator with transforming properties. *Current Topics in Microbiology and Immunology*. 255: 245-260.

Lee, L.F., A.M. Fadly, and H.D. Hunt. 2000. Avian leukosis virus subgroup J envelope gene product for diagnosis and immunogenic composition. United States Patent #6,146,641.

Lee, L.F., P. Wu, D. Sui, D. Ren, J. Kamil, H.J. Kung, and R.L. Witter. 2000. The complete unique long sequence and the overall genomic organization of the GA strain of Marek's disease virus. *Proceedings of the National Academy of Sciences USA* 97(11): 6091- 6096.

Lupiani, B., **L.F. Lee**, and S.M. Reddy. 2001. Protein-coding content of the sequence of Marek's disease virus. *Current Topics in Microbiology and Immunology* 255: 159-190.

Parcells, M.S., S.F. Lin, R.L. Dienglewicz, V. Majerciak, D.R. Robinson, H.C. Chen, Z. Wu, G.R. Dubyak, P. Brunovskis, H.D. Hunt, **L.F. Lee**, and H.J. Kung. 2001. Marek's disease virus (MDV) encodes an interleukin-8 homolog (vIL-8): Characterization of the vIL-8 protein and a vIL-8 deletion mutant MDV. *Journal of Virology* 75(11): 5159-5173.

Silva, R.F., **L.F. Lee**, and G.F. Kutish. 2001. The genomic structure of Marek's disease virus. *Current Topics in Microbiology and Immunology* 255: 143-158.

Wu, P., W.M. Reed, and **L.F. Lee**. 2001. Glycoproteins H and L of Marek's disease virus form a hetero-oligomer essential for

translocation and cell surface expression. Archives of Virology 146: 983-992.

Reddy SM

Lupiani, B., L.F. Lee, and **S.M. Reddy**. 2001. Protein-coding content of the sequence of Marek's disease virus. Current Topics in Microbiology and Immunology 255: 159- 190.

Miles, A.M., **S.M. Reddy**, R.W. Morgan. 2001. Coinfection of specific-pathogen-free chickens with Marek's disease virus (MDV) and chicken infectious anemia virus: effect of MDV pathotype. Avian Diseases 45(1): 9-18.

Reddy, S.M., R.L. Witter, and I.M. Gimeno. 2000. Development of a quantitative- competitive polymerase chain reaction assay for serotype 1 Marek's disease virus. Avian Diseases 44(4): 770-775.

Silva RF

Fadly, A.M., **R.F. Silva**, and L.F. Lee. 2000. Antigenic characterization of selected field isolates of subgroup J avian leukosis virus. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 13-22.

Silva, R.F., L.F. Lee, and G.F. Kutish. 2001. The genomic structure of Marek's disease virus. Current Topics in Microbiology and Immunology 255: 143-158.

Witter, R.L., L.D. Bacon, H.D. Hunt, **R.F. Silva**, and A.M. Fadly. 2001. Avian leukosis virus subgroup J infection

profiles in broiler breeder chickens: associations with virus transmission to progeny. Avian Disease 44: 913-931.

Witter RL

Lee, L.F., P. Wu, D. Sui, D. Ren, J. Kamil, H.J. Kung, and R.L. Witter. 2000. The complete unique long sequence and the overall genomic organization of the GA strain of Marek's disease virus. Proceedings of the National Academy of Sciences USA 97(11): 6091- 6096.

Reddy, S.M., **R.L. Witter**, and I.M. Gimeno. 2000. Development of a quantitative-competitive polymerase chain reaction assay for serotype 1 Marek's disease virus. Avian Diseases 44(4): 770-775.

Witter, R.L. 2001. Part II: Marek's disease vaccines - how the virus fights back. World Poultry 17(2): 38-39.

Witter, R.L. 2001. Part 1: Marek's disease vaccines - the viral battlefield. World Poultry 17(2): 42-44.

Witter, R.L., L.D. Bacon, H.D. Hunt, R.E. Silva, and A.M. Fadly. 2001. Avian leukosis virus subgroup J infection profiles in broiler breeder chickens: associations with virus transmission to progeny. Avian Disease 44: 913-931.

Witter, R.L. 2001. Protective efficacy of Marek's disease vaccines. Current Topics in Microbiology and Immunology 255: 58-90.

Witter, R.L. 2000. Determinants of early transmission of ALV-J in commercial broiler breeder chickens. In E.F. Kaleta, L.N. Payne, and U. Heffels-Redmann, editors, Proceedings of the International Symposium on ALV-J and Other Avian Retroviruses, Rauschholzhausen, Germany, p. 216-225.

Mississippi State, Mississippi

Poultry Research Unit

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=64-06-20-00>

Branton SL

Branton, S. L. 2001. *Mycoplasma* vaccines in layer chickens. 45th Annual Meeting of State Contact Representatives and Inspectors. Gulfport, Mississippi, May 15, 2001. National Poultry Improvement Plan.

Branton, S. L. 2001. Making sense of sensitive *Mycoplasma*. *Tecnica International* January: 56-58.

Branton, S.L., B.D. Lott, J.D. May, W.R. Maslin, G.T. Pharr, S.D. Bearson, S.D. Collier, and D.L. Boykin. 2000. The effects of ts-11 strain *Mycoplasma gallisepticum* vaccination in commercial layers on egg production and selected egg quality parameters. *Avian Diseases* 44: 618-623.

Stoneville, Mississippi

Catfish Genetics Research Unit

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=64-02-35-00>

Bilodeau AL

Bilodeau, A.L., B.C. Small, and W.R. Wolters. 2003. Pathogen loads, clearance and plasma cortisol response in channel catfish, *Ictalurus punctatus* (Rafinesque), following challenge with *Edwardsiella ictaluri*. *Journal of Fish Diseases* 26(7): 433-437.

different genes encode heavy chains of membrane and secreted IgD. *Journal of Immunology* 169(5): 2488-2497.

Quiniou, S.M. 2001. Physiological and Genetic Aspects of Channel Catfish Immunity: I. Effect of Low Temperatures Stress on Distribution of Epidermal Mucus Cells as a Factor in Saprolegniosis and II. Linkage and Function of Major Histocompatibility Genes. Thesis.

Quiniou SM

Barker, K.S., **S.M. Quiniou**, M.R. Wilson, E. Bengten, T.B. Stuge, G.W. Warr, L.W. Clem, and N.W. Miller. 2000. Telomerase expression and telomere length in immortal leukocyte lines from channel catfish. *Developmental and Comparative Immunology* 24(6-7): 583-595.

Shen, L., T.B. Stuge, H. Zhou, M. Khayat, K.S. Barker, **S.M. Quiniou**, M. Wilson, E. Bengten, V.G. Chinchar, L.W. Clem, and N.W. Miller. 2002. Channel catfish cytotoxic cells: a mini-review. *Developmental and Comparative Immunology* 26(2): 141-149.

Bengten, E., **S.M. Quiniou**, T.B. Stuge, T. Katagiri, N.W. Miller, L.W. Clem, G.W. Warr, and M. Wilson. 2002. The IgH locus of the channel catfish, *Ictalurus punctatus*, contains multiple constant region gene sequences:

Waldbieser GC

Chen, H., **G.C. Waldbieser**, C.D. Rice, B. Elibol, W.R. Wolters, and L.A. Hanson. 2002. Isolation and characterization of channel catfish natural resistance associated macrophage protein gene. *Developmental and Comparative Immunology* 26(6): 517-531.

Yang, F., T. Ventura-Holman, **G.C. Waldbieser**, and C.J. Lobb. 2003. Structure, genomic organization, and

phylogenetic implications of six new VH families in the channel catfish. *Molecular Immunology* 40(5): 247-260.

Columbia, Missouri

Animal Physiology Research Unit

http://www.agron.missouri.edu/ars_columbia/apru.html

Carroll JA

Carroll, J.A., D. Alberts, D.J. Parzik, D.M. Stocco, and T.H. Welsh. 2001. Influence of corticotropin-releasing hormone (CRH) on the expression of steroidogenic acute regulatory (StAR) protein in neonatal pigs derived by Caesarian section or natural birth. *Journal of Animal Science* 79(Suppl.1): 384.

Carroll, J.A., K.L. Fritsche, J.D. Spencer, A.M. Gaines, G.L. Allee, R.L. Matteri, H.G. Kattesh, M.P. Roberts, L.A. Beausang, and M.E. Zannelli. 2001. Effect of fish oil supplementation on the young pig's immunological response to an endotoxic challenge. *Journal of Animal Science* 79(Suppl.2): 34.

Carroll, J.A., R.L. Matteri, C.J. Dyer, L.A. Beausang, and M.E. Zannelli. 2001. Impact of environmental temperature on response of neonatal pigs to an endotoxin challenge. *American Journal of Veterinary Research*. 62: 561-566.

Carroll, J.A., K.J. Touchette, G.L. Allee, D. Alberts, D.M. Stocco, and T.H. Welsh. 2001. Effect of E. Coli exposure on production of ACTH, steroidogenic acute regulatory (StAR) protein and cortisol in neonatal pigs. *Journal of Animal Science* 79(Suppl.2): 20.

Maddock, K.R., E.P. Berg, C.A. Stahl, M.L. Linville, and **J.A. Carroll**. 2001.

The effects of alpha-lipoic acid (LA) on performance and health of weaned neonatal pigs. *Journal of Animal Science* 79(Suppl. 1): 398.

Matteri, R.L., **J.A. Carroll**, and C.J. Dyer. 2000. Neuroendocrine responses to stress. In G.P. Moberg and J.A. Mench, editors, *The Biology of Animal Stress: Basic Principles and Implications for Animal Welfare*, CABI Publishing: Oxon, UK, p. 43-76.

Matteri, R.L., C.J. Dyer, K.J. Touchette, **J.A. Carroll**, and G.L. Allee. 2000. Effects of weaning on somatotrophic gene expression and circulating levels of insulin-like growth factor-1 (IGF-1) and IGF-2 in pigs. *Domestic Animal Endocrinology* 19: 247-259.

McKee, C.A., **J.A. Carroll**, S.D. Eicher, M.E. Zannelli, L.A. Beausang, and R.L. Matteri. 2001. Supplemental vitamin C and beta-glucan alter growth and the LPS-induced immunological responses in young pigs. *Journal of Animal Science* 79(Suppl.2): 34.

Seaman, J.S., E.P. Berg, T.J. Safranski, and **J.A. Carroll**. 2001. Effect of dexamethasone treatment on growth in neonatal swine. *Journal of Animal Science* 79(Suppl.1): 30.

Seaman, J.S., T.J. Safranski, R.L. Matteri, and **J.A. Carroll**. 2001. Dexamethasone treatment and increased growth in neonatal piglets. *Journal of Animal Science* 79(Suppl.2): 46.

Strauch, T.A., **J.A. Carroll**, C.A. Abbey, N.H. McArthur, S.K. Durham, M.C. Lucy, R.D. Randel, and T.H. Welsh. 2001. Evaluation of somatotrophic gene expression and function in 3/4 Angus or Brahman steers. *Journal of Animal Science* 79(Suppl.2): 19.

Whitley, N.C., C.R. Barb, R.R. Kraeling, J.B. Barrett, G.B. Rampacek, **J.A. Carroll**, and D.H. Keisler. 2000. Feed intake and serum GH, LH and cortisol in gilts after intracerebroventricular or intravenous injection of urocortin. *Domestic Animal Endocrinology* 19: 209-221.

Donald Danforth Plant Science Center, Plant Genetics Research Unit

http://www.agron.missouri.edu/ars_columbia/pgru.html

Herman EM

Helm R.M., G. Cockrell, C. Connaughton, C.M. West, **E.M. Herman**, H.A. Sampson, G.A. Bannon, and A.W. Burks. 2000. Mutational analysis of the IgE-binding epitopes of P34/Gly m Bd 30K. *Journal of Allergy and Clinical Immunology* 105(2 pt 1): 378-384.

Herman, E.M. 2004. Allergenic responses to legume proteins. In C. Brummer and R. Wilson, editors, *Genomics for Legume Crops*, AOCS Press: Champagne, Illinois, pp. 326-348.

Herman, E.M. 2003. Genetically modified soybeans and food allergies. *Journal of Experimental Botany* 54(386): 1317-1319.

Herman, E.M., R.M. Helm, R. Jung, and A.J. Kinney. 2003. Genetic modification removes an immunodominant allergen from soybean. *Plant Physiology* 132(1): 36-43.

Okinaka, Y., Y. Ching-Hong, **E.M. Herman**, A. Kinney, and N.T. Keen. 2002. The P34 syringolide elicitor receptor interacts with a soybean photorespiration enzyme, NADH-dependent hydroxypyruvate reductase. *Plant Molecular Microbe Interactions* 15: 1213-1218.

Clay Center, Nebraska

Roman L. Hruska U.S. Meat Animal Research Center

<http://www.marc.usda.gov/>

Chitko-McKown CG

Heaton, M.P., W.M. Grosse, S.M. Kappes, J.W. Keele, **C.G. Chitko-McKown**, L.V. Cundiff, A. Braun, D.P. Little, and W.W. Laegreid. 2001. Estimation of DNA sequence diversity in bovine cytokine genes. *Mammalian Genome* 12(1): 32-37.

Laegreid, W.W., M.P. Heaton, J.E. Keen, W.M. Grosse, **C.G. Chitko-McKown**, T.P. Smith, J.W. Keele, G.L. Bennett, and T.E. Besser. 2002. Association of bovine neonatal Fc receptor alpha-chain gene (FCGRT) haplotypes with serum IgG concentration in newborn calves. *Mammalian Genome* 13(12): 704-710.

Orient Point, New York

Plum Island Animal Disease Center

<http://www.ars.usda.gov/plum/>

Barrera J

Rodriguez, L.L., **J. Barrera**, E. Kramer, J. Lubroth, F. Brown, and W.T. Golde. 2003. A synthetic peptide containing the consensus sequence of the G-H loop region of foot-and-mouth disease virus type-O VP1 and a promiscuous T-helper epitope induces peptide-specific antibodies but fails to protect cattle against viral challenge. *Vaccine* 21(25-26): 3751-3756.

Bautista EM

Bautista, E.M., G.S. Ferman, and W.T. Golde. 2003. Induction of lymphopenia and inhibition of T cell function during acute infection of swine with foot and mouth disease virus (FMDV). *Veterinary Immunology and Immunopathology* 92(1-2): 61-73.

Bautista, E.M., D. Gregg, and W.T. Golde. 2002. Characterization and functional analysis of skin-derived dendritic cells from swine without a requirement for in vitro propagation. *Veterinary Immunology and Immunopathology* 88(3-4): 131-148.

Bautista, E.M., K.S. Faaberg, D. Mickelson, and E.D. McGruder. 2002. Functional properties of the predicted helicase of porcine reproductive and respiratory syndrome virus. *Virology* 298(2): 258-270.

Baxt B

Duque, H. and **B. Baxt**. 2001. Analysis of the bovine integrin subunits beta1 and

beta5 as receptors for foot-and-mouth disease virus. *American Society for Virology*, p. 120.

Neff, S. and **B. Baxt**. 2001. The ability of integrin alpha V beta 3 to function as a receptor for foot-and-mouth disease virus is not dependent on the presence of complete subunit cytoplasmic domains. *Journal of Virology* 75: 527-532.

Brum MC

Wu, Q., **M.C. Brum**, L. Caron, M. Koster, M.J. Grubman. 2003. Adenovirus-mediated type I interferon expression delays and reduces disease signs in cattle challenged with foot-and-mouth disease virus. *Journal of Interferon and Cytokine Research* 23(7): 359-368.

Caron L

Caron, L., E.F. Flores, R. Weiblen, C.F.C. Scherer, L.F. Irigoyen, P.M. Roehle, A. Odeon, and J.H. Sur. 2002. Latent infection by bovine herpesvirus type-5 in experimentally infected rabbits: virus reactivation, shedding and recrudescence of neurological disease. *Veterinary Microbiology* 84(4): 285-295.

Golde WT

Mbow, M.L., R.D. Gilmore Jr., B. Stevenson, **W.T. Golde**, J. Piesman, and B.J. Johnson. 2002. *Borrelia burgdorferi*-specific monoclonal antibodies derived from mice primed with Lyme disease spirochete-infected *Ixodes scapularis* ticks. *Hybridoma and Hybridomics* 21(3): 179-182.

Grubman MJ

Chinsangaram, J., M. Koster, and **M.J. Grubman**. 2001. Inhibition of L-deleted foot- and-mouth disease virus replication by interferon alpha/beta involves double-stranded RNA dependent protein kinase. *Journal of Virology* 75: 5498-5503.

Grubman, M J., J. Chinsangaram, M. Koster, and M.P. Moraes. 2001. Foot and Mouth Disease Virus Vaccine, Provisional Patent, Docket Number: 0128.01, Serial Number: 60/286, 345, Date Filed: 4/26/2001.

Mayr, G.A., V. O'Donnell, J. Chinsangaram, P.W. Mason, and **M.J. Grubman**. 2001. Immune responses and protection against foot-and-mouth disease virus (FMDV) challenge in swine vaccinated with adenovirus-FMDV constructs. *Vaccine* 19: 2152-2162.

Mason PW

Henry, T., V. O'Donnell, J. Pacheco, J. Gregory, and **P. Mason**. 2001. Analyses of foot-and-mouth disease clones expressing truncated nonstructural 3A proteins reveal genetic plasticity. *American Society for Virology*, p. 205.

Knowles, N., P. Davies, T. Henry, V. O'Donnell, J. Pacheco, and **P. Mason**. 2001. Emergence in Asia of foot-and-mouth disease viruses with altered host Range: characterization of alterations in the 3A protein. *Journal of Virology* 75: 1551-1556.

Mason, P. and N. Knowles. 2000. Emergence and re-emergence of foot-and-mouth disease in Asia: identification and characterization of new strains of an old enemy. *United States Animal Health Association*, p. 291-300.

Mayr, G.A., V. O'Donnell, J. Chinsangaram, **P.W. Mason**, and M.J. Grubman. 2001. Immune responses and protection against foot-and-mouth disease virus (FMDV) challenge in swine vaccinated with adenovirus-FMDV constructs. *Vaccine* 19: 2152-2162.

O'Donnell, V., J. Pacheco, T. Henry, and **P. Mason**. 2001. Attenuation of O Taiwan foot-and-mouth disease virus (FMDV) in bovines is not associated with an altered subcellular distribution of the 3A protein. *American Society for Virology*, p. 136.

Pacheco, J. and **P. Mason**. 2001. Multiple factors affect bovine attenuation of Asian FMDV strains. *American Society for Virology*, p. 179.

Neilan JG

Sur, J.H., T.G. Burrage, **J.G. Neilan**, D.L. Rock, and L. Zsak. 2001. African Swine Fever Virus multigene family 360 and 530 genes affect virulence, pathogenesis and immune response in swine. *American Society for Virology*, p. 84.

Zsak, L., Z. Lu, T.G. Burrage, **J.G. Neilan**, G.F. Kutish, D.M. Moore, and D.L. Rock. 2001. Africa Swine Fever virus multigene family 360 and 530 genes are novel macrophage host range determinants. *Journal of Virology* 75: 3066-3076.

Zsak, L., J.H. Sur, T.G. Burrage, **J.G. Neilan**, and D.L. Rock. 2001. African Swine Fever Virus multigene families 360 and 530 genes promote infected macrophage survival. *Miami Nature Biotechnology Winter Symposium*, p. 97.

Rock DL

Sur, J.H., T.G. Burrage, J.G. Neilan, **D.L. Rock**, and L. Zsak. 2001. African Swine Fever Virus multigene family 360 and 530 genes affect virulence, pathogenesis and

immune response in swine. American Society for Virology, p. 84.

Tulman, E.R. and **D.L. Rock**. 2001. Novel virulence and host range genes of African Swine Fever Virus. *Current Opinion in Microbiology* 4: 456-461.

Zsak, L., Z. Lu, T.G. Burrage, J.G. Neilan, G.F. Kutish, D.M. Moore, and **D.L. Rock**. 2001. Africa Swine Fever Virus multigene family 360 and 530 genes are novel macrophage host range determinants. *Journal of Virology* 75: 3066-3076.

Zsak, L., J.H. Sur, T.G. Burrage, J.G. Neilan, and **D.L. Rock**. 2001. African Swine Fever Virus multigene families 360 and 530 genes promote infected macrophage survival. Miami Nature Biotechnology Winter Symposium, p. 97.

Rodriguez LL

Flanagan, E.B., J.M. Zamparo, L.A. Ball, **L.L. Rodriguez**, and G.W. Wertz. 2001. Rearrangement of the genes of vesicular stomatitis virus eliminates clinical disease in the natural host: new strategy for vaccine development. *Journal of Virology* 75(13): 6107-6114.

Rodriguez, L.L. 2002. Emergence and re-emergence of vesicular stomatitis in the United States. *Virus Research* 85(2): 211-219.

Rodriguez, L.L., J. Barrera, E. Kramer, J. Lubroth, F. Brown, and W.T. Golde. 2003. A synthetic peptide containing the consensus sequence of the G-H loop region of foot-and-mouth disease virus type-O VP1 and a promiscuous T-helper epitope induces peptide-specific antibodies but fails to protect cattle against viral challenge. *Vaccine* 21(25-26): 3751-3756.

Rodriguez, L.L., S.J. Pauszek, T.A. Bunch, and K.R. Schumann. 2002. Full-length genome analysis of natural isolates of vesicular stomatitis virus (Indiana 1 serotype) from North, Central and South America. *Journal of General Virology* 83(Pt 10): 2475-2483.

Scherer C

Scherer, C.F.C., E.F. Flores, R. Weiblen, L. Caron, L.F. Irigoyen, J.P. Neves, and M.N. Maciel. 2001. Experimental infection of pregnant ewes with bovine viral diarrhea virus type-2 (BVDV-2): effects on the pregnancy and fetus. *Veterinary Microbiology* 79(4): 285-299.

Wu Q

Wu, Q., M.C. Brum, L. Caron, M. Koster, and M.J. Grubman. 2003. Adenovirus-mediated type I interferon expression delays and reduces disease signs in cattle challenged with foot-and-mouth disease virus. *Journal of Interferon and Cytokine Research* 23(7): 359-68.

Zsak L

Sur, J.H., T.G. Burrage, J.G. Neilan, D.L. Rock, and **L. Zsak**. 2001. African Swine Fever virus multigene family 360 and 530 genes affect virulence, pathogenesis and immune response in swine. American Society for Virology, p. 84.

Zsak, L., Z. Lu, T.G. Burrage, J.G. Neilan, G.F. Kutish, D.M. Moore, and D.L. Rock. 2001. Africa Swine Fever virus multigene family 360 and 530 genes are novel macrophage host range determinants. *Journal of Virology* 75: 3066-3076.

Zsak, L., J.H. Sur, T.G. Burrage, J.G. Neilan, and D.L. Rock. 2001. African Swine Fever virus multigene families 360 and 530 genes promote infected macrophage survival.

College Station, Texas

Southern Plains Agricultural Research Center

<http://sparc.usda.gov/>

Crippen TL

Crippen, T.L., K.M. Bischoff, V.K. Lowry, and M.H. Kogut. 2003. rP33 activates bacterial killing by chicken peripheral blood heterophils. *Journal of Food Protection* 66(5): 787-792.

Crippen, T.L., C.L. Sheffield, H. He, V.K. Lowry, and M.H. Kogut. 2003. Differential nitric oxide production by chicken immune cells. *Developmental and Comparative Immunology* 27(6-7): 603-610.

Farnell, M.B., **T.L. Crippen**, H. He, C.L. Swaggerty, and M.H. Kogut. 2003. Oxidative burst mediated by toll like receptors (TLR) and CD14 on avian heterophils stimulated with bacterial toll agonists. *Developmental and Comparative Immunology* 27(5): 423-429.

Ferro PJ

Peterson, M.J., R. Aguirre, **P.J. Ferro**, D.A. Jones, T.A. Lawyer, M.N. Peterson, and N.J. Silvy. 2002. Infectious disease survey of Rio Grande wild turkeys in the Edwards Plateau of Texas. *Journal of Wildlife Diseases* 38(4): 826-833.

Peterson, M.J., **P.J. Ferro**, M.N. Peterson, R.M. Sullivan, B.E. Toole, and N.J. Silvy. 2002. Infectious disease survey of lesser prairie chickens in north Texas. *Journal of Wildlife Diseases* 38(4): 834-839.

Swaggerty, C.L., I.Y. Pevzner, **P.J. Ferro**, T.L. Crippen, and M.H. Kogut. 2003. Association between in vitro heterophil function and the feathering gene in commercial broiler chickens. *Avian Pathology* 32(5): 483-488.

He H

Farnell, M.B., **H. He**, K. Genovese, and M.H. Kogut. 2003. Pharmacological analysis of signal transduction pathways required for oxidative burst in chicken heterophils stimulated by a Toll-like receptor 2 agonist. *International Immunopharmacology* 3(12): 1677-1684.

He, H., T.L. Crippen, M.B. Farnell, and M.H. Kogut. 2003. Identification of CpG oligodeoxynucleotide motifs that stimulate nitric oxide and cytokine production in avian macrophage and peripheral blood mononuclear cells. *Developmental and Comparative Immunology* 27(6-7): 621-627.

He, H., M.B. Farnell, and M.H. Kogut. 2003. Inflammatory agonist stimulation and signal pathway of oxidative burst in neonatal chicken heterophils. *Comparative Biochemistry and Physiology. Part-A, Molecular and Integrative Physiology* 135(1): 177-184.

He, H., M.H. Kogut. 2003. CpG-ODN-induced nitric oxide production is mediated through clathrin-dependent endocytosis, endosomal maturation, and activation of PKC, MEK1/2 and p38 MAPK, and NF-

kappaB pathways in avian macrophage cells (HD11). *Cell Signalling* 15(10): 911-917.

Kogut MH

Alfano, J.C., V.M. Petrone, T. Fehervari, G. Nava, **M.H. Kogut**, D. Nisbet, and G. Tellez. 2002. Resistance to velogenic Newcastle Disease virus in Leghorn chickens by use of prophylactic lymphokines. *Avian Diseases* 46: 525-534.

Farnell, M.B., T.L. Crippen, H. Haiqi, C.L. Swaggerty, and **M.H. Kogut**. 2003. Oxidative burst mediated by Toll-like receptors (TLR) and CD14 on avian heterophils stimulated with bacterial Toll agonists. *Developmental and Comparative Immunology* 27: 423-429.

Farnell, M.B., H. He, K. Genovese, and **M.H. Kogut**. 2003. Pharmacological analysis of signal transduction pathways required for oxidative burst in chicken heterophils stimulated by a Toll-like receptor-2 agonist. *International Immunopharmacology* 3: 1677-1684.

Farnell, M.B., **M.H. Kogut**, and H. He. 2003. Differential activation of signaling transduction pathways mediating oxidative burst by chicken heterophils in response to stimulation with lipopolysaccharide and lipoteichoic acid. *Inflammation* 27: 225-231.

Kogut, M.H. 2002. Dynamics of a protective avian inflammatory response: the role of an IL-8-like cytokine in the recruitment of heterophils to the site of organ invasion by *Salmonella enteritidis*. *Comparative Immunology, Microbiology, & Infectious Diseases* 25: 159-172.

Kogut, M.H. 2002. Immunomodulation of the avian innate immune response to

prevent bacterial infections in poultry. *Feedinfo.com* 1/29/2002.

Kogut, M.H. 2001. Signal transduction pathways activated by engaging immunoglobulin Fc receptors on chicken heterophils. *Developmental and Comparative Immunology* 25: 639-646.

Kogut, M.H., T.L. Crippen, V.K. Lowry, and M. Farnell. 2002. Selective pharmacological inhibitors reveal the role of Syk tyrosine kinase, phospholipase C, phosphatidylinositol 3'-kinase, and p38 mitogen-activated protein kinase in Fc receptor-mediated signaling of chicken heterophil degranulation. *International Immunopharmacology* 2: 963-973.

Kogut, M.H., M. Farnell, and V.K. Lowry. 2003. The use of selective pharmacological inhibitors to delineate signal transduction pathways activated during complement receptor-mediated degranulation in chicken heterophils. *International Immunopharmacology* 3: 693-706.

Kogut, M.H., K.J. Genovese, and V.K. Lowry. 2001. Differential activation of signal transduction pathways mediating phagocytosis, oxidative burst, and degranulation by chicken heterophils in response to stimulation with opsonized *Salmonella enteritidis*. *Inflammation* 25: 7-15.

Kogut, M.H., K.J. Genovese, and D.J. Nisbet. 2001. Signal transduction pathways activated by engaging immunoglobulin Fc receptors on chicken heterophils. *Developmental and Comparative Immunology* 25(7): 639-646.

Kogut, M.H., K.J. Genovese, and L.S. Stanker. 2001. Immune lymphokine-mediated control of salmonellosis in swine. U.S. Patent 6,221,348.

Kogut, M., V.K. Lowry, M. Farnell. 2002. Selective pharmacological inhibitors reveal the role of Syk tyrosine kinase, phospholipase C, phosphatidylinositol-3'-kinase, and p38 mitogen-activated protein kinase in Fc receptor-mediated signaling of chicken heterophil degranulation. *International Immunopharmacology* 2(7): 963-973.

Kogut, M.H., E.J. Pishko, B. Kaspers, and K.C. Weining. 2001. Modulation of functional activities of chicken heterophils by recombinant chicken IFN-gamma. *Journal of Interferon and Cytokine Research* 21: 85- 92.

Kogut, M.H., L. Rothwell, and P. Kaiser. 2003. Differential regulation of cytokine gene expression by avian heterophils during receptor-mediated phagocytosis of opsonized and nonopsonized *Salmonella enteritidis*. *Journal of Interferon and Cytokine Research* 23: 319-327.

Kogut, M.H., L. Rothwell, and P. Kaiser. 2003. Priming by recombinant chicken interleukin-2 induces selective expression of IL-8 and IL-18 mRNA in chicken heterophils during receptor-mediated phagocytosis of opsonized and nonopsonized *Salmonella enterica* serovar *Enteritidis*. *Molecular Immunology* 40: 603-610.

Kogut, M.H., L. Rothwell, and P. Kaiser. 2002. Differential effects of age on chicken heterophil functional activation by recombinant chicken interleukin-2. *Developmental and Comparative Immunology* 26: 817-830.

Lowry VK

Crippen, T.L., C.L. Sheffield, H. He, **V.K. Lowry**, and M.H. Kogut. 2003.

Differential nitric oxide production by chicken immune cells. *Developmental and Comparative Immunology* 27(6-7): 603-610.

Kogut, M., V.K. Lowry, and M. Farnell. 2002. Selective pharmacological inhibitors reveal the role of Syk tyrosine kinase, phospholipase C, phosphatidylinositol-3'-kinase, and p38 mitogen-activated protein kinase in Fc receptor-mediated signaling of chicken heterophil degranulation. *International Immunopharmacology* 2(7): 963-973.

Nisbet DJ

Kogut, M.H., K.J. Genovese, and D.J. Nisbet. 2001. Signal transduction pathways activated by engaging immunoglobulin Fc receptors on chicken heterophils. *Developmental and Comparative Immunology* 25(7): 639-646.

Nava M.G., **D.J. Nisbet**, V.K. Lowry, and I.G. Tellez. 2001. Effect of prophylactic administration of defined probiotic on mortality and horizontal transmission by *Salmonella gallinarum* in broiler chicks. *Fiftieth Proceedings of the Western Poultry Disease Conference, University of California at Davis*, p. 155-158.

Swaggerty CL

Swaggerty, C.L., I.Y. Pevzner, P.J. Ferro, T.L. Crippen, and M.H. Kogut. 2003. Association between in vitro heterophil function and the feathering gene in commercial broiler chickens. *Avian Pathology* 32(5): 483-488.

Swaggerty, C.L., I.Y. Pevzner, V.K. Lowry, M.B. Farnell, and M.H. Kogut. 2003. Functional comparison of heterophils isolated from commercial broiler chickens. *Avian Pathology* 32(1): 95-102.

Houston, Texas

Children's Nutrition Research Center

<http://www.bcm.tmc.edu/cnrc/index.html>

Smith C.W.

Anderson, D.C. and **C.W. Smith**. 2001. Leukocyte adhesion deficiencies. In C.R. Sriver, A.L. Beaudet, W.S. Sly, D. Valle, B. Childs, and B. Vogelstein, editors, *The Metabolic and Molecular Bases of Inherited Disease*, 8th Edition, Volume III, McGraw-Hill: New York, Chapter 188, pp. 4829-4856.

Burns, A.R., **C.W. Smith**, and D. Walker. 2003. Unique structural features that influence neutrophil emigration into the lung. *Physiological Reviews* 83: 309-336.

Chakraborty, A., E.R. Hentzen, S.M. Seo, and **C.W. Smith**. 2003. Granulocyte colony-stimulating factor promotes adhesion of neutrophils. *American Journal of Physiology: Cell Physiology* 284: C103-C110.

Collins, R.G., U. Jung, M. Ramirez, D.C. Bullard, M.J. Hicks, **C.W. Smith**, K. Ley, and A.L. Beaudet. 2001. Dermal and pulmonary inflammatory disease in E-selectin and P-selectin double-null mice is reduced in triple-selectin-null mice. *Blood* 98: 727-735.

Li, Z., C.A. Rivera, A.R. Burns, and **C.W. Smith**. 2004. Hindlimb unloading depresses corneal epithelial wound

healing in mice. *Journal of Applied Physiology* 97: 641-647.

Mariscalco, M.M., W. Vergara, J. Mei, E. Smith, and **C.W. Smith**. 2002. Mechanisms of decreased leukocyte localization in the developing host. *American Journal of Physiology: Heart and Circulatory Physiology* 282: H636-H644.

Prince, J.E., C.F. Brayton, M.C. Fossett, J.S. Durand, L. Kaplan, **C.W. Smith**, and C.M. Ballantyne. 2001. The differential roles of LFA-1 and Mac-1 in host defense against systemic infection with *Streptococcus pneumoniae*. *Journal of Immunology* 166: 7362-7369.

Rivera, C.A., M.H. Tchamitchi, L. Mendoza, and **C.W. Smith**. 2003. Endotoxemia and hepatic injury in a rodent model of hindlimb unloading. *Journal of Applied Physiology* 95: 1656-1663.

Robker, R.L., R.G. Collins, A.L. Beaudet, H.J. Mersmann, and **C.W. Smith**. 2004. Leukocyte migration in adipose tissue of mice null for ICAM-1 and Mac-1 adhesion receptors. *Obesity Research* 12(6): 936-940.

Seo, S.M., L.V. McIntire, and **C.W. Smith**. 2001. Effects of IL-8, Gro-alpha, and LTB4 on the adhesive kinetics of LFA-1 and Mac-1 on human neutrophils. *American Journal of Physiology: Cell Physiology* 281: C1568-C1578.

Logan, Utah

Poisonous Plant Research

<http://www.ars.usda.gov/pandp/locations/locations.htm?modecode=54-28-20-00>

Gardner DR

Panter, K.E., **D.R. Gardner**, L.F. James, B.L. Stegelmeier, and R.J. Molyneux. 2000. Natural toxins from poisonous plants affecting reproductive function in livestock. In A.T. Tu and W. Gaffield, editors, *Natural and Selected Synthetic Toxins: Biological Implications*, American Chemical Society: Washington, D.C., pp. 154-172.

Pfister, J.A., K.E. Panter, **D.R. Gardner**, B.L. Stegelmeier, R.J. Molyneux, M.H. Ralphs, and S.T. Lee. 2001. Alkaloids as anti-quality factors in plants on western U.S. rangelands. *Journal of Range Management* 54: 447-461.

James LF

Panter, K.E., D.R. Gardner, **L.F. James**, B.L. Stegelmeier, and R.J. Molyneux. 2000. Natural toxins from poisonous plants affecting reproductive function in livestock. In A.T. Tu and W. Gaffield, editors, *Natural and Selected Synthetic Toxins: Biological Implications*, American Chemical Society: Washington, D.C., pp. 154-172.

Ralphs, M.H., G. Greathouse, A.P. Knight, and **L.F. James**. 2001. Cattle preference for Lambert locoweed over white locoweed. *Journal of Range Management* 54: 265-268.

Panter KE

Panter, K.E. and B.L. Stegelmeier. 2000. Toxicoses of food animals.

Veterinary Clinics of North America: Food Animal Practice 6(3): 531-544.

Panter, K.E., D.R. Gardner, L.F. James, B.L. Stegelmeier, and R.J. Molyneux. 2000. Natural toxins from poisonous plants affecting reproductive function in livestock. In A.T. Tu and W. Gaffield, editors, *Natural and Selected Synthetic Toxins: Biological Implications*, American Chemical Society: Washington, D.C., pp. 154-172.

Pfister, J.A., **K.E. Panter**, D.R. Gardner, B.L. Stegelmeier, R.J. Molyneux, M.H. Ralphs, and S.T. Lee. 2001. Alkaloids as anti-quality factors in plants on western U.S. rangelands. *Journal of Range Management* 54: 447-461.

Pfister JA

Launchbaugh, K., F.D. Provenza, and **J.A. Pfister**. 2001. Animal response to anti-quality factors in forages. *Journal of Range Management* 54: 431-440.

Pfister, J.A., K.E. Panter, D.R. Gardner, B.L. Stegelmeier, R.J. Molyneux, M.H. Ralphs, and S.T. Lee. 2001. Alkaloids as anti-quality factors in plants on western U.S. rangelands. *Journal of Range Management* 54: 447-461.

Ralphs MH

Pfister, J.A., K.E. Panter, D.R. Gardner, B.L. Stegelmeier, R.J. Molyneux, **M.H. Ralphs**, and S.T. Lee. 2001. Alkaloids as anti-quality factors in plants on western U.S. rangelands. *Journal of Range Management* 54: 447-461.

Ralphs, M.H., G. Greathouse, A.P. Knight, and L.F. James. 2001. Cattle preference for

Lambert locoweed over white locoweed.
Journal of Range Management 54: 265-268.

Stegelmeier BL

Panther, K.E. and **B.L. Stegelmeier**.
2000. Toxicoses of food animals.
Veterinary Clinics of North America:
Food Animal Practice 6(3): 531-544.

Panther, K.E., D.R. Gardner, L.F. James,
B.L. Stegelmeier, and R.J. Molyneux.

Pullman, Washington

Animal Disease Research Unit

<http://pwa.ars.usda.gov/pullman/adru/web2.htm>

Goff WL

Boonchit, S., X. Xuan, N. Yokoyama,
W.L. Goff, S.D. Waghela, G. Wagner,
and I. Igarashi. 2004. Improved enzyme-
linked immunosorbent assay using C-
terminal truncated recombinant antigens
of *Babesia bovis* rhoptry-associated
protein 1 for detection of specific
antibodies. Journal of Clinical
Microbiology 42: 1601-1604.

Boonchit, S., X. Xuan, N. Yokoyama,
W.L. Goff, G. Wagner, and I. Igarashi.
2002. Evaluation of an enzyme-linked
immunosorbent assay with recombinant
rhoptry-associated protein-1 antigen
against *Babesia bovis* for the detection
of specific antibodies in cattle. Journal
of Clinical Microbiology 40: 3771-3775.

Davidson, W.R. and **W.L. Goff**. 2001.
Anaplasmosis. In E.S. Williams and I.K.
Barker, editors, Infectious Diseases of
Wild Mammals, Third Edition, Iowa
State University Press: Ames, Iowa,
chapter 27.

2000. Natural toxins from poisonous plants
affecting reproductive function in livestock.
In A.T. Tu and W. Gaffield, editors, Natural
and Selected Synthetic Toxins: Biological
Implications, American Chemical Society:
Washington, D.C., pp. 154-172.

Pfister, J.A., K.E. Panther, D.R. Gardner, **B.L. Stegelmeier**, R.J. Molyneux, M.H. Ralphs,
and S.T. Lee. 2001. Alkaloids as anti-quality
factors in plants on western U.S. rangelands.
Journal of Range Management 54: 447-461.

Goff, W.L., W.C. Johnson, R.H. Horn, G.M.
Barrington, and D.P. Knowles. 2003. The
innate immune response in calves to
Boophilus microplus tick transmitted *Babesia*
bovis involves type-1 cytokine induction and
NK-like cells in the spleen. Parasite
Immunology 25: 185-188.

Goff, W.L., W.C. Johnson, S.M. Parish,
G.M. Barrington, T.H. Elsasser, W.C. Davis,
and R.A. Valdez. 2002. IL-4 and IL-10
inhibition of IFN-gamma- and TNF-alpha-
dependent nitric oxide production from
bovine mononuclear phagocytes exposed to
Babesia bovis merozoites. Veterinary
Immunology and Immunopathology 84: 237-
251.

Goff, W.L., W.C. Johnson, S.M. Parish,
G.M. Barrington, W. Tuo, and R.A. Valdez.
2001. The age-related immunity in cattle to
Babesia bovis infection involves the rapid
induction of IL-12, IFN-gamma and iNOS
mRNA expression in the spleen. Parasite
Immunology 23: 453-461.

Goff, W.L., W.C. Johnson, W. Tuo, R.A. Valdez, S.M. Parish, G.M. Barrington, and W.C. Davis. 2002. Age-related innate immune response in calves to *Babesia bovis* involves IL-12 induction and IL-10 modulation. *Annals of the New York Academy of Science* 969: 164-168.

Goff, W.L., T.F. McElwain, C.E. Suarez, W.C. Johnson, W.C. Brown, J. Norimine, and D.P. Knowles. 2003. Competitive enzyme-linked immunosorbent assay based on a rhoptry-associated protein 1 epitope specifically identifies *Babesia bovis*-infected cattle. *Clinical and Diagnostic Laboratory Immunology* 10: 38-43.

Johnson, W.C., S. Lasri, L.S. Kappmeyer, A. Rhalem, H. Sahibi, D.P. Knowles, and **W.L. Goff**. 2001. Factors influencing the deployment of a *Babesia caballi* cELISA in tropical areas. *Proceedings of the 10th International Symposium of Veterinary Laboratory Diagnosticians*, p. 322-323.

Rhalem, A., H. Sahibi, S. Lasri, W.C. Johnson, L.S. Kappmeyer, A. Hamidouch, D.P. Knowles, and **W.L. Goff**. 2001. Validation of a competitive enzyme-linked immunosorbent assay for diagnosing *Babesia equi* infections of Moroccan origin and its use in determining the seroprevalence of *B. equi* in Morocco. *Journal of Veterinary Diagnostic Investigations* 13: 249-251.

Herrmann LM

Herrmann, L.M., T.V. Baszler, and D.P. Knowles. 2000. PrP(c) mRNA, but not PrP(Sc) is found in the salivary glands of scrapie-infected sheep. *Biochimica et Biophysica Acta* 1479: 147-154.

Herrmann, L.M., W.C. Davis, D.P. Knowles, K.J. Wardrop, M.S. Sy, P. Gambetti, and K.I. O'Rourke. 2001. Cellular prion protein is expressed on peripheral blood mononuclear cells but not platelets of normal and scrapie-infected sheep. *Haematologica* 86: 146-153.

Knowles DP Jr

Bradway, D.S., S. Torioni de Echaide, **D.P. Knowles**, S.G. Hennager, and T.F. McElwain. 2001. Sensitivity and specificity of the complement fixation test for detection of *Anaplasma marginale* persistently infected cattle. *Journal of Veterinary Diagnostic Investigation* 13: 77-79.

Brayton, K.A., **D.P. Knowles**, T.C. McGuire, and G.H. Palmer. 2001. Efficient use of small genome to generate antigenic diversity in tick-borne ehrlichial pathogens. *Proceedings National Academy Science* 98: 4130-4135.

Herrmann, L.M., T.V. Baszler, and **D.P. Knowles**. 2000. PrP(c) mRNA, but not PrP(Sc) is found in the salivary glands of scrapie-infected sheep. *Biochimica et Biophysica Acta* 1479: 147-154.

Herrmann, L.M., W.C. Davis, **D.P. Knowles**, K.J. Wardrop, M.S. Sy, P. Gambetti, and K.I. O'Rourke. 2001. Cellular prion protein is expressed on peripheral blood mononuclear cells but not platelets of normal and scrapie-infected sheep. *Haematologica* 86: 146-153.

Johnson, W.C., S. Lasri, L.S. Kappmeyer, A. Rhalem, H. Sahibi, **D.P. Knowles**, and W.L. Goff. 2001. Factors influencing the deployment of a *Babesia caballi* cELISA in tropical areas. *Proceedings of the 10th International Symposium of Veterinary Laboratory Diagnosticians*, p. 322-323.

Rhalem, A., H. Sahibi, S. Lasri, W.C. Johnson, L.S. Kappmeyer, A. Hamidouch, **D.P. Knowles**, and W.L. Goff. 2001.

Validation of a competitive enzyme-linked immunosorbent assay for diagnosing *Babesia equi* infections of Moroccan origin and its use in determining the seroprevalence of *B. equi* in Morocco. *Journal of Veterinary Diagnostic Investigations* 13: 249-251.

Shoda, L.K., K.A. Kegerreis, C.E. Suarez, W. Mwangi, **D.P. Knowles**, and W.C. Brown. 2001. Inclusion of a immunostimulatory CpG oligodeoxynucleotide sequence in plasmid DNA induced IL-12, TNF- α , and NO production by bovine monocyte-derived macrophages. *Journal of Leukocyte Biology* 70(1): 103-112.

Tuo, W., D. Zhuang, **D.P. Knowles**, W.P. Cheevers, M.S. Sy, and K.I. O'Rourke. 2001. Prp-c and Prp-Sc at the fetal-maternal interface. *Journal of Biological Chemistry* 276: 18229-18234.

Valdez, R.A., T.C. McGuire, W.C. Brown, W.C. Davis, and **D.P. Knowles**. 2001. Long term in vivo depletion of functional CD4⁺ T lymphocytes from calves requires both thymectomy and anti CD4 monoclonal antibody treatment. *Immunology* 102: 426-433.

O'Rourke KI

Hamir, A.N., R.C. Cutlip, J.M. Miller, E.S. Williams, M.J. Stack, M.W. Miller, **K.I. O'Rourke**, and M.J. Chaplin. 2001. Preliminary findings on the experimental transmission of chronic wasting disease agent of mule deer to cattle. *Journal of Veterinary Diagnostic Investigation* 13: 91-96.

Herrmann, L. M., W.C. Davis, D.P. Knowles, K.J. Wardrop, M.S. Sy, P. Gambetti, and **K.I. O'Rourke**. 2001. Cellular prion protein is expressed on peripheral blood mononuclear cells but not

platelets of normal and scrapie-infected sheep. *Haematologica* 86: 146-153.

Kim, H., **K.I. O'Rourke**, M. Walter, H.G. Purchase, J. Enck, and T.K. Shin. 2001. Immunohistochemical detection of scrapie prion proteins in clinically normal sheep in Pennsylvania. *Journal of Veterinary Diagnostic Investigation* 13: 89-91.

Koo, H.C., Y.H. Park, B.C. Lee, C. Chae, **K.I. O'Rourke**, and T.V. Baszler. 2001. Immunohistochemical detection of prion protein (PrP-Sc) and epidemiological study of BSE in Korea. *Journal of Veterinary Science* 2: 25-31.

Raymond, G.J., A. Bossers, L.D. Raymond, **K.I. O'Rourke**, L.E. McHolland, P.K. Bryant III, M.W. Miller, E.S. Williams, M. Smits, and B. Caughey. 2000. Evidence of a molecular barrier limiting susceptibility of humans, cattle and sheep to chronic wasting disease. *EMBO Journal* 19: 4425-4430.

Tuo, W., D. Zhuang, D.P. Knowles, W.P. Cheevers, M.S. Sy, and **K.I. O'Rourke**. 2001. Prp-c and Prp-Sc at the fetal-maternal interface. *Journal of Biological Chemistry* 276: 18229-18234.

Suarez CE

Camacho-Nuez, M., M.L. Munoz, **C.E. Suarez**, T.C. McGuire, W.C. Brown, and G.H. Palmer. 2000. Expression of polymorphic msp1b genes during acute *Anaplasma marginale* rickettsemia. *Infection and Immunity* 68(4): 1946-1952.

Florin-Christensen, J., **C.E. Suarez**, M. Florin-Christensen, S.A. Hines, T.F. McElwain, and G.H. Palmer. 2000. Phosphatidylcholine formation is the predominant lipid biosynthetic event in the hemoparasite *Babesia bovis*. *Molecular and Biochemical Parasitology* 106(1): 147-156.

Florin-Christensen, J., **C.E. Suarez**, M. Florin-Christensen, M. Wainszelbaum, W.C. Brown, T.F. McElwain, and G.H. Palmer. 2001. A unique phospholipid organization in bovine erythrocyte membranes. Proceedings of the National Academy of Sciences of the United States of America 98(14): 7736-7741.

Shoda, L.K., K.A. Kegerreis, C.E. Suarez, W. Mwangi, D.P. Knowles, and W.C. Brown. 2001. Inclusion of a immunostimulatory CpG oligodeoxynucleotide sequence in plasmid DNA induced IL-12, TNF- α , and NO production by bovine monocyte-derived macrophages. Journal of Leukocyte Biology 70(1): 103-112.

Shoda, L.K., K.A. Kegerreis, **C.E. Suarez**, I. Roditi, R.S. Corral, G.M. Bertot, J. Norimine, and W.C. Brown. 2001. DNA from protozoal parasites *Babesia bovis*, *Trypanosoma cruzi* and *T. brucei* is mitogenic for B lymphocytes and stimulates macrophage expression of interleukin-12, tumor necrosis factor α and nitric oxide. Infection and

Immunity 69(4): 2162-2171.

Suarez, C.E., M. Florin-Christensen, S.A. Hines, G.H. Palmer, W.C. Brown, and T.F. McElwain. 2000. Characterization of allelic variation in the *Babesia bovis* merozoite surface antigen-1 (msa-1) gene locus and identification of a cross-reactive, inhibition sensitive MSA-1 epitope. Infection and Immunity 68(12): 6865-6870.

Valdez RA

Goff, W.L., W.C. Johnson, S.M. Parish, G.M. Barrington, W. Tuo, and **R.A. Valdez**. 2001. The age-related immunity in cattle to *Babesia bovis* infection involves the rapid induction of IL-12, IFN- γ and iNOS mRNA expression in the spleen. Parasite Immunology 23: 453-461.

Valdez, R.A., T.C. McGuire, W.C. Brown, W.C. Davis, and D.P. Knowles. 2001. Long term in vivo depletion of functional CD4⁺ T lymphocytes from calves requires both thymectomy and anti CD4 monoclonal antibody treatment. Immunology 102: 426-433.

Kearneysville, West Virginia

National Center for Cool and Cold Water Aquaculture

<http://ncccwa.ars.usda.gov/>

Palti Y

Palti, Y., K.M. Nichols, K.I. Waller, J.E. Parsons, and G.H. Thorgaard. 2001. Association between DNA polymorphisms tightly linked to MHC class II genes and IHN virus resistance in backcrosses of rainbow and cutthroat trout. Aquaculture 194(3-4): 283-289.

Thorgaard, G.H., G.S. Bailey, D. Williams, D.R. Buhler, S.L. Kaattari, S.S. Ristow, J.D. Hansen, J.R. Winton, J.L. Bartholomew, J.J. Nagler, P.J.

Walsh, M.M. Vijayan, R.H. Devlin, R.W. Hardy, K.E. Overturf, W.P. Young, B.D. Robison, C. Rexroad, and **Y. Palti**. 2002. Status and opportunities for genomics research with rainbow trout. Comparative Biochemistry and Physiology. Part-B, Biochemistry and Molecular Biology 133(4): 609-646.

Wiens GD

Wiens, G.D., M. Brown, and M.B. Rittenberg. 2003. Repertoire shift in the humoral response to phosphocholine-keyhole

limpet hemocyanin: VH somatic mutation in germinal center B cells impairs T15 Ig function. *Journal of Immunology* 170(10): 5095-5102.

Wiens, G.D., R. Pascho, and J.R. Winton. 2002. A single Ala139-to-Glu

substitution in the *Renibacterium salmoninarum* virulence-associated protein p57 results in antigenic variation and is associated with enhanced p57 binding to chinook salmon leukocytes. *Applied and Environmental Microbiology* 68(8): 3969-3977.

Laramie, Wyoming

Arthropod-Borne Animal Diseases Research Laboratory

http://uwadmnweb.uwyo.edu/Usda_Ars/home.htm

Mecham JO

Gerry, A.C., B.A. Mullens, N.J. MacLachlan, and **J.O. Mecham**. 2001. Seasonal transmission of bluetongue virus by *Culicoides sonorensis* (Diptera: Ceratopogonidae) at a southern California dairy and evaluation of vectorial capacity as a predictor of bluetongue virus transmission. *Journal of Medical Entomology* 38(2): 197-209.

Mecham, J.O. 2001. Neutralization of bluetongue virus in invertebrate cells. American Society for Virology: Madison, Wisconsin, Abstract P23-1.

Mecham, J.O., et al. 2000. Report of the committee on bluetongue and bovine retroviruses. *Proceedings of the United States Animal Health Association* 104: 159-168.

Mecham, J.O. and M.M. Jochim. 2000. Development of an enzyme-linked immunosorbent assay for the detection of antibody to epizootic hemorrhagic disease of deer virus. *Journal Of Veterinary Diagnostic Investigation* 12: 142-145.

Mecham, J.O. and W.C. Wilson. 2004. Antigen capture competitive enzyme-linked immunosorbent assays using

baculovirus-expressed antigens for diagnosis of bluetongue virus and epizootic hemorrhagic disease virus. *Journal of Clinical Microbiology* 42(2): 518-523.

Mecham, J.O. and W.C. Wilson. 2001. Application of baculovirus expressed antigens in enzyme-linked immunosorbent assays for detection of antibody to bluetongue and epizootic hemorrhagic disease viruses. Abstracts from the scientific Sessions of the 44th American Association of Veterinary Laboratory Diagnosticians, Hershey, Pennsylvania, p. 20.

Meissner, J.D., **J.O. Mecham**, and W.C. Wilson. 2001. Verification of bluetongue virus S9 segment nucleotide sequences. *Virus Research* 81(1-2): 93-101.

Wilson, W.C. and **J.O. Mecham**. 2000. Molecular evolution of orbiviruses. *Proceedings of the United States Animal Health Association* 104: 169-180.

Wilson, W.C., H.C. Ma, E.H. Venter, A.A. van Djik, B.S. Seal, and **J.O. Mecham**. 2001. Erratum to 'Phylogenetic relationships of bluetongue viruses based on gene S7'. *Virus Research* 73: 201-202.

Wilson WC

Johnson, D.J., **W.C. Wilson**, and P.S. Paul. 2000. Validation of a reverse transcriptase multiplex PCR test for the serotype determination of U.S. isolates of bluetongue virus. *Veterinary Microbiology* 76: 105-115.

Meissner, J.D., J.O. Mecham, and **W.C. Wilson**. 2001. Verification of

bluetongue virus S9 segment nucleotide sequences. *Virus Research* 81(1-2): 93-101
Wilson, W.C., H.C. Ma, E.H. Venter, A.A. van Dijk, B.S. Seal, and J.O. Mecham. 2001. Erratum to 'Phylogenetic relationships of bluetongue viruses based on gene S7'. *Virus Research* 73: 201-202.

Wilson, W.C. and J.O. Mecham. 2000. Molecular evolution of orbiviruses. *Proceedings of the United States Animal Health Association* 104: 169-180.

APPENDIX

- List of Participants
- Author Index

List of Participants

Uma S. Babu

Food and Drug Administration
8301 Muirkirk Road
Laurel, MD 20708
301-827-8515
301-594-0517
usb@cfsan.fda.gov

Cynthia Baldwin

University of Massachusetts
410 Paige Laboratory
Amherst, MA 01003
413-545-3167
413-545-6326
cbaldwin@vasci.umass.edu

Douglas Bannerman

USDA, ARS, ANRI, BFGL
BARC – East
Building 1040, Room 2
Beltsville, MD 20705
301-504-5066
301-504-8330
dbanner@anri.barc.usda.gov

Rick Barb

USDA, ARS, SAA, Animal Physiology
Research Unit
Russell Research Center
P.O. Box 5677
Athens, GA 30604-5677
706-583-8276
706-542-0399
rbarb@saa.ars.usda.gov

José Barrera

USDA, ARS, Plum Island Animal Disease Ctr
P. O. Box 848
Greenport, NY 11944-0848
631-323-3266
631-323-3262

Elida M. Bautista

USDA, ARS, Plum Island Animal Disease Ctr
P. O. Box 848
Greenport, NY 11944-0848
631-323-3292
631-323-3295
ebautista@piadc.ars.usda.gov

Jay A. Berzofsky

Center for Cancer Research, National Cancer
Institute
National Institute of Health
Building 10, Room B12
Bethesda, MD 20892
301-496-6874
307-496-9956
berzofsky@helix.nih.gov

Joseph T. Bielitzki

Defense Advanced Research Projects
Agency/DSO
3701 North Fairfax Drive
Arlington, VA 22203-1714
703-696-5278
jbielitzki@darpa.mil

Lanie Bilodeau

USDA, ARS, Catfish Genetics Research Unit
141 Experiment Station Road
Stoneville, MS 38776
662-686-3591
662-686-3567
abilodeau@ars.usda.gov

Willi K. Born

National Jewish Medical & Research Center
1400 Jackson Street
Denver, CO 80220
303-398-1087
303-398-1396
birn@ujc.org

Sonia de Avila Botton

USDA, ARS, Plum Island Animal Disease
Center
31 Clearwater Road
Old Saybrook, CT 06475
631-323-3325
631-323-3006
sbotton@piadc.ars.usda.gov

David A. Brake

Pfizer Animal Health R&D
Pfizer Global Research and Development,
Pfizer, Inc.
M.S. 81180C2
Groton, CT 06340
860-441-8230
860-715-7936
david_brake@groton.pfizer.com

Deborah L. Brennan

USDA, ARS, MSA
P. O. Box 225
141 Experiment Station Road
Stoneville, MS 38776
662-686-5266
662-686-5459
dbrennan@ars.usda.gov

Robert E. Briggs

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7762
515-663-7458
bbriggs@nadc.ars.usda.gov

Kim A. Brogden

USDA, ARS, National Animal Disease Center
P. O. Box 70, 2300 Dayton Avenue
Ames, IA 50010
515-663-7980
515-663-7458
kbrogden@nadc.ars.usda.gov

Wendy C. Brown

Washington State University
Department of Veterinary Microbiology and
Pathology
Pullman, WA 99164-7040
509-335-6067
509-335-8529
wbrown@vetmed.wsu.edu

Mario Celso Sporotto Brum

USDA, ARS, Plum Island Animal Disease
Center
Foot and Mouth Disease Unit
P. O. Box 848
Greenport, NY 11922
631-323-3320
631-323-3006
mbrum@piadc.ars.usda.gov

John Butler

Department of Microbiology
The University of Iowa
51 Newton Road, 3550 Bowen Science Building
Iowa City, IA 52242
319-335-7776
john-butler@uiowa.edu

Luizinho Caron

USDA, ARS, PIADC, UNICAMP-IB-PPGBM
1146 Boston Post Road
Old Saybrook, CT 06475
631-323-3320
631-323-3355
lcaron@piadc.ars.usda.gov

Carol G. Chitko-McKown

USDA, ARS, U. S. Meat Animal Research
Center
P. O. Box 166, State Spur 18D
Clay Center, NE 68933-0166
402-762-4372
402-762-4375
mckown@email.marc.usda.gov

Michelle M. Colby

Office of Science and Technology Policy
438 Eisenhower Executive Office Building
Washington, DC 20502
202-456-6067
202-456-6027
mcolby@ostp.eop.gov

Darrell F. Cole

USDA, ARS, MWA
1815 N. University Street
Peoria, IL 61604
309-681-6600
309-681-6684
dfc@mwa.ars.usda.gov

Stephanie D. Collier

USDA, ARS, Poultry Research Unit
P. O. Box 5367
Mississippi State, MS 39762
662-320-7483
662-320-7589
scollier@msa-msstate.ars.usda.gov

Tawni Crippen

USDA, ARS, SPARC
2881 F&B Road
College Station, TX 77845
979-260-3773
979-260-9332

Rami A. Dalloul

USDA, ARS, ANRI, Animal Parasitic Diseases
Laboratory
Bldg 1043, Room 101, BARC-East
Beltsville, MD 20705
301-504-8771
301-504-5103
rdalloul@anri.barc.usda.gov

Bernadette Davillier

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2170D
Beltsville, MD 20705-5138
301-504-4698
301-504-5467

Mari Ellen Davis

University of Arkansas
Department of Animal Science, AFLS B-107
Fayetteville, AR 72701
479-575-5036
479-575-5756

Henry Dawson

USDA, ARS, Nutrient Requirements and
Functions Laboratory
10300 Baltimore Avenue
Bldg 307, Room 206, BARC-East
Beltsville, MD 20705
301-504-9412 ext. 278
301-504-9062
dawson@307.bhnrc.usda.gov

Annie De Groot

EpiVax, Inc.
16 Bassett Street
Providence, RI 02903
401-272-2123
401-272-7562
Anne_DeGroot@Brown.edu

Ann M. Donoghue

USDA, ARS, Poultry Production and Product
Safety Research
POSC 303 University of Arkansas
Fayetteville, AR 72701
479-575-2413
479-575-4202
donoghue@uark.edu

Susan Eicher

USDA, ARS, Livestock Behavior Research Unit
125 S. Russell Street, 216 Poultry, Purdue
University
Lafayette, IN 47907
765-496-3665
765-496-1993
spruiett@purdue.edu

Theodore Elsasser

USDA, ARS, ANRI, Growth Biology
Laboratory
Bldg 20, Room 201, BARC-East
Beltsville, MD 20705
301-504-8281
301-504-8623
elsasser@anri.barc.usda.gov

Gisela F. Erf

University of Arkansas
Department of Poultry
Fayetteville, AR 72701
479-575-8664
479-575-7139
gferf@uark.edu

Mark Estes

University of Texas Medical Branch
2-330F Children's Hospital - Department of
Pediatrics
Galveston, TX 77555-0732
409-772-0434
409-772-0460
dmestes@utamb.edu

Matthew J. Fenton

University of Maryland School of Medicine
685 W. Baltimore, MSTF-800
Baltimore, MD 21201-1192
410-706-0159
410-706-8162
mfenton@medicine.umaryland.edu

Brad Fenwick

USDA, CSREES
800 9th Street, SW
Waterfront Center
Washington, DC 20024
202-401-5022
202-401-2653
bfenwick@csrees@usda.gov

Geoffrey Ferman

USDA, Plum Island Animal Diseases Center
P. O. Box 848
Greenport, NY 11944-0848
631-323-3285
631-323-3295
gferman@piadc.ars.usda.gov

Pamela J. Ferro

USDA, ARS, SPARC
2881 F & B Road, Building 1
College Station, TX 77845
979-260-9525
979-260-9332
ferro@ffsru.tamu.edu

Janet Fulton

Hy-Line International
P. O. Box 310, 240th Street
Dallas Center, IA 50063
515-992-4173
jfulton@hyline.com

Louis Gasbarre

USDA, ARS, Bovine Functional Genomics
Laboratory
Bldg. 200, BARC-East, Room 6
Beltsville, MD 20705
301-504-8509, 301-504-6426
lgasbarr@anri.barc.usda.gov

Cyril Gay

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2194
Beltsville, MD 20705-5138
301-504-4786
301-504-5467
cgg@ars.usda.gov

Jesse P. Goff

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7547
515-663-7669
jgoff@nadc.ars.usda.gov

Will L. Goff

USDA, ARS, Animal Disease Research Unit
3003 ADBF, Washington State University
Pullman, WA 99164-6630
509-335-6003
509-335-8328
wgoff@vetmed.wsu.edu

William T. Golde

USDA, ARS, Plum Island Animal Diseases
Center
P. O. Box 848
Greenport, NY 11944-0848
631-323-3249
631-323-3005
wgolde@piadc.ars.usda.gov

Ronald D. Green

USDA, ARS, NPS, APPVS
GWCC, 5601 Sunnyside Avenue, Bldg 4-2164
Beltsville, MD 20705-5138
301-504-4736
rdg@ars.usda.gov

Adam Grow

USDA, APHIS, VS, NCAHP, EST
4700 River Road, Unit 43
Riverdale, MD 20737
301-736-3752
adam.g.grow@aphis.usda.gov

Andrew Hammond

USDA, ARS, SAA
P. O. Box 5677
950 College Station Road
Athens, GA 30604
706-546-3311
706-546-3398
ACH@saa.ars.usda.gov

James A. Harp

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7533
515-663-7669
jharp@nadc.ars.usda.gov

Gary J. Hausman

USDA, ARS, Russell Research Center
Animal Physiology Research Unit
950 College Station Road
Athens, GA 30605-2720
706-583-8275
706-542-0399
ghausman@saa.ars.usda.gov

Haiqi He

USDA, ARS
2881 F&B Road
College Station, TX 77845
979-260-3771
979-260-9332
he@ffsru.tamu.edu

Robert Heckert

USDA, ARS, NPS, APPVS
GWCC, 5601 Sunnyside Avenue, Bldg 4-2176
Beltsville, MD 20705-5138
301-504-5774
301-504-5467
rah@ars.usda.gov

Eliot Herman

USDA, ARS
Plant Genetic Research Unit, Donald Danforth
Plant Science Center
975 N. Warson Road
St. Louis, MO 63105
314-587-1292
314-587-1392
eherman@danforthcenter.org

Lynn M. Herrmann

USDA, ARS, Animal Disease Research Unit
Washington State University
ADBF, Room 3003
Pullman, WA 991640-6630
509-335-6068
509-335-8328
lherrman@vetmed.wsu.edu

Adrianna D. Hewings
USDA, ARS, MWA
1815 N. University Street
Peoria, IL 61604
309-681-6602
309-681-6684
Hewingsa@mwa.ars.usda.gov

Ronald L. Horst
USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7312
515-663-7669
rhorst@nadc.ars.usda.gov

Geraldine Huff
USDA, ARS, Poultry Production and Product
Safety Research
POSC 303 University of Arkansas
Fayetteville, AR 72701
479-575-7966
479-575-4202
grhuff@uark.edu

Henry D. Hunt
USDA, ARS, Avian Diseases and Oncology
Research
3606 E. Mount Hope Road
East Lansing, MI 48823-5338
517-337-6834
517-337-6776
hunthd@msu.edu

Mark Jenkins
USDA, ARS, APDL
BARC – East, Building 1040
Beltsville, MD 20705
301-504-8300
301-504-6273
mjenkins@anri.barc.usda.gov

Peter J. Johnson
USDA, CSREES
800 9th Street, S.W.
Washington, DC 20024
202-401-1896
pjohnson@csrees.usda.gov

Pete Kaiser
Institute for Animal Health
Compton, Berkshire RG20 7NN, UK
+44 1635 577277
pete.kaiser@bbsrc.ac.uk

Ildy M. Katona
Uniformed Services University of the Health
Sciences
Department of Pediatrics at USUHS
4301 Jones Bridge Road, Room A3007
Bethesda, MD 20814
301-295-3365
301-295-3898
dthomas@usuhs.mil

Marcus Kehrli
USDA, ARS, National Animal Disease Ctr
2300 Dayton Avenue
Ames, IA 50010
515-663-7254
mkehrli@nadc.ars.usda.gov

Kayoko Kimura
USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7419
515-663-7669

Kirk C. Klasing
University of California
Department of Animal Science
1 Shields Avenue
Davis, CA 95616
530-752-1901
530-752-0175
kcklasing@ucdavis.edu

Donald P. Knowles
USDA, ARS, Animal Disease Research Unit
Washington State University
ADBF, Room 3003
Pullman, WA 99164-6630
509-335-6022
509-335-8328
dknowles@vetmed.wsu.edu

Michael H. Kogut

USDA, ARS, Southern Plains Agricultural
Research Center
Food and Feed Safety Research
2881 F&B Road
College Station, TX 77845-9594
979-260-3772
979-260-9332
kogut@ffsru.tamu.edu

John Korslund

USDA, APHIS, VS, AHP, NCAAP, EST
4700 River Road
Riverdale, MD 20737
301-734-5914
301-734-7984
john.a.korslund@aphis.usda.gov

Tim R. Kramer

USDA, ARS
900 S. Shackleford Road
Suite 509
Little Rock, AR 72211
501-954-8102
501-954-9596
tkramer@spa.ars.usda.gov

Arthur M. Kreig

Coley Pharmaceutical Group
93 Worcester Street, Suite 101
Wellesley, MA 02481
781-431-9000 ext.1284
781-431-0951
akreig@coleypharma.com

William W. Laegreid

USDA, ARS, Meat Animal Research Center
P. O. Box 166, State Spur 18D
Clay Center, NE 68933-0166
402-762-4177
402-762-4375
laegreid@email.marc.usda.gov

Kelly M. Lager

USDA, ARS, National Animal Disease Center
P. O. Box 70
Ames, IA 50010
515-63-7371
515-663-7458
klager@nadcr.ars.usda.gov

Scott LaPatra

Clear Springs Food, Inc.
P.O. Box 712
Buhl, ID 83316
208-543-3456
208-543-4146
scottl@clearsprings.com

Jean Larson

USDA, Animal Welfare Information Center
10301 Baltimore Avenue
Beltsville, MD 20705
301-504-5215
301-504-7122
jlarson@nal.usda.gov

Beth Lautner

National Pork Board
1776 NW 114th Street
Clive, IA 50325
515-223-2623
515-223-2646
beth.lautner@porkboard.org

Robert Li

USDA, ARS, Catfish Genetics Research Unit
141 Experiment Station Road
Stoneville, MS 38776
662-686-3535
662-686-3567
rli@ars.usda.gov

Hyun Lillehoj

USDA, ARS, ANRI, APDL
Building 1040, BARC-East
Beltsville, MD 20705
301-504-8771
301-504-5103
hlilleho@anri.barc.usda.gov

James Lindsay

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2182
Beltsville, MD 20705-5138
301-504-4674
301-504-5467
jal@ars.usda.gov

John D. Lippolis

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7446
515-663-7669

Linda Logan-Henfrey

Pathobiology College of Veterinary Medicine
Mailstop 1961, Texas A&M University
College Station, TX 77843-1255
979-845-2660
979-862-3128
llogan@cvm.tamu.edu

Virginia K. Lowry

Texas A&M University
USDA, ARS, SPARC
2881 F & B Road
College Station, TX 77845
979-260-9263
979-260-9332
lowry@ffsru.tamu.edu

Joan K. Lunney

USDA, ARS, Animal Parasitic Diseases Lab.
10300 Baltimore Avenue
Bldg 1040, Rom 107, BARC-East
Beltsville, MD 20705
301-504-9368
301-504-5306
jlunney@anri.barc.usda.gov

Lolita Mangra

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2172A
Beltsville, MD 20705-5138
301-504-4778
301-504-5467
lsm@ars.usda.gov

William Martin

EpiVax, Inc.
16 Bassett Street
Providence, RI 02903
401-272-2123
401-272-5762
martinb@epivax.com

Wilda Martinez

USDA, ARS, NAA
600 East Mermaid Lane
Wyndmoor, PA 19038
215-233-6593
215-233-6719
wilda.martinez@ars.usda.gov

Charles V. Maxwell

University of Arkansas
Department of Animal Science, AFLS B-106
Fayetteville, AR 72701
479-575-2111
479-575-5756
cmaxwell@uark.edu

Gregory Mayer

USDA, APHIS, FADDL, DSS
P.O. Box 848
Greenport, NY 11944
631-323-3206
631-323-3366
gregory.a.mayer@aphis.usda.gov

Scott McVey

Pfizer Animal Health
601 W. Cornhusker Highway
Lincoln, NE 68527
402-441-2817
402-441-2760
d.scott.mcvey@pfizer.com

James O. Mecham

USDA, ARS, Arthropod-Borne Animal Diseases
Research Laboratory
P. O. Box 3965, University Station
Laramie, WY 82071
307-766-3620
307-766-3500
jmecham@uwyo.edu

Jeremy Merchant-Forde

USDA, ARS, LBRU
Poultry Science Building, Purdue University
125 S. Russell street
West Lafayette, IN 47907
705-494-6358
705-496-1993
merchant@perdue.edu

Simin Meydani

USDA, HNRC at Tufts University
711 Washington Street
Boston, MA 02111
617-556-3129
617-556-3224
simin.meydani@tufts.edu

Wongi Min

USDA, ARS, Animal Parasitic Diseases Lab.
Animal and Natural Resources Institute
Building 1040, BARC-East
Beltsville, MD 20705
301-504-8771
301-504-5103
wongimin@anri.barc.usda.gov

Albert Morgan

USDA, APHIS, VA
4700 River Road, Unit 148
Riverdale, MD 20737
301-734-8245
301-734-4314
albert.p.morgan@aphis.usda.gov

Brian Murphy

National Institutes of Health (NIH)
National Institute of Allergies and Infectious
Diseases (NIAID)
50 South Drive MSC 8007
Bethesda, MD 20892
301-594-1616
bm25f@nih.gov

Michael Murtaugh

Department of Veterinary Pathobiology
University of Minnesota
1971 Commonwealth Avenue
St. Paul, MN 55108
612-625-6735
612-625-5203
murta001@umn.edu

Peter L. Nara

Biological Mimetics, Inc.
124 Byte Drive
Frederick, MD 21702
301-620-7691
301-620-1827

John Neill

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7730
515-663-7458
jneill@nadc.ars.usda.gov

Eric Newmann

National Pork
1776 NW 114 th
Clive, IA 50325
515-889-2791
515-223-2646
eric.neuman@porkboard.org

Brian J. Nonnecke

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7311
515-663-7669

Robert Nordgren

R&D Merial, LTD
1210 Thornwell Drive
Athens, GA 30606

Evelyn A. Nystrom

USDA, ARS, National Animal Disease Ctr.
P. O. Box 70
Ames, IA 50010
515-663-7376
515-663-7458
enystrom@nadc.ars.usda.gov

Steven Olsen

USDA, ARS
141 Experiment
Stoneville, MS 38701
662-686-3546
662-686-3567
solsen@nadc.ars.usda.gov

Max Paape

USDA, ARS, NCCWA
BARC – East, Building 1040, Room 2
Beltsville, MD 20705
301-504-8302
301-504-8330
mpaape@anri.barc.usda.gov

Yniv Palti

USDA, ARS, National Center for Cool and
Cold Water Aquaculture
11876 Leetown Road
Kearneysville, WV 25430
304-724-8340
304-725-0351
ypalti@nccwa.ars.usda.gov

Sylvie Quiniou

USDA, ARS
141 Experiment
Stoneville, MS 38701
662-686-3546
662-686-3567
squiniou@msa-stoneville.ars.usda.gov

Muquarrab A. Qureshi

USDA, CSREES
1400 Independence Avenue S.W.
Washington, DC 20250-2220
202-401-4895
202-401-1602
mqureshi@csrees.usda.gov

Narayan Rath

USDA, ARS, Poultry Production and Product
Safety Research
POSC 303 University of Arkansas
Fayetteville, AR 72701
479-575-6189
479-575-4202
nrath@uark.edu

Richard B. Raybourne

Food and Drug Administration
8301 Muirkirk Road
Laurel, MD 20708
301-827-8099
301-594-0517
Richard.Raybourne@cfsan.fda.gov

Timothy A. Reinhardt

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7540
515-663-7669
treinhar@nadc.ars.usda.gov

Juergen A. Richt

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7366
515-663-7548
jricht@nadc.ars.usda.gov

Julia Ridpath

USDA, ARS, National Animal Disease Center
P.O. Box 70, 2300 Dayton Avenue
Ames, IA 50010
515-663-7586
515-663-7458
jridpath@nadc.ars.usda.gov

Jane Robens

USDA, ARS, NPS, APPVS
5601 Sunnyside Avenue, Bldg 4-2184
Beltsville, MD 20705-5138
301-504-5381
301-504-5467
jfr@ars.usda.gov

M. Fernanda Rodriguez

USDA, ARS, National Center for Cool and
Cold Water Aquaculture
11876 Leetown Road
Kearneysville, WV 25430
301-724-8340 ext 2184
frodriguez@nccwa.ars.usda.gov

Gary A. Rohrer

USDA, ARS, Meat Animal Research Center
Spur 18D, P. O. Box 166
Clay Center, NE 68933-0166
402-762-4365
402-762-4390
rohrer@email.marc.usda.gov

Ronald Rosenberg

USDA, ARS, NPS, APPVS
GWCC

5601 Sunnyside Avenue, Bldg 4-2178
Beltsville, MD 20705-5138

301-504-5771

301-504-5467

rmr@ars.usda.gov

James A. Roth

Iowa State University

VMPM, College of Veterinary Medicine

Ames, IA 50011

515-294-8459

515-294-8259

jaroeth@iastate.edu

Randy E. Sacco

USDA, ARS, National Animal Disease Center

2300 Dayton Avenue

Ames, IA 50010

515-663-7354

515-663-7458

rsacco@nadc.ars.usda.gov

Charles Fernando Capinos Scherer

USDA, ARS, PIADC, University of Missouri

97 High Street, Route 81

Clinton, CT 06413-1733

860-669-0137

csherer@piadc.ars.usda.gov

Ronald D. Schultz

University of Wisconsin-Madison

Department of Pathobiological Sciences, School
of Veterinary Medicine

2015 Linden Drive

Madison, WI 53706-1102

608-263-9888

608-262-1739

manningj@svm.vetmed.wisc.edu

Richard Shelby

USDA, ARS, Aquatic Animal Health

Laboratory

P. O. Box 952

Auburn, AL 36830

334-887-3741

334-887-2983

Franklin A. Sher

National Institutes of Health, National Institute
of Allergy and Infectious Diseases

50 South Drive, MSC-8002, Building. 50,
Room 6140,

Bethesda, MD 20892

301-496-3535

301-402-0890

asher@niaid.nih.gov

Lawrence Silbart

University of Connecticut

1390 Storrs Road, Unit 4163

AG Biotech Building

Storrs, CT

860-486-6073

860-486-5067

lsilbart@canr.unconn.edu

C. Wayne Smith

USDA, ARS, Baylor College of Medicine

Immunology & Medicine-Thrombosis Research,

Head, Section of Leukocyte Biology

1100 Bates, Room 6014

Houston, TX 77030-2600

713-798-0311

713-798-0337

cwsmith@bcm.tmc.edu

Douglas Smith

Baylor University Medical Center

3500 Gaston Avenue

Dallas, TX 75246

214-820-2119

214-820-6384

dsmith@baylorhealth.edu

Lewis Smith

USDA, ARS, NPS, APPVS

GWCC

5601 Sunnyside Avenue, Bldg 4-2168

Beltsville, MD 20705-5138

301-504-5925

301-504-5467

lws@ars.usda.gov

Wenxia Song

University of Maryland
2107 Microbiology Building
Dept. CBMG
College Park, MD 20742
301-405-7552
301-314-9489
ws90@umail.umd.edu

Joseph T. Spence

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2186
Beltsville, MD 20705-5138
301-504-7050
301-504-4725
jts@ars.usda.gov

Phillip J. Sprino

Colorado Serum Company
4950 York Street
Denver, CO 80216-0428
303-295-7527
303-295-1923
psprino@colorado-serum.com

Judith R. Stabel

USDA, ARS, National Animal Disease Center
2300 Dayton Road
Ames, IA 50010
515-663-7304
515-663-7458
jstabel@nadc.ars.usda.gov

Charles Stephensen

USDA, ARS, Western Human Nutrition
Research Center
Nutrition Department
1 Shields Avenue, University of California
Davis, CA 95616
530-754-9266
530-752-8966
cstephensen@ucdavis.edu

Donice Stewart

USDA, ARS, NPS, APPVS
GWCC
5601 Sunnyside Avenue, Bldg 4-2174
Beltsville, MD 20705-5138
301-504-4583
301-504-5467
dcs@ars.usda.gov

M. Suresh

Department of Pathobiological Sciences
School of Veterinary Medicine, University of
Wisconsin-Madison
2015 Linden Drive
Madison, WI 53706-1102
608-265-9791
608-263-0438
sureshm@svm.vetmed.wisc.edu

Christina Swaggerty

USDA, ARS, SPARC, FFSRU
2881 F & B Road
College Station, TX 77845
979-260-9397
979-260-9332
swaggerty@ffsru.tamu.edu

David E. Swayne

USDA, ARS, Southeast Poultry Research
Laboratory
RBRRRC, 950 College Station Road
Athens, GA 32604
706-546-3433
706-546-3161
dswayne@seprl.usda.gov

Wenbin Tuo

USDA, ARS, APDL, ANRI
Bldg 1040, Room 1, BARC-East
Beltsville, MD 20705
301-504-8258
wtuo@anri.barc.usda.gov

Joseph F. Urban

USDA, ARS, Nutrient Requirements and
Functions Laboratory
Bldg 307-C, Room 214, BARC-East
Beltsville, MD 20705
301-504-5528 ext. 267
301-504-9062
urban@307.bhnrc.usda.gov

Ray Waters

USDA, ARS, National Animal Disease Center
2300 Dayton Avenue
Ames, IA 50010
515-663-7756
515-663-7458
rwaters@nadc.ars.usda.gov

David White

USDA, ARS, ABADRL
Department 3965, 1000 E. University Avenue
Laramie, WY 82071
307-766-3601
307-766-3500
dmwhite@uwyo.edu

Ann L. Wiegers

USDA, APHIS, VS
510 South 17th Street, Suite 104
Ames, IA 50010
515-232-5785 ext.503
515-232-7120
ann.l.wiegers@aphis.usda.gov

Gregory Wiens

USDA, ARS, National Center for Cool and
Cold Water Aquaculture
11876 Leetown Road
Kearneysville, WV 25430
304-724-8340
304-725-0351
gwiens@ncccw.ars.usda.gov

David Woodland

Trudeau Institute
154 Algonquin Avenue
Soranac Lake, NY 12983
518-891-3080
518-891-5126
dwoodland@trudeauinstitute.org

Dayong Wu

USDA, ARS, Human Nutrition Research Center
on Aging at Tufts University
711 Washington Street
Boston, MA 02111
617-556-3368
617-556-3224
dayong.wu@tufts.edu

Max (Min-Tsung) Yeh

USDA, ARS, RVSS, FADDL, VS
P.O. Box 848
Greenport, NY 11944
631-323-3256
631-323-3366
max.yeh@aphis.usda.gov

Lijuan Yuan

The Ohio State University
FAHRP/OARDC 1680 Madison Avenue
Wooster, OH 44691
330-263-3740
330-264-3677
yuan.21@osu.edu

Xiaoping Zhu

University of Maryland at College Park
Lab of Immunology, VA-MD Regional College
of Veterinary Medicine
8075 Greenmead Drive
College Park, MD 20742
301-314-6814
301-314-6855
xz60@umail.umd.edu

Federico A. Zuckerman

University of Illinois
College of Veterinary Medicine
2001 South Lincoln Avenue
Urbana, IL 61802
217-333-7767
217-244-7421
fazaaa@uiuc.edu

Kurt Zuelke
USDA, ARS, ANRI, Biotechnology &
Germplasm Laboratory
Bldg 200, Room 124, BARC-East

Beltsville, MD 20705
301-504-5161
301-504-5123
kzuelke@anri.barc.usda.gov

Author Index

Abdel-Azim, GA	60	Cheng HH	21, 142
Adolfsson O	55	Cheng HW	35, 105
Allen PC	125	Cheng, H.	43
Andersen AA	106	Chitko-McKown CG	44, 149
Araujo R	28	Chung S	123
Araujo, R	61	Collier SD	38
Arns, CW	63	Collins, R.	58
Augustine PC	125	Crippen TL	35, 153
Babu, Uma	35	Dalloul R.	15, 54
Bacon LD	142	Dalloul, Rami	35
Badger TM	96	Davis, W.C.	39
Baldwin, C. L.	31	Dawson H	24, 27, 48, 52, 126
Balog JM	56, 93	Dawson, H	58
Bannantine JP	52	Dawson, H.	54
Bannerman DD	44, 125	De Groot, Annie [.....	21
Barrera J	38, 150	Dean-Nystrom EA	31, 113
Bautista EM	16, 63, 150	DeCunha, Cristina W.	39
Bautista, Elida M.	35	Dellagostin, O.A.	39
Baxt B	150	Donoghue AM	93
Belisle S	55	Dubey JP	18, 48, 126
Beshah E	24, 52	Dubey, JB	52
Beshah, E	58	Eiben, L.	43
Bilodeau AL	60, 147	Eicher SD	14, 35, 105
Bono JL	44	Eicher, S.D.	36
Born, Willi	17	Eiting, K.T.	44
Botton SA	63	Elsasser TH	127
Boyd P	18	Emara, M	61
Boyd, P.	52	Estes, D. Mark	19
Branton SL	147	Evans JD	127
Bricker BJ	106	Evans JJ	88
Briggs RE	106	Fach, SJ	39
Brockmeier SL	44, 107	Fadly AM	143
Brogden KA	107	Fayer R	127
Brown, Wendy C.	17	Ferman GS	16
Brum MC	16, 63, 150	Ferman, Douglas Gregg	35
Brum, Mario C.	35	Ferman, Geoffrey S.	35
Burton, JL	60	Ferro PJ	29, 56, 62, 153
Butte, N.	58	Ferro, PJ	61
Call J	24	Fetterer RH	18, 127
Call, J	58	Fetterer, R.	52
Caron L	63, 150	Foote, M.R.	41
Carroll JA	148	Foote, MR	58
Casey TA	64	Fowler, M.A.	41
Champagne ET	123	Fowler, MA	58
Cheevers, W.P.	43	Fox JM	44

Frank GH	108	Jenkins MC	128
Freeman, AE	60	Jiang X	58
Freytag T	58	Johnson WC	36
Gardner DR	157	Johnson, T. A.	36
Gasbarre LC	18, 28, 128	Johnson, T.R.	36
Gasbarre, LC	61	Kaiser P	37
Gasbarre. L.	52	Kaiser, P.	61
Gast RK	100	Kapczynski DR	64, 67, 101
Gause, W.	50	Kappmeyer LS	39
Gay RT	25, 55	Keen JE	44
Gay, R.T.	58	Kehrli ME Jr	46, 60, 110
Goff JP	40, 46	Kelley DS	98
Goff WL	36, 158	Kelm, SC	60
Goff, J. P.	19	Khalifeh MS	51
Goff, J.P.	40	Kim, JK	65
Golde WT	16, 38, 50, 63, 150	Kimura K	46, 110
Golde, William T.	35	Kimura, K.	19, 40
Gregg D	16, 50	King DJ	66, 101
Grosse WM	44	Klasing, Kirk C.	25
Grubman MJ	16, 63, 151	Klesius PH	88
Grubman, Marvin	35	Klinman, Dennis M.	35
Hakkak R	96	Knowles DP Jr	22, 36, 39, 159
Han SN	25, 55, 140	Knowles, D.P.	43
Han, SN	58	Kogut MH	29, 32, 35, 37, 56, 62, 154
Hansen, J.D.	43	Kogut, MH	61
Hare W	24	Koster MJ	63
Hare, W	58	Kuck, AL	60
Harlan, J.M.	44	Laegreid WW	44
Harp JA	40, 108	Lager KM	44, 110
He H	32, 35, 153	Ledbetter T	24
Heaton MP	44	Ledbetter, T	58
Heckert, Robert A.	35	Lee CW	66
Herman EM	149	Lee LF	145
Herrmann LM	22, 159	Lee, Jai Wei	37
Herrmann, L.M.	43	Lehmkuhl HD	111
Hines, SA	39	Lillehoj HS	15, 30, 32, 54, 61, 65, 129
Holt PS	40, 100	Lillehoj, HS	61, 65
Horst RL	109	Lillehoj, Hyun S.	35
Horton R	67	Lim CE	90
Hovatter K	42	Lippolis JD	46
Huff GR	56, 93	Lopez, A.M.	39
Huff WE	56, 94	Lowry VK	56, 155
Hunt HD	21, 144	Lunney JK	48, 135
Hunt, H.	43	Madden, K.	50
Hwang DH	97	Maleki SJ	57, 124
Jacob RA	97	Marko M	55
James A. Harp	64	Mason PW	151
James LF	157	Mathews B	52
Janutenaite, J.	44	Matthews BF	30

Matthews, BF.....	61	Reddy SM.....	146
Maue A.....	65	Register KB.....	115
McGuire, Travis C.....	39	Reinhardt TA.....	46, 115
McMunn KA.....	14	Rexroad C III.....	42
McMunn, K.A.....	36	Richt JA.....	115
Mecham JO.....	41, 162	Ridpath JF.....	47, 49, 116
Mengeling WL.....	111	Ristow, S.S.....	43
Mersmann, H.....	58	Robker, R.....	58
Meydani S.....	25, 53, 55, 141	Rock DL.....	151
Meydani, SN.....	58	Rodriguez LL.....	38, 50, 152
Miller JM.....	52, 112	Rothwell L.....	37
Miller LC.....	44	Rutherford, Mark.....	28
Miller, B.L.....	41	Sacco RE.....	39, 47, 117
Miller, BL.....	58	Scherer C.....	50, 152
Min W.....	30, 32, 61, 65	Schnell, S.....	60
Min, W.....	61, 65	Schoene N.....	24, 52
Moraes MP.....	63	Schoene, N.....	58
Morash, M.....	43	Schultz-Cherry S.....	66
Morimoto M.....	52	Seal BS.....	102
Morimoto, M.....	50	Shafer-Weaver KA.....	138
Murtaugh, Michael.....	28	Shea-Donohue T.....	24, 50, 52
Neilan JG.....	151	Shea-Donohue, T.....	58
Neill JD.....	47, 49, 112	Shelby R.A.....	91
Nielsen NC.....	105	Shoemaker CA.....	91
Nisbet DJ.....	155	Silva RF.....	146
Nishi SM.....	48	Small B.....	60
Noakes, M.....	43	Smith CW.....	58, 156
Nonnecke BJ.....	19, 26, 41, 67, 113	Solano-Aguilar GI.....	24, 52, 138
Nonnecke, BJ.....	58	Solano-Aguilar, G.....	58
Nonnecker, B.J.....	40	Sonstegard T.....	28
O'Rourke KI.....	160	Sonstegard, T.....	61
Oesper JC.....	64	Stabel JR.....	51, 52, 118
Olsen SC.....	65, 114	Stabel TJ.....	44, 118
Ordovas J.....	55	Staeheli, P.....	65
Paape MJ.....	37, 137	Stanton TB.....	118
Palmer MV.....	39, 52, 65, 67, 114	Stegelmeier BL.....	158
Palti Y.....	42, 43, 161	Stephensen CB.....	58, 98
Panter KE.....	157	Stoffegren WS.....	65
Perdue ML.....	102	Suarez CE.....	160
Pesch B.....	44	Suarez DL.....	66, 103
Pevzner, IY.....	61	Suresh, M.....	33
Pfister JA.....	157	Swaggerty CL.....	29, 62, 155
Phillips, R.....	43	Swaggerty, CL.....	61
Prior RL.....	96	Swayne DE.....	66, 67, 104
Quiniou SM.....	147	Tabatabai LB.....	119
Rahner TE.....	40	Thorgaard, G.H.....	43
Ralphs MH.....	157	Trout JM.....	139
Rath NC.....	48, 56, 95	Tumpey T.....	64, 66, 67, 104
Raybourne, Richard.....	35	Tuo W.....	139

Tuo, wenbin	52	Woodland, David L.	17
Tuo, Wenbin	18	Wu DY	25, 53, 142
Urban JF Jr.....	24, 48, 50, 52, 139	Wu Q.....	152
Urban, J.....	58	Wu, D.....	58
Valdez RA	161	Xiao, Zhengguo	28
Van Tassel, C.....	61	Xie, Hang	35
Van Tassell C.....	28	Xu D	92
Waldbieser GC	147	Yersin AG.....	122
Waters WR	26, 39, 41, 52, 65, 67, 119	Zarlenga DS.....	50, 52, 140
Waters, WR	58	Zhang H.	21
Welch T	42	Zhang, H.	43
Wesley RD.....	121	Zhao, A.	50
Whipple DL	67, 122	Zhao, Xin.....	37
Wiens GD	42, 161	Zimmerman, A.....	43
Wilson WC	163	Zsak L	152
Winn, R.K.....	44	Zuerner RL	122
Witter RL	146	Zunino SJ.....	99
Wolters W	60		

NATIONAL AGRICULTURAL LIBRARY



1022508831